

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annales@mbss.org, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / *INDICE GENERALE* / *CONTENTS*BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
*BIOTIC GLOBALIZATION***Okan AKYOL, Oğuzhan TAKICAK & Hasan TARUN**

A Fugitive Lessepsian Fish in a Sea-Cage Farm in the Aegean Sea: *Stephanolepis diaspros* (Monacanthidae) 1
Ubežni lesepski migrant iz ribogojnice v Egejskem morju: Stephanolepis diaspros (Monacanthidae)

Nicola BETTOSO, Lisa FARESI, Valentina TORBOLI & Jose A. CUESTA

Additional Record of the Pea Crab *Pinnotheres bicristatus* (Brachyura: Pinnotheridae) in the Adriatic Sea 7
Dodatni zapis o pojavljanju stražne rakovice vrste Pinnotheres bicristatus (Brachyura: Pinnotheridae) v Jadranskem morju

Alan DEIDUN, Sarah BAUMANN, Bruno ZAVA & Maria CORSINI-FOKA

Confirming the Occurrence of the Non-Indigenous *Pteragogus trispilus* (Actinopterygii: Labridae) within Maltese Waters 13
Potrditev pojavljanja tujerodne ustnače vrste Pteragogus trispilus (Actinopterygii: Labridae) znotraj malteških voda

Deniz ERGÜDEN, Yusuf Kenan BAYHAN, Sibel ALAGÖZ ERGÜDEN & Deniz AYAS

A New Ichthyological Record and Distributional Update for *Epigonus denticulatus* Dieuzeide, 1950 in Turkish Mediterranean Waters 21
Nov ihtiološki zapis in podatki o razširjenosti rjavega veleokca, Epigonus denticulatus Dieuzeide, 1950 v turških sredozemskih vodah

Sara LADOUL, Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On the Occurrence of Cornish Blackfish *Schedophilus medusophagus* (Osteichthyes: Centrolophidae) from the Maghreb Shore (Southwestern Mediterranean Sea) 27
Potrjena prisotnost meduzojeda Schedophilus medusophagus (Osteichthyes: Centrolophidae) z magrebske obale (jugozahodno Sredozemsko morje)

Christina MICHAIL & Francesco TIRALONGO

First Occurrence of Ariidae in Cypriot Waters – a Major Contribution to Biodiversity 35
Prvo pojavljanje predstavnikov iz družine Ariidae v ciprskih vodah – velik prispevek k biodiverziteti

SREDOZEMSKA HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Lovrenc LIPEJ, Riccardo BATTISTELLA, Borut MAVRIČ & Danijel IVAJNŠIČ**

An Insight into the Diet of the Bull Ray, *Aetomylaeus bovinus* (Geoffroy Saint-Hilaire, 1817) in the Northern Adriatic Sea 43
Vpogled v prehranjevalne navade kljunatega morskega goloba, Aetomylaeus bovinus (Geoffroy Saint-Hilaire, 1817) v severnem Jadranu

Cem ÇEVİK, Deniz ERGÜDEN & Deniz AYAS

A New Capture Record of *Alopias superciliosus* Lowe, 1841 from the Turkish Coast (Northeastern Mediterranean) 55
Nov ulov velikooke morske lisice Alopias superciliosus Lowe, 1841 iz turške obale (severovzhodno Sredozemlje)

Cem DALYAN, N. Bikem KESİCİ, Elif YÜCEDAĞ BAKIR, Yunus GÖNÜL & Hakan KABASAKAL

No Longer as Common as its Name: a Review of the Occurrence of *Torpedo torpedo* (Linnaeus, 1758) (Chondrichthyes: Elasmobranchii) in Turkish Waters, with Photographic Evidence 65
Ni več tako pogost kot njegovo ime: pregled pojavljanja okatega električnega skata Torpedo torpedo (Linnaeus, 1758) (Chondrichthyes: Elasmobranchii) v turških vodah s fotografskimi dokazi

Deniz ERGÜDEN, Cemal TURAN, Servet Ahmet DOĞDU & Deniz AYAS

Disc Deformity in a Juvenile Female Brown Ray, *Raja miraletus* (Family: Rajidae), from Northeastern Mediterranean (Türkiye) 73
Deformacija diska pri juvenilni samici modropege raže, Raja miraletus (družina: Rajidae), iz severovzhodnega Sredozemskega morja (Turčija)

Farid HEMIDA, Christian REYNAUD & Christian CAPAPÉ

On an Old Record of the Smalltooth Sand Tiger Shark *Odontaspis ferox* (Chondrichthyes: Odontaspidae) from the Algerian Coast (Southwestern Mediterranean Sea) 83
O starem zapisu o drobn zobem morskem biku Odontaspis ferox (Chondrichthyes: Odontaspidae) z alžirske obale (jugozahodno Sredozemsko morje)

Hakan KABASAKAL, Uğur UZER & F. Saadet KARAKULAK

Plastic Debris-Induced Fin Damage in the Smoothhound, *Mustelus mustelus* 91
Poškodbe plavuti pri navadnem morskem psu, Mustelus mustelus, zaradi plastičnih odpadkov

Nicolas ZIANI, Florane TONDU, Rémi BRU, Chloé MOSNIER, Sarah FOXONET, Ruben Bao GALLIEN, Mathias POULY, Modan Lou TONIETTO, Lucille VERDON, Eloïse DEYSSON, Alessandro DE MADDALENA & Hakan KABASAKAL

Bite Marks Observed on a Large Female White Shark *Carcharodon carcharias* Off Camargue, France Provide Potential Insights into the Reproduction of the Mediterranean Population 97
Sledovi ugrizov na veliki samici belega morskega volka Carcharodon carcharias pri Camargu (Francija) kažejo na možno razmnoževanje sredozemske populacije

**MORSKA FAVNA
FAUNA MARINA
MARINE FAUNA****Sihem RAFRAFI-NOUIRA, RIMEL BENMESSAOUD, Mourad CHÉRIFF, Christian REYNAUD & Christian CAPAPÉ**

Morphological Deformities in a Common Two-Banded Sea Bream, *Diplodus vulgaris* (Osteichthyes: Sparidae), from Northern Tunisian Waters (Central Mediterranean Sea) 109
Morfološke deformacije pri fratru, Diplodus vulgaris (Osteichthyes: Sparidae), iz severnih tunizijskih vod (osrednje Sredozemsko morje)

Abdelkarim DERBALI, Aymen HADJ TAIEB & Wassim KAMMOUN

The Current Status of *Polititapes aureus* (Mollusca: Bivalvia) in the Coastal Zone of Sfax, Tunisia (Central Mediterranean) 117
Trenutno stanje vrste Polititapes aureus (Mollusca: Bivalvia) na obalnem območju Sfaksa v Tuniziji (osrednje Sredozemlje)

Neža LEBAN & Valentina PITACCO

Current Knowledge on the Distribution of the Poorly Known Echiurid Species *Maxmuelleria gigas* (M. Müller, 1852) in the Slovenian Sea 125
Trenutno poznavanje prostorske razporeditve manj poznane vrste zvezdaša Maxmuelleria gigas (M. Müller, 1852) v slovenskem morju

Jan MALEJ, Tjaša KOGOVŠEK, Martin VODOPIVEC, Janja FRANCÉ, Patricija MOZETIČ, Matevž MALEJ & Alenka MALEJ

Long-Term Study of Zooplankton Biomass in the Gulf of Trieste (Adriatic Sea) 133
Dolgoročna študija zooplanktonske biomase v Tržaškem zalivu (Jadransko morje)

Sihem RAFRAFI-NOUIRA, Rimel BENMESSAOUD, Mourad CHÉRIFF, Christian REYNAUD & Christian CAPAPÉ

Occurrence of the Longjaw Snake Eel, *Ophisurus serpens* (Ophichthidae), in Tunisian Waters (Central Mediterranean Sea) 145
Pojavljanje zobate jegulje, Ophisurus serpens (Ophichthidae), iz tunizijskih voda (osrednje Sredozemsko morje)

FLORA
FLORA
FLORA

**Martina ORLANDO-BONACA, Artur BONACA,
Diego BONACA & Ana ROTTER**

Seagrasses: a Promising Source of
Bioactive Compounds for Human
Health Applications 153
*Morske cvetnice: obetaven vir
bioaktivnih spojin za uporabo
v zdravstvu*

OCENE IN POROČILA
RECENSIONI E RELAZIONI
REVIEWS AND REPORTS

Shin-ichi Uye

Book review: Mirrors of the Sea: When
Science and Art Meet. 30 Years of the
Unesco Intergovernmental Oceanographic
Commission in Slovenia 167
Kazalo k slikam na ovitku 169
Index to images on the cover 169

received: 2025-02-14

DOI 10.19233/ASHN.2025.08

A NEW CAPTURE RECORD OF *ALOPIAS SUPERCILIOSUS* LOWE, 1841 FROM THE TURKISH COAST (NORTHEASTERN MEDITERRANEAN)

Cem ÇEVİK

Faculty of Fisheries, Çukurova University Balcalı Campus, 01330, Sarıçam/Adana, Türkiye

Deniz ERGÜDEN

Marine Science Department, Faculty of Marine Science and Technology, Iskenderun Technical University, 31200 Iskenderun, Hatay, Türkiye

e-mail: deniz.erguden@iste.edu.tr; derguden@gmail.com

Deniz AYAS

Faculty of Fisheries, Mersin University Yenişehir Campus, 33160, Mersin, Türkiye

ABSTRACT

A single male bigeye thresher, Alopias superciliosus, was incidentally captured by a commercial trawler. The specimen measured 310 cm in total length and weighed 156.0 kg. The capture occurred on 16 January 2025 in the coastal region of Akyatan, at a depth of 30 m. Prior to this study, 15 records of A. superciliosus had been documented along the northeastern Mediterranean coast of Türkiye. This study contributes to the existing body of knowledge and marks a notable milestone by documenting the first recorded occurrence of an adult male bigeye thresher in the northeastern Mediterranean, specifically along the Akyatan coast of Türkiye. It also presents a comprehensive analysis of the specimen's morphometric measurements in comparison with existing Mediterranean records.

Key words: bigeye thresher, Alopiidae, record, Akyatan coast, Türkiye

NUOVA CATTURA DI *ALOPIAS SUPERCILIOSUS* LOWE, 1841 LUNGO LA COSTA TURCA (MEDITERRANEO NORD-ORIENTALE)

SINTESI

Un singolo esemplare maschio di Alopias superciliosus è stato catturato accidentalmente da un peschereccio commerciale. L'esemplare misurava 310 cm di lunghezza totale e pesava 156,0 kg. La cattura è avvenuta il 16 gennaio 2025 nella regione costiera di Akyatan, a una profondità di 30 m. Prima d'ora erano stati documentati 15 esemplari di A. superciliosus lungo la costa mediterranea nord-orientale della Turchia. Questo studio contribuisce alle conoscenze esistenti e segna una pietra miliare notevole, documentando la prima presenza registrata di un maschio adulto di squalo volpe occhione nel Mediterraneo nord-orientale, in particolare lungo la costa di Akyatan in Turchia. Viene inoltre presentata un'analisi completa delle misure morfometriche dell'esemplare rispetto ai dati esistenti nel Mediterraneo.

Parole chiave: squalo volpe occhione, Alopiidae, cattura, costa di Akyatan, Turchia

INTRODUCTION

The family Alopiidae comprises three globally distributed species: the bigeye thresher (*Alopias superciliosus*), the common thresher (*Alopias vulpinus*), and the pelagic thresher (*Alopias pelagicus*) (Ebert *et al.*, 2021; Froese & Pauly, 2025). Of these shark species, *A. superciliosus* and *A. vulpinus* have been recorded in the Mediterranean Sea and along the Turkish coastline, while *A. pelagicus* is distributed in the Indo-Pacific Ocean (Ebert *et al.*, 2021).



Fig. 1: Map showing the capture site (•) of the *A. superciliosus* along the Akyatan coast, Türkiye.

Sl. 1: Zemljevid obravnavanega območja z lokaliteto ulova (•) primerka vrste *A. superciliosus* vzdolž obale Akyatan, Turčija.

The bigeye thresher, *A. superciliosus*, is a large shark from the order Lamniformes. It is highly migratory and widely distributed in warm temperate and tropical waters (Compagno, 2001). This species occurs both in the pelagic and benthopelagic zones of the continental shelf at depths ranging from 0 to 955 m, with a higher prevalence in the 0–100 m range (Compagno, 2001; Coelho *et al.*, 2015). *A. superciliosus* primarily feeds on teleost fishes (lancetfish, herring, mackerel, etc.) and small billfishes in the pelagic zone, as well as on bottom fish such as the European hake and squid in the benthic zone (Compagno, 1984).

In the Mediterranean Sea, the bigeye thresher shark was first reported from Italian waters (Cigala-Fulgosi, 1983), with subsequent records extending from the west to the east of the Basin (Corsini-Foka & Sioulas, 2009; Kabasakal *et al.*, 2011; Farrag *et al.*, 2017; Kabasakal *et al.*, 2017; Kleitou *et al.*, 2017; Lanteri *et al.*, 2017; Ergüden *et al.*, 2020).

The first documented sighting of the bigeye thresher in Turkish waters occurred along the Marmaris and Gökova coasts in 2005 (Mater, 2005; Clo *et al.*, 2008), followed by a recorded capture in Silivri (Kabasakal & Karhan, 2008). Subsequent reports include records from Silivri, Sivrice, and Fethiye in 2011 and 2016 (Kabasakal *et al.*, 2011), from the Gulf of Antalya in 2015 and 2016 (Gökoğlu *et al.*, 2017; Kabasakal, 2017; Lanteri *et al.*, 2017), as well as from Tasucu and off Fethiye (Ayas *et al.*, 2020; Akyol *et al.*, 2020). The most recent documented capture in Turkish waters was an adult female incidentally caught off the coast of Çevlik in Iskenderun Bay (Ergüden *et al.*, 2020).

The present paper reports the first incidental capture of an adult male bigeye thresher, *A. superciliosus*, along the Akyatan coast (northeastern Mediterranean, Türkiye) and the morphometric measurements of the specimen.

MATERIAL AND METHODS

The bigeye thresher was captured at night by a commercial trawler at a depth of 30 m on the sandy-muddy bottom of the Akyatan coast (at



Fig. 2: The *A. superciliosus* recorded along the Akyatan coast, northeastern Mediterranean Sea (scale bar: 0.3 cm).
Sl. 2: Primerek vrste *A. superciliosus*, ujet vzdolž obale Akyatan, severovzhodno Sredozemsko morje (merilo: 0,3 cm).

Tab. 1: Comparison of selected morphometric measurements of the captured *A. superciliosus* with six previously reported specimens from the Mediterranean Sea. Abbreviations; TL (total length), FDFL (first dorsal fin length), FDFBL (first dorsal fin base length), PFL (pectoral fin length), PFBL (pectoral fin base length), VFL (ventral fin length), VFBL (ventral fin base length), DFVF (distance between dorsal-fin origin and ventral fin origin), PFVF (distance between pectoral fin origin and ventral fin origin), PDL (predorsal length), PVL (preventral length), PPL (pre pectoral length), IOS (Inter orbital space), EY (Eye length), EH (Eye height), MW (mouth width, ML (Mouth length), TailL (tail length), CL (Clasper length). *Lanteri et al. (2017) did not provide percentage TL values in their records on the collected head and pectoral fins of the studied specimen.

Tab. 1: Primerjava izbranih morfometričnih meritev ujetega primerka vrste *A. superciliosus* s šestimi predhodno ugotovljenimi primerki iz Sredozemskega morja. Okrajšave; TL (totalna dolžina), FDFL (dolžina prve hrbtnje plavuti), FDFBL (dolžina korena prve hrbtnje plavuti), PFL (dolžina prsne plavuti), PFBL (dolžina korena prsne plavuti), VFL (dolžina trebušne plavuti), VFBL (dolžina osnove trebušne plavuti), DFVF (razdalja med začetkom hrbtnje plavuti in začetkom trebušne plavuti), PFVF (razdalja med začetkom prsne plavuti in začetkom trebušne plavuti), PDL (predorzalna dolžina), PVL (preventralna dolžina), PPL (prepektoralna dolžina), IOS (medočesni prostor), EY (dolžina oči), EH (višina oči), MW (širina ust), ML (dolžina ust), TailL (dolžina repa), CL (dolžina klasperja). *Lanteri in sod. (2017) v svojih zapisih niso navedli vrednosti TL v odstotkih za glavo in prsne plavuti proučevanega osebk.

Character (cm)	This study	Corsini-Foka & Sioulas (2009)	Kabasakal et al. (2011)	*Lanteri et al. (2017)	Farrag et al. (2017)	Ayas et al. (2020)	Ergüden et al. (2020)
Location	Karataş coast	Dodocenasae waters	Fethiye coast	Ligurian Sea	Egyptian coast	Tasucu coast	Çevlik coast
Sex	Male	Male	Female	Female	-	Female	Female
TL	310.0	310.0	450.0	-	180.0	240.0	472.0
FDFL	25.0 (8.06 %TL)	27.0 (8.7 %TL)	41.0 (9.1 %TL)	-	-	17.9 (7.4 %TL)	40.0 (8.5 %TL)
FDFBL	22.0 (7.9 %TL)	21.0 (6.8 %TL)	32.0 (7.1 %TL)	-	-	13.6 (5.6 %TL)	32.0 (6.8 %TL)
PFL	60.0 (19.3 %TL)	62.0 (20.0 %TL)	82.0 (18.2 %TL)	25.4	35.0 (19.4 %TL)	51.5 (21.4 %TL)	83.0 (17.5 %TL)
PFBL	25.0 (8.06 %TL)	21.0 (6.8 %TL)	30.0 (6.7 %TL)	-	-	20.1 (8.4 %TL)	40.0 (8.5 %TL)
VFL	26.0 (8.04 %TL)	26.0 (8.4 %TL)	39.0 (8.7 %TL)	-	-	20.6 (8.6 %TL)	37.0 (7.8 %TL)
VFBL	25.0 (8.06 %TL)	21.0 (6.8 %TL)	30.0 (6.7 %TL)	-	-	15.4 (6.4 %TL)	30.0 (6.4 %TL)
DFVF	36.0 (11.6 %TL)	23.0 (7.4 %TL)	-	-	-	17.8 (7.4 %TL)	-
PFVF	57.0 (18.4 %TL)	77.0 (24.8 %TL)	-	-	-	68.9 (28.7 %TL)	204.0 (43.2 %TL)
PDL	97.0 (31.3 %TL)	94.0 (30.3 %TL)	128.0 (28.4 %TL)	-	57.0 (31.6 %TL)	87.7 (36.5 %TL)	-
PVL	124.0 (40.0 %TL)	118.0 (38.1 %TL)	165.0 (36.7 %TL)	-	70.0 (38.8 %TL)	105.3 (43.9 %TL)	-
PPL	52.0 (16.7 %TL)	45.0 (14.5 %TL)	55.0 (12.2 %TL)	47.7	29.0 (16.1 %TL)	43.7 (18.2 %TL)	-
IOS	13.1 (4.2 %TL)	-	-	7.4	-	-	-
EL	7.6 (2.4 %TL)	-	-	7.4	-	-	-
EH	5.4 (1.7 %TL)	-	-	4.7	-	-	-
MW	11.0 (3.5 %TL)	-	-	10.4	-	-	20.0 (4.2 %TL)
ML	8.0 (2.6 %TL)	-	-	7.3	-	-	-
TailL	160.0 (51.6 %TL)	143.0 (46.1 %TL)	217.0 (48.2 %TL)	54.4	-	103.8 (43.3 %TL)	240.0 (50.5 %TL)
CL	30.2	-	-	-	-	-	-

36°28'20.52" N, 34°55'20.99" E) on 16 January 2025 (Fig. 1). Upon being brought aboard, the specimen was found deceased. Since the capture occurred nocturnally, the individual was stored in a refrigerated vessel by the crew until its delivery to the laboratory the following day. Upon arrival at the laboratory, the specimen was immediately examined and identified. It was then preserved at the Museum of the Faculty of Fisheries, Cukurova University, registered under catalogue number CSFM-PIS/18.01.2025 (see Figs. 2 and 3).

The morphometric measurements of the captured *A. superciliosus* were recorded to the nearest 0.01 mm using a calliper. All subsequent morphological descriptions and coloration of the specimen were consistent with those previously documented for this species by Gruber & Compagno (1981) and Compagno (2001).

Nineteen morphometric measurements were taken, including total length (TL), first dorsal fin length (FDFL), first dorsal fin base length (FDFBL), pectoral fin length (PFL), pectoral fin base length (PFBL), ventral fin length (VFL), ventral fin base length (VFBL), distance between dorsal fin origin and ventral fin origin (DFVF), distance between pectoral fin origin and ventral fin origin (PFVF), predorsal length (PDL), preventral length (PVL),

prepectoral length (PPL), interorbital space (IOS), eye length (EYL), eye height (EYH), mouth width (MOW), mouth length (MOL), tail length (TAIL), and clasper length (CL). The complete list of morphometric measurements can be found in Table 1.

RESULTS AND DISCUSSION

The male bigeye thresher specimen measured 310 cm in total length (TL) and weighed 156.0 kg. Several morphometric measurements (expressed as percentages of TL) were recorded as follows: first dorsal fin length (FDFL), 8.06%; first dorsal fin base length (FDFBL), 7.90%; pectoral fin length (PFL), 19.30%; pectoral fin base length (PFBL), 8.06%; ventral fin length (VFL), 8.04%; ventral fin base length (VFBL), 8.06%; distance between dorsal fin origin and ventral fin origin (DFVF), 11.60%; distance between pectoral fin origin and ventral fin origin (PFVF), 18.40%; predorsal length (PDL), 31.3%; preventral length (PVL), 40.00%; prepectoral length (PPL), 16.70%; and tail length (TAIL), 16.00% (Tab. 1).

A comparison of morphometric measurements was made between the male specimen from this study and available Mediterranean records of both male and female bigeye thresher specimens (Tab. 1). Table



Fig. 3: Head view of the captured *A. superciliosus*.
Sl. 3: Glava ujetega primerka vrste *A. superciliosus*.

Tab. 2: Historical capture records of *A. superciliosus* in Turkish Mediterranean waters.**Tab. 2: Zgodovinski podatki o pojavljanju vrste *A. superciliosus* v sredozemskih vodah Turčije.**

References	Locality	Date	Number of Specimen	Depth	Fishing Gear	Sex	Length (cm) /Weight (kg)
Mater (2005)	Gökova	23 May 2005	1	12	Shrimp net	-	350/150.0
Clo <i>et al.</i> (2009)	Marmaris	April 2004	1	-	Gill net	-	-/160.0
Kabasakal & Karhan (2008)	Silivri	23 February 2007	1	-	Purse-seine	-	450/-
Kabasakal <i>et al.</i> (2011)	Fethiye	28 February 2011	1	110	Trammel net	Female	430/300.0
Kabasakal <i>et al.</i> (2011)	Silivri	02 July 2011	1	-	Purse-seine	-	250/65.0
Kabasakal <i>et al.</i> (2011)	Sivrice	21 May 2016	1	100	Stationary net	-	400/-
Gökoglu <i>et al.</i> (2017)	Antalya	19 March 2015-15 July 2015	2	600-700	Bottom trawl-	Female	180-299/19.0-65.0
		19 March 2015-12 April 2015	2		Long-line	Male	180-293/15.5-57.0
Kabasakal (2017)	Antalya	22 December 2017	1		Pelagic swordfish longline	Female	342.4/-
Lanteri <i>et al.</i> (2017)	-	25 February 2013	1	-	-	-	370/250.0
Lanteri <i>et al.</i> (2017)	Yeşilköy	13 March 2013	1	-	-	-	200/110.0
Lanteri <i>et al.</i> (2017)	Yediburunlar	15 July 2013	1	-	-	-	500/395.0
Lanteri <i>et al.</i> (2017)	Antalya	15 April 2015	1	-	-	-	-
Akyol <i>et al.</i> (2020)	off Fethiye	23 September 2012	1	1100	Pelagic longline	Female	150/30.2
Ergüden <i>et al.</i> (2020)	Çevlik	9 April 2019	1	20	Purse seine	Female	472/400.0
Ayas <i>et al.</i> (2020)	Taşucu	2 January 2020	1	25	Trammel net	Female	240/48.0
This study	Akyatan	16 January 2025	1	30	Bottom trawl	Male	310/156.0

2 presents detailed capture data for the incidentally caught specimen, including: locality, capture date, depth, region, fishing gear, sex, length, and weight, along with comparable data from previous records in Mediterranean waters.

The following characteristics are diagnostic for this shark species: head flattened dorsally and ventrally; lateral grooves situated above the branchial region; snout bulbous; eyes large and extending into the dorsal part of the head; first dorsal fin positioned more posteriorly, with the midpoint of its origin markedly closer to pelvic fin origin; dorsal extension of caudal fin approximates precaudal length; pectoral fins notably pointed; upper lobe of the caudal fin characterised by elongated, strap-like appearance nearly equalling the length of the remainder of the body; lower lobe comparatively short but well-developed (Compagno, 1998).

In terms of coloration, the dorsolateral region of the specimen exhibited a brownish-grey hue and the ventral side a whitish tone with greyish undertones. The posterior margins of the pectoral, pelvic, and first dorsal fin showed dusky pigmentation.

The bigeye thresher is readily distinguishable from the other two thresher shark species by its notably large eyes and the prominent lateral groove that originates behind the eyes and extends along both sides of the head above the gill openings (Ebert *et al.*, 2021). The specimen examined in this study was consistent with the descriptions of *A. superciliosus* reported in the existing literature (Gruber & Compagno, 1981; Compagno, 1984; Compagno, 2001).

In previous studies, maximum total length (TL) values for bigeye thresher sharks have been reported as 484 and 488 cm (Ebert & Stehmann, 2013; Froese & Pauly, 2025), with Lanteri *et al.* (2017) documenting specimens reaching up to 500 cm TL. Sexual maturity is reported to occur at 245–300 cm TL in males and 282–355 cm TL in females (Chen *et al.*, 1997). Size at birth ranges from 64 to 140 cm TL (Bauchot, 1987; Golani, 1996). The gestation period of *A. superciliosus* is 12 months, with an average litter size of 2–4 pups, typically two. The species is believed to have an annual reproductive cycle (Compagno, 1984; Compagno, 2001). Fernandez-Carvalho *et al.* (2011) report the maximum age at 22 years for females and 17 years for males. Based on its size and developmental characteristics, the specimen presented herein was likely a newly matured adult male.

Table 1 presents a comparative analysis of morphometric measurements from six selected Mediterranean studies alongside our findings. While our Akyatan male specimen shares identical TL (310 cm) with the male specimen from the Dodecanese

region reported by Corsini-Foka & Sioulas (2009), it shows slightly lower values for FDFL, PFL, and PFVF, but exceeds the Dodecanese specimen in other measured parameters. These observed variations may reflect various ecological and biological factors, including feeding ability, habitat, environmental conditions, gonadal maturity, etc.

In the Mediterranean Sea, *A. superciliosus* is considered particularly susceptible to overfishing. In 2014 and 2016, thresher sharks were listed in Appendix II of the Convention on the Conservation of Migratory Species of Wild Animals (CMS), reflecting the Parties' commitment to regional conservation efforts. Additionally, thresher sharks were included in Appendix II of the Convention on International Trade in Endangered Species Wild Fauna and Flora (CITES), requiring exports from CITES Parties to be accompanied by permits demonstrating legal and sustainable sourcing.

The 2018 IUCN Red List classified *A. superciliosus* as vulnerable (VU) globally under criteria A2bd (Rigby *et al.*, 2019) and endangered (EN) in the Mediterranean and European regions (Walls & Soldo, 2016). In the Mediterranean Sea, the thresher shark is both targeted and caught as bycatch in a variety of commercial fishing practices, including longline, bottom trawl, purse seine, and gillnet fisheries (Serena, 2005; Bariche, 2012). While Serena (2005) considers *A. superciliosus* to be an uncommon species in the Mediterranean – captured only sporadically and poorly documented – De Maddalena & Baensch (2005), Clo *et al.* (2008), and Corsini-Foka & Sioulas (2009) argue that the species is not rare in the region and is, in fact, abundant in the eastern Mediterranean Basin.

Incidental captures of sharks – apex predators in the marine food chain – have the potential to disrupt the ecosystem balance and threaten the sustainability of shark populations in the wild (Nurastrì *et al.*, 2024). According to Serena (2005), the sharks' limited restocking capacity and slow population recovery following overfishing are consequences of their late sexual maturity and low fecundity.

Recent years have seen growing emphasis on researching sharks environmental preferences, developing methods to reduce their incidental catch in commercial fisheries, and implementing sustainable conservation measures for shark species.

A more comprehensive dataset is essential for accurately assessing the scale of shark bycatch mortality in Turkish fisheries. Current evidence suggests that *A. superciliosus* and other shark populations in the Eastern Mediterranean face significant threats. Thus, targeted investigation is necessary to ascertain these species' conservation status specifically within the context of Turkish fisheries.

While *A. superciliosus* was classified as endangered in the Mediterranean region during its most recent IUCN Red List assessment in 2018 (Rigby *et al.*, 2019), its current conservation status remains uncertain. The morphometric data obtained from an incidentally captured male specimen in this study therefore contribute valuable new information to the scientific literature

on this protected species, which maintains a limited population in Mediterranean waters.

ACKNOWLEDGMENTS

The authors would like to thank the boat captain and staff for their help in providing the specimen.

NOV ULOV VELIKOOKE MORSKE LISICE *ALOPIAS SUPERCILIOSUS* LOWE, 1841 IZ TURŠKE OBALE (SEVEROVZHODNO SREDOZEMLJE)

Cem ÇEVİK

Faculty of Fisheries, Çukurova University Balcalı Campus, 01330, Sarıçam/Adana, Türkiye

Deniz ERGÜDEN

Marine Science Department, Faculty of Marine Science and Technology, Iskenderun Technical University, 31200 Iskenderun, Hatay, Türkiye

e-mail: deniz.erguden@iste.edu.tr; derguden@gmail.com

Deniz AYAS

Faculty of Fisheries, Mersin University Yenişehir Campus, 33160, Mersin, Türkiye

POVZETEK

*Samec velikooke morske lisice se je po naključju ujel v vlečno mrežo komercialnega plovila. Meril je 310 cm v skupno dolžino in tehtal 156,0 kg. Ulov se je zgodil 16. januarja 2025 v obalni regiji Akyatan, na globini 30 m. Pred pričujočo študijo je bilo vzdolž severovzhodne sredozemske obale Turčije dokumentiranih 15 zapisov o pojavljanju vrste *A. superciliosus*. Ta študija dopolnjuje obstoječe znanje in pomeni pomemben mejnik z dokumentiranjem prvega zabeleženega pojava odraslega samca velikooke morske lisice v severovzhodnem Sredozemlju, natančneje ob obali Akyatan v Turčiji. Predstavlja tudi celovito analizo morfometričnih meritev osebkov v primerjavi z obstoječimi sredozemskimi zapisi o pojavljanju.*

Ključne besede: velikooka morska lisica, Alopiidae, zapis o pojavljanju, obala Akyatan, Türkiye

REFERENCES

- Akyol, O., T. Ceyhan & C. Capapé (2020):** Capture of a bigeye thresher shark *Alopias superciliosus* (Alopiidae) in Turkish Waters (Eastern Mediterranean Sea). *Annales, Ser. Hist. Nat.*, 30(1), 31-36.
- Ayas, D., H.D. Akbora & B.A. Çiçek (2020):** Occurrence of the young thresher shark *Alopias superciliosus* Lowe, 1841 (Lamniformes: Alopiidae) in the Northeastern Mediterranean Sea. *Comm. J. Biol.*, 4, 28-31.
- Bariche, M. (2012):** Field identification guide to the living marine resources of the Eastern and Southern Mediterranean. FAO Species Identification Guide for Fishery Purposes. Rome, FAO, 610 p.
- Bauchot, M.L. (1987):** Raies at autres batoidés. In: Fisher, M., M. Schneider & M.-L. Bauchot (eds.): Fiches FAO d'Identification des Espèces pour les Besoins de la Pêche. Méditerranée et Mer Noire. Zone de Pêche 37. Revision 1. II, FAO, Rome, pp. 847-885.
- Fernandez-Carvalho, J., R. Coelho, K. Erzini & M.N. Santos (2011):** Age and growth of the bigeye thresher shark, *Alopias superciliosus*, from the pelagic longline fisheries in the tropical northeastern Atlantic Ocean, determined by vertebral band counts. *Aquat. Living Resour.*, 24, 359-368.
- Chen, C.T., K.M. Liu & Y.C. Chang (1997):** Reproductive biology of the bigeye thresher shark, *Alopias superciliosus* (Lowe, 1839) (Chondrichthyes: Alopiidae), in the northwestern Pacific. *Ichthyol. Res.*, 44, 227-235.
- Cigala-Fulgosi, F. (1983):** First record of *Alopias superciliosus* (Lowe, 1840) in the Mediterranean, with notes on some fossil species of the genus *Alopias*. *Ann. Mus. Civ. Stor. Nat. Genova*, 84, 211-229.
- Compagno, L.J.V. (1984):** FAO Species Catalogue. Vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 1 - Hexanchiformes to Lamniformes. FAO Fish. Synop. 125(4/1), FAO, Rome, 249 pp.
- Compagno, L.J.V. (1998):** Alopiidae. Thresher sharks. In: Carpenter K. E. & Niem V. H. (eds.): FAO identification guide for fishery purposes. The Living Marine Resources of the Western Central Pacific. FAO, Rome, pp. 1269-1273.
- Compagno, L.J.V. (2001):** Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Vol. 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). FAO Spec. Cat. Fish. Purp. 1(2), FAO, Rome, 269 pp.
- Coelho, R., J. Fernandez-Carvalho & M.N. Santos (2015):** Habitat use and diel vertical migration of bigeye thresher shark: Overlap with pelagic longline fishing gear. *Mar. Environ. Res.*, 112, 91-99.
- Corsini-Foka, M. & A. Sioulas (2009):** On two old specimens of *Alopias superciliosus* (Chondrichthyes: Alopiidae) from the Aegean waters. *Mar. Biodiv. Rec.*, 2, e72.
- Clo, S., R. Bonfil & E. de Sabata (2009):** Additional records of the bigeye thresher shark, *Alopias superciliosus* from the central and eastern Mediterranean Sea. *JMBA2- Biodiv. Rec.*, 2, e20.
- De Maddalena, A. & H. Baensch (2005):** Haie im Mittelmeer. Frackh-Kosmos Verlags-GmbH & Co., Stuttgart, 240 pp.
- Ebert, D.A. & M. Stehmann (2013):** Sharks, batoids and chimaeras of the North Atlantic. FAO Species Catalogue for Fishery Purposes. No. 7. FAO, Rome, 523 pp.
- Ebert, D.A., M. Dando & S. Fowler (2021):** Sharks of the world: A complete guide. Princeton University Press, 608 pp.
- Ergüden D., M. İğde, C. Turan, D. Ayas & H. Kabasakal (2020):** Occurrence of a large bigeye thresher shark, *Alopias superciliosus* (Lamniformes: Alopiidae), in the northeastern Levantine Sea (İskenderun Bay, Eastern Mediterranean Sea, Turkey). *Annales, Ser. Hist. Nat.*, 30(2), 157-164.
- Farrag, M.M.S. (2017):** New record of the bigeye thresher shark, *Alopias superciliosus* Lowe, 1841 (Family: Alopiidae) from the eastern Mediterranean Sea, Egypt. *Int. J. Fish. Aquat. Stud.*, 5, 316-318.
- Fernandez-Carvalho, J., R. Coelho, K. Erzini & M. Nevs Santos (2011):** Age and growth of the bigeye thresher shark, *Alopias superciliosus*, from the pelagic longline fisheries in the tropical northeastern Atlantic Ocean, determined by vertebral band counts. *Aquat. Liv. Res.*, 24(4), 359-368.
- Froese, R. & D. Pauly (Eds.) (2025):** Fishbase. World Wide Web Electronic Publication. www.fishbase.org. version (10/2024) (accessed: 07 February 2025).
- Gökoğlu, M., S. Teker & D. Julian (2017):** First report of thresher sharks (Alopiidae) in the Gulf of Antalya. *Iran. J. Fish. Sci.*, 16, 1108-1113.
- Gruber S.H. & L.J.V. Compagno (1981):** Taxonomic status and biology of the bigeye thresher, *Alopias superciliosus* (Lowe, 1839). *Fish. Bull., Nat. Mar. Fish. Ser.*, 79, 617-640.
- Kabasakal, H. & S.Ü. Karhan (2008):** On the occurrence of the bigeye thresher shark, *Alopias superciliosus* (Chondrichthyes: Alopiidae), in Turkish waters. *Mar. Biodiv. Rec.*, 1, e69.
- Kabasakal, H., C. Dalyan & A. Yurtsever (2011):** Additional records of the bigeye thresher shark *Alopias superciliosus* (Lowe, 1839) (Chondrichthyes: Lamniformes: Alopiidae) from Turkish waters. *Annales, Ser. Hist. Nat.*, 21, 143-148.
- Kabasakal, H. (2017):** Notes on historical and contemporary catches of lamniform sharks in Turkish waters. *Annales, Ser. Hist. Nat.*, 27, 51-58.
- Kabasakal, H., S.Ü. Karhan & S. Sakıman (2017):** Review of the distribution of large sharks in the seas of Turkey (Eastern Mediterranean). *Cah. Biol. Mar.*, 58, 219-228. <https://doi.org/10.21411/CBM.A.96D9F948>.

Kleitou, P.C., C.I. Antoniou, I. Giovos & D. Kletou (2017): How accurately are we describing the longline bycatch? The case of the ‘rare’ shark *Alopias superciliosus* in eastern Mediterranean. *Int. J. Fish. Aquat. Stud.*, 5, 375-378.

Lanteri, L., L. Castellano & F. Garibaldi (2017): New record of *Alopias superciliosus* Lowe, 1841 in the North-Western Mediterranean and annotated review of the Mediterranean records. *Acta Adriat.*, 58, 313-324.

Mater, S. (2005): Denizlerimizde yeni bir köpekbalığı: *Alopias superciliosus* (Lowe, 1841). *Aquaculture* 1, 1-10. (in Turkish).

Nurastrı, Vi.D., S.B. Prayitno & S W Saputra (2024): Biological characteristics of bigeye thresher shark (*Alopias superciliosus*) landed in Cilacap Ocean fishing port, Cilacap Regency, Central Java, Indonesia. *Asian J. Curr. Res.*, 9(4), 197-206.

Rigby, C.L., R. Barreto, J. Carlson, D. Fernando, S. Fordham, M.P. Francis, K. Herman, R.W. Jabado, K.M. Liu, A. Marshall, N. Pacoureaux, E. Romanov, R.B. Sherley & H. Winker (2019): *Alopias superciliosus*. The IUCN Red List of Threatened Species 2019: e.T161696A894216. https://dx.doi.org/10.2305/IUCN.UK.2019_3.RLTS.T161696A894216.en.

Serena, F. (2005): Field identification guide to the sharks and rays of the Mediterranean and Black Sea. *FAO Species Identification Guide for Fishery Purposes*. FAO, Rome, 87 pp.

Walls, R. & A. Soldo (2016): *Alopias superciliosus* (Europe assessment). The IUCN Red List of Threatened Species 2015: e.T161696A48907814.