

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Založnika/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 6. 12. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / CONTENTSSREDOZEMSKA HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Hakan KABASAKAL**

Analysis of Confirmed Shark Attacks
in the Eastern Mediterranean Sea and
the Sea of Marmara (1827–2025) 169
*Analiza potrjenih napadov morskih psov
v vzhodnem Sredozemskem morju
in Marmarskem morju (1827–2025)*

Alen SOLDO

Observations of a Juvenile Basking Shark
Cetorhinus maximus in the Adriatic Sea
Support the Hypothesis of a Distinct
Mediterranean Population 187
*Opazovanja mladega morskega psa
orjaka (Cetorhinus maximus)
v Jadranskem morju podpirajo
hipotezo o ločeni sredozemski
populaciji*

**Jacopo BERNARDI, Pero UGARKOVIĆ &
Ilija ČETKOVIĆ**

An Unusual Encounter with a Juvenile Kitefin
Shark, *Dalatias licha*, in Shallow Coastal
Waters of Montenegro (Adriatic Sea) 197
*Nenavadno srečanje z mladim primerkom
klinoplavutega morskega psa, Dalatias licha,
v plitvih obalnih vodah Črne gore
(Jadransko morje)*

**Cemal TURAN, Alen SOLDO, Servet A. DOĞDU,
Funda TURAN, Aysegül ERGENLER & Ali UYAN**

Identification of a Potential Nursery Ground
of the Spiny Butterfly Ray, *Gymnura altavela*,
in the Northeastern Mediterranean Sea, Türkiye 205
*Identifikacija potencialnih jaslčic za
metuljastega skata, Gymnura
altavela, v severovzhodnem
Sredozemskem morju,
Turčija*

**Alen SOLDO, Eleonora DE SABATA &
Simona CLO**

Recent Records of Critically Endangered
Common Guitarfish *Rhinobatos rhinobatos*
(Linnaeus, 1758) in the Northern
Mediterranean 213
*Nedavni zapiski o pojavljanju kritično
ogroženega navadnega goslaša
Rhinobatos rhinobatos
(Linnaeus, 1758) v
severnem Sredozemlju*

Hakan KABASAKAL & F. Saadet KARAKULAK

Demersal Elasmobranchs of the Sea of Marmara:
Updated Inventory, Taxonomic Issues and
Environmental Implications 221
*Pridnene hrustančnice v Marmarskem morju:
posodobljena inventarizacija, taksonomska
vprašanja in okoljske posledice*

IHTIOLOGIJA**ITTILOGIA****ICHTHYOLOGY****Jamila RIZGALLA, Amani FITORI, Francesco
TIRALONGO & Abdalrh BEN ABDALAH**

First Records of Blennies (Suborder
Blennioidea) off the Coast of Libya 245
*Prvi zapisi o pojavljanju babic
(podred Blennioidea) ob
libijski obali*

**Farid HEMIDA, Sara LADOUL, Christian
REYNAUD & Christian CAPAPÉ**

Confirmed Occurrence of the Mediterranean
Spearfish *Tetrapturus belone* (Osteichthyes:
Istiophoridae) from the Algerian Coast
(Southwestern Mediterranean Sea) 257
*Potrjeno pojavljanje sredozemske
jadrovnice, Tetrapturus belone
(Osteichthyes: Istiophoridae) iz
alžirske obale (jugovzhodno
Sredozemsko morje)*

Lana KHREMA, Adib SAAD & Christian CAPAPÉ

First Substantiated Record of Blackfish
Centrolophus niger (Centrolophidae)
from the Syrian Coast (Eastern
Mediterranean Sea) 263
Prvi potrjen zapis črnuha, Centrolophus niger
(*Centrolophidae*), s sirske obale
(vzhodno Sredozemsko morje)

Amir IBRAHIM, Chirine HUSSEIN, Firas ALSHAWY & Alaa ALCHEIKH AHMAD

Marine Fishes (Teleostei / Osteichthyes)
of Syria (Eastern Mediterranean):
an Updated Checklist 269
Morske ribe (Teleostei / Osteichthyes)
Sirije (vzhodno Sredozemsko morje):
posodobljeni seznam vrst

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO, Enrico LA SPINA, Daniele TIBULLO & Francesco TIRALONGO

Cassiopea andromeda at the Southernmost
Tip of Italy: a Recent Arrival or an
Overlooked Resident? 287
Tujerodni klobučnjak Cassiopea andromeda
na najjužnejši konici Italije: nedavni
prišlek ali spregledan prebivalec?

Mourad CHÉRIF, Rimel BENMESSAOUD, Sihem RAFRAFI-NOUIRA, Mohamed Nouri AYADI & Christian CAPAPÉ

First Substantiated Record of the Golden-Banded
Goatfish *Upeneus moluccensis* (Osteichthyes:
Mullidae) from the Coast of Tunisia
(Central Mediterranean Sea) 295
Prvi potrjen zapis o zlatoprogem bradaču
Upeneus moluccensis (Osteichthyes: Mullidae)
z obale Tunizije (osrednje Sredozemsko morje)

Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ČETKOVIĆ, Borut MAVRIČ & Lovrenc LIPEJ

New Record of the Long-Jawed Squirrelfish,
Holocentrus adscensionis (Osbeck, 1765),
in the Adriatic Sea 303
Novi zapis o pojavljanju veveričjaka vrste
Holocentrus adscensionis (Osbeck, 1765)
v Jadranskem morju

MORSKA FAVNA
FAUNA MARINA
MARINE FAUNA

Nicola BETTOSO, Lisa FARESI, Saul CIRIACO, Marco FANTIN & Marco SEGARICH

New Findings of the Cup-Shaped Demosponge
Calyx nicaeensis (Risso, 1826) on the Rocky
Outcrops in the Gulf of Trieste
(Northern Adriatic Sea) 311
Nove najdbe morskega keliha Calyx nicaeensis
(*Risso, 1826*) *na skalnih osamelcih v Tržaškem*
zalivu (severno Jadransko morje)

Aleksandra V. BORODINA & Yuri O. VELIAEV

Features of Lipid Accumulation in Striped Venus
Clam *Chamelea gallina* in the Sublittoral
Zone of the Crimean Coast (Black Sea) 319
Značilnosti kopičenja lipidov v navadni
venerici (Chamelea gallina) v obrežnem
pasu krimske obale (Črno morje)

Okan AKYOL & Oğuzhan TAKICAK

Recent Observations on *Monachus monachus*
(Phocidae) at Sea-Cage Fish Farms in Izmir
(Turkish Aegean Sea) 329
Nedavna opažanja primerkov Monachus monachus
(*Phocidae*) *v ribogojnicah z morskimi kletkami v*
Izmirju (turško Egejsko morje)

FLORA
FLORA
FLORA

Amelio PEZZETTA & Marco PAOLUCCI

La Flora di Palena (Parco Nazionale della
Majella): Aggiornamento Floristico 337
Flora Palene (Nacionalni park Majella):
floristična posodobitev

DELO NAŠIH ZAVODOV IN DRUŠTEV
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS AND
ASSOCIATIONS

Bojana LIPEJ

Reviving Landscapes, Connecting Species:
Lessons from the ReCo Project 383

IN MEMORIAM

Tom TURK

V spomin Marjanu Richterju (1935–2025) 389

Kazalo k slikam na ovitku 391
Index to images on the cover 391

received: 2025-06-16

DOI 10.19233/ASHN.2025.25

DEMERSAL ELASMOBRANCHS OF THE SEA OF MARMARA: UPDATED INVENTORY, TAXONOMIC ISSUES, AND ENVIRONMENTAL IMPLICATIONS

Hakan KABASAKAL

Istanbul University, Institute of Graduate Studies in Sciences, Fisheries Technologies and Management Program, Istanbul, Türkiye
WWF Türkiye, İstanbul, Türkiye
e-mail: kabasaka.hakan@gmail.com

F. Saadet KARAKULAK

Department of Fisheries Technologies and Management, Faculty of Aquatic Sciences, İstanbul University, İstanbul, Türkiye

ABSTRACT

A total of 16 elasmobranch species (nine shark and seven batoid species) were sampled in the Sea of Marmara (SoM) between 2014 and 2024 for this study. During the same period, an additional nine species (six shark and three batoid species) were recorded from the same area by other researchers. Over the past 100 years, 32 species of demersal elasmobranchs (16 shark and 16 batoid species) have been reported from the SoM, but applying “evidence-based confirmed or unconfirmed presence” criteria reduces the current number to 25 (15 shark species, 10 batoid species). Principal Component Analysis indicates that adaptability to shallower waters is now critical to the spatial distribution of demersal elasmobranchs in the SoM. This shift clearly shows how dramatically the environmental degradation, marine pollution, and deoxygenation have reduced the region’s elasmobranch fauna primarily to demersal species over the last 40 years, underscoring the potentially devastating impact of such degradation on biodiversity.

Key words: Elasmobranchii, survival, conservation, inventory, Sea of Marmara, Turkey

ELASMOBRANCHI DEMERSALI DEL MARE DI MARMARA: INVENTARIO AGGIORNATO, QUESTIONI TASSONOMICHE E IMPLICAZIONI AMBIENTALI

SINTESI

Per questo studio, tra il 2014 e il 2024 sono state campionate nel Mar di Marmara (SoM) un totale di 16 specie di elasmobranchi (nove specie di squali e sette specie di batoidi). Nello stesso periodo, altri ricercatori hanno registrato altre nove specie (sei specie di squali e tre specie di batoidi) nella stessa area. Negli ultimi 100 anni, nel SoM sono state segnalate 32 specie di elasmobranchi demersali (16 specie di squali e 16 specie di batoidi), ma applicando i criteri di “presenza confermata o non confermata basata su prove” il numero attuale si riduce a 25 (15 specie di squali e 10 specie di batoidi). L’analisi delle componenti principali indica che l’adattabilità alle acque meno profonde è ora fondamentale per la distribuzione spaziale degli elasmobranchi demersali nel Mar di Marmara. Questo cambiamento mostra chiaramente come il degrado ambientale, l’inquinamento marino e la deossigenazione abbiano ridotto drasticamente la fauna di elasmobranchi della regione, principalmente alle specie demersali, negli ultimi 40 anni, sottolineando l’impatto potenzialmente devastante di tale degrado sulla biodiversità.

Parole chiave: Elasmobranchii, sopravvivenza, conservazione, inventario, Mar di Marmara, Turchia

INTRODUCTION

Fauna inventories provide the most direct means of assessing the components of animal diversity in a specific biome or locality at a given time (Silveira *et al.*, 2010). From this perspective, the elasmobranch species inventory of the Sea of Marmara (SoM) has been updated many times in studies by various researchers over the past 100 years. These studies either (1) include elasmobranchs as part of the region's general ichthyofauna (Ninni, 1923; Deveciyan, 1926; Rhasis Erazi, 1942; Kocataş *et al.*, 1993; Eryılmaz & Meriç, 2005; Torcu Koç *et al.*, 2012; Daban *et al.*, 2021; Bilecenoğlu, 2024), or (2) focus specifically on the elasmobranch fauna of the SoM (Kabasakal, 2016, 2022; Artüz & Fricke, 2024; Karadurmuş & Sarı, 2024; Kabasakal & Karakulak, 2024). In one of the earliest general ichthyological studies listing sharks and batoids, Ninni (1923) reported 21 elasmobranch species among the 162 fish species landed at Istanbul Wholesale Fish Market. In another pioneering study, Deveciyan (1926) provided detailed information on 16 species of sharks and batoids landed at the same market. However, neither author specified the exact locations in the SoM where these elasmobranchs were captured. Although the most recent surveys report 35 (Bilecenoğlu, 2024) and 38 (Artüz & Fricke, 2024) elasmobranch species in the ichthyofauna of the SoM, the number of species with confirmed current presence in the region decreases to 25 (Kabasakal & Karakulak, 2024) when evidence-based confirmation criteria are applied (Kovačić *et al.*, 2020).

Based on commercial landings data, several researchers have pointed to a dramatic reduction (of up to 98%) in demersal elasmobranch species in the SoM over the past three decades (Demirel *et al.*, 2020; Gül & Demirel, 2021). It has recently been suggested that this decline is primarily due to overfishing and deteriorating environmental conditions, particularly the deoxygenation of bottom waters (Kabasakal, 2025). According to Silveira *et al.* (2010), fauna inventories are essential for assessing projects with myriad environmental impacts, many of which may be significant and irreversible. Therefore, creating an accurate 'snapshot' of the specific locality to be impacted is a task of utmost importance and responsibility. Given that proper species identification is the foundational step in effective fisheries management (Stauffer Jr. & Kovcovsky, 2007), updating the inventory of demersal elasmobranchs in the environmentally degraded and deoxygenated SoM is a critical initiative that extends far beyond mere taxonomic curiosity. The present article provides an updated and confirmed list of demersal elasmobranch species in the SoM, based on the latest environmental monitoring and

stock assessment surveys. In addition, it discusses key environmental threats to the future survival of these species in the SoM, as well as factors contributing to taxonomic inconsistencies.

MATERIAL AND METHODS

An overview of the SoM

A significant environmental issue currently threatening marine ecosystems is the emergence and expansion of 'dead zones' – hypoxic or anoxic marine areas caused by human activity (Diaz, 2016). The SoM is experiencing severe hypoxia in its deep waters, with conditions in some areas progressing towards anoxia (Mantıkçı *et al.*, 2022; Salihoğlu *et al.*, 2022). Hypoxia in seawater is defined as a decrease in dissolved oxygen (DO) concentration to below 2 mg/L (or 80 µM; Diaz & Rosenberg, 2008; Vaquer-Sunyer & Duarte, 2008). The SoM, designated as geographical subarea (GSA) 28 by the General Fisheries Commission for the Mediterranean (GFCM, 2018), is one of the most damaged marine ecosystems in the Mediterranean Basin (Saygu *et al.*, 2023).

It is a relatively small marine basin with a surface area of 11,500 km² and a maximum depth of 1,390 m, connected to the Mediterranean and Black Seas through narrow straits (Beşiktepe *et al.*, 1994; Fig. 1). Its hydrographic characteristics are largely shaped by water exchange through these straits. A distinctive feature of this system is the persistent oxygen deficiency in the deeper sub-halocline layers (Ünlüata & Özsoy, 1988; Beşiktepe *et al.*, 1994).

Acting as a receiving and settling basin for organic matter transported from the Black Sea via the Bosphorus Strait (Ünlüata & Özsoy, 1988; Yılmaz *et al.*, 1990), the SoM has experienced substantial ecological degradation over the past four decades due to increasing anthropogenic pressures. As a result of eutrophication, DO concentrations – particularly in bathyal waters – have fallen below the hypoxia threshold (Yılmaz *et al.*, 1990; Mantıkçı *et al.*, 2022). In the eastern and northeastern parts of the SoM, where deoxygenation is most severe, hypoxic conditions now extend onto the continental shelf (ÇŞİDB, TUBİTAK-MAM, 2021, p. 27; Kabasakal, 2025).

Bottom trawl surveys

The bottom trawl surveys conducted throughout the SoM (Fig. 1) between 2014 and 2024 were mainly carried out as part of two projects: (1) the Stock Assessment of Demersal Fish Species in the Eastern Marmara Sea (Project No. 51922), and (2) the Marine Monitoring (DEN-İZ) project of the Ministry of Environment, Urbanisation and Climate Change.




Fig. 1: Map of the study area showing sampling stations. Solid red circles denote the fixed stations of Project 51922, and solid purple squares indicate the fixed stations of DEN-İZ project.

Sl. 1: Zemljevid obravnavanega območja raziskave, ki prikazuje vzorčevalne postaje. Rdeči krogi označujejo fiksne postaje projekta 51922, vijolični kvadrati pa fiksne postaje projekta DEN-İZ.

Sampling was performed on board *R/V Yunus-S*, a 510 hp stern trawler operated by the Istanbul University's Faculty of Aquatic Sciences. A bottom trawl with a cod-end mesh size of 14 mm (approximately 22 mm mesh opening) was used for the hauls following the MEDITS protocol (Anonymous, 2017). Haul duration was set at 30 min for depths under 200 m and 60 min for depths over 200 m, with a standard towing speed of 3 knots (Anonymous, 2017). Oceanographic parameters (salinity, S; temperature, T; and dissolved oxygen, DO) were recorded using a SeaBird CTD probe. The start and end coordinates of each trawl haul, along with the corresponding seafloor depths, were determined using the vessel's onboard GPS and echo sounder.

Handling of specimens and species identification

To reduce at-vessel mortality and increase post-release survival of captured sharks and rays, we followed best-practice procedures proposed by Saygu and Deval (2014) and Ellis *et al.* (2016a). An on-board survival tank was improvised using a large container with a continuous supply of fresh seawater via a hose. Large specimens were handled according to the FAO and ACCOBAMS (2018) guides. All individuals were counted, weighed, and measured as soon as possible. With the exception of a few voucher samples, the remaining elasmobranch catch was gently released. Voucher samples are preserved at the laboratory of

the Department of Fisheries Technologies and Management, Faculty of Aquatic Sciences, Istanbul University. For very large specimens (e.g., *Hexanchus griseus* or *Echinorhinus brucus*), photographs were taken and are kept in the authors' personal archives. Species identification is based on Serena (2005), Ebert and Stehmann (2013), Last *et al.* (2016), Ebert *et al.* (2021), and Barone *et al.* (2022). Scientific names follow WoRMS Editorial Board (2025).

The current status of elasmobranch species not sampled in the present survey, but known to occur in the SoM from previous studies (Ninni, 1923; Rhasis Erazi, 1942; Geldiay, 1969; Anonymous, 1974; JICA, 1993; Benli *et al.*, 1993; Meriç, 1995; Uysal *et al.*, 1996; Zengin *et al.*, 2004; Yaka & Yüce, 2006; Torcu Koç *et al.*, 2012; Yıldız *et al.*, 2016; Bilecenoğlu, 2019; Kabasakal *et al.*, 2023a, 2024a,b; Karadurmuş & Sarı, 2024), was assessed using evidence-based criteria for confirmed and unconfirmed presence (Kovačić *et al.*, 2020). Detailed descriptions of these criteria are provided by Kovačić *et al.* (2020).

Dispersal/settlement success of demersal elasmobranch species

Principal Component Analysis (PCA) was used to examine the effect of life history characteristics and bathymetric distribution on the past and present abundance of demersal elasmobranch species in

Tab. 1: Species-specific life history traits used in the PCA of demersal elasmobranch species of the SoM. TL – total length; L_mM , L_mF length at 50% sexual maturity (males, females); GL – generation length. Continues in the next section.

Tab. 1: Vrstno specifične značilnosti življenjskega cikla, uporabljene pri PCA pridnenih vrst rib v Marmarskem morju. TL – skupna dolžina; L_mM , L_mF – dolžina pri doseženi 50% spolni zrelosti (samci, samice); GL – dolžina generacije. Nadaljevanje v naslednjem razdelku.

Species	FAO code	TL (cm)	L_mM (cm)	L_mF (cm)	GL (years)
<i>Hexanchus griseus</i> (Bonnaterre, 1788)	SBL	600	400	330	53
<i>Galeus melastomus</i> Rafinesque, 1810	SHO	90	42	45	5.7
<i>Scyliorhinus canicula</i> (Linnaeus, 1758)	SYC	80	56	60	?
<i>Scyliorhinus stellaris</i> (Linnaeus, 1758)	SYT	162	77	79	20
<i>Mustelus asterias</i> Cloquet, 1819	SDS	140	85	96	13
<i>Mustelus mustelus</i> (Linnaeus, 1758)	SMD	175	112	124	17.8
<i>Mustelus punctulatus</i> Risso, 1827	MPT	190	88.5	100	17.8
<i>Galeorhinus galeus</i> (Linnaeus, 1758)	GAG	195	170	185	30
<i>Dalatias licha</i> (Bonnaterre, 1788)	SCK	182	100	120	28.7
<i>Oxynotus centrina</i> (Linnaeus, 1758)	OXY	150	60	66	20
<i>Centrophorus uyato</i> (Rafinesque, 1810)	CPU	128	91	89	20
<i>Squalus acanthias</i> Linnaeus, 1758	DGS	195	64	93	8
<i>Squalus blainville</i> (Risso, 1827)	QUB	92	56	60	19.5
<i>Echinorhinus brucus</i> (Bonnaterre, 1788)	SHB	394	190	230	30
<i>Squatina oculata</i> Cuvier, 1829	SUT	160	82	100	12
<i>Squatina squatina</i> (Linnaeus, 1758)	AGN	244	132	169	11
<i>Tetronarce nobiliana</i> (Bonaparte, 1835)	TTO	180	?	?	?
<i>Torpedo marmorata</i> Risso, 1810	TTR	100	44	49	16
<i>Torpedo torpedo</i> (Linnaeus, 1758)	TTV	60	?	?	12.5
<i>Dipturus batis</i> (Linnaeus, 1758)	RJB	285	197.5	185.5	20
<i>Dipturus oxyrinchus</i> (Linnaeus, 1758)	RJO	150	103.5	91	10
<i>Leucoraja naevus</i> (Müller & Henle, 1841)	RJN	81	56.16	54.96	8
<i>Raja asterias</i> Delaroche, 1809	JRS	75	56	52	4.36
<i>Raja clavata</i> Linnaeus, 1758	RJC	105	85	75	11.5
<i>Raja miraletus</i> Linnaeus, 1758	JAI	63	41.8	34.3	7.2
<i>Raja montagui</i> Fowler, 1910	RJM	102	60.7	53.74	7
<i>Raja radula</i> Delaroche, 1809	JAR	70	46	40	9
<i>Dasyatis pastinaca</i> (Linnaeus, 1758)	JDP	69.5	32	38	7.5
<i>Dasyatis tortonesei</i> Capapé, 1975	JDO	80	38	47	?
<i>Gymnura altavela</i> (Linnaeus, 1758)	RGL	400	105	102	7
<i>Aetomylaeus bovinus</i> (Geoffroy St. Hilaire, 1817)	MPO	222	90	90	15
<i>Myliobatis aquila</i> (Linnaeus, 1758)	MYL	183	40	60	11

Tab. 1 (Continued): MRD – maximum recorded depth range; CDR – common depth range; TRL – trophic level. Continues in the next section.

Tab. 1 (nadaljevanje): MRD – največji zabeleženi razpon globin; CDR – običajni razpon globin; TRL – trofična raven. Nadaljevanje v naslednjem razdelku.

Species	MDR (m)			MDR (m)	CDR (m)			CDR (m)	TRL
	200–500	501–1000	>1000		0–100	101–200	>200		
<i>H. griseus</i>	0	0	1	2500	0	0	1	200–1100	4.48
<i>G. melastomus</i>	0	0	1	2000	0	0	1	200–500	4.2
<i>S. canicula</i>	0	1	0	800	1	1	1	0–450	3.8
<i>S. stellaris</i>	1	0	0	380	1	0	0	20–63	4
<i>M. asterias</i>	1	0	0	200	1	1	0	0–200	3.6
<i>M. mustelus</i>	0	1	0	800	1	0	0	5–50	4.3
<i>M. punctulatus</i>	1	0	0	200	1	1	0	0–200	3.8
<i>G. galeus</i>	0	1	0	800	1	1	1	0–800	4.3
<i>D. licha</i>	0	0	1	1800	0	0	1	1800	4.2
<i>O. centrina</i>	0	1	1	805	1	1	1	35–805	3.1
<i>C. uyato</i>	0	0	1	1400	0	1	1	115–745	4.5
<i>S. acanthias</i>	0	0	1	1978	1	1	1	0–600	4.4
<i>S. blainville</i>	0	0	1	1500	1	1	1	15–1500	4
<i>E. brucus</i>	0	0	1	1214	1	1	1	10–1214	4.4
<i>S. oculata</i>	1	0	0	500	1	0	0	50–100	4
<i>S. squatina</i>	1	0	0	150	1	1	0	1–150	4.1
<i>T. nobiliana</i>	0	1	0	800	1	1	0	10–150	4.5
<i>T. marmorata</i>	1	0	0	370	1	1	0	2–370	4.5
<i>T. torpedo</i>	1	0	0	400	1	0	0	2–70	4.5
<i>D. batis</i>	0	1	0	1000	1	1	1	30–600	4.1
<i>D. oxyrinchus</i>	0	1	1	1461	0	0	1	200	3.5
<i>L. naevus</i>	0	1	0	900	1	1	0	20–250	3.6
<i>R. asterias</i>	0	1	0	700	1	0	0	20–50	3.5
<i>R. clavata</i>	0	1	1	1020	1	0	0	10–60	4.2
<i>R. miraletus</i>	1	0	0	462	1	1	0	50–150	3.8
<i>R. montagui</i>	1	1	0	530	1	1	0	20–120	4
<i>R. radula</i>	1	0	0	300	1	1	1	0–300	3.7
<i>D. pastinaca</i>	1	0	0	200	1	0	0	20–35	4.1
<i>D. tortonesei</i>	1	0	0	200	0	1	0	100–200	4
<i>G. altavela</i>	1	0	0	100	1	0	0	5–100	4.5
<i>A. bovinus</i>	1	0	0	150	1	1	0	10–150	3.8
<i>M. aquila</i>	1	0	0	300	1	1	0	1–300	3.6

Tab. 1 (Continued): DFRSoM – Date of first record in the Sea of Marmara; DLRSoM – Date of last record in the Sea of Marmara; SLF – continental shelf; SUS – shelf-upper slope; DWT – deep bathyal water. Data on L_mM , L_mF , GL, and TRL were obtained from the literature cited in the reference column.

Tab. 1 (nadaljevanje): DFRSoM – Datum prvega zapisa v Marmarskem morju; DLRSoM – Datum zadnjega zapisa v Marmarskem morju; SLF – kontinentalni prag; SUS – kontinentalni prag-zgornje pobočje; DWT – globoki batial. Podatki o L_mM , L_mF , GL in TRL so bili pridobljeni iz literature, navedene v referencah.

Species	Before 1990	DFR SoM	DLR SoM	After 2000	SLF	SUS	DWT	Reference
<i>H. griseus</i>	1	1923	2022	1	0	1	1	Finucci <i>et al.</i> (2020)
<i>G. melastomus</i>	0	1993	2017	1	0	1	1	Abella <i>et al.</i> (2016)
<i>S. canicula</i>	1	1923	2024	1	1	1	0	Serena <i>et al.</i> (2015)
<i>S. stellaris</i>	1	1923	2023	1	1	0	0	Ellis <i>et al.</i> (2016b)
<i>M. asterias</i>	1	1969	2024	1	1	0	0	Farrell <i>et al.</i> (2016a)
<i>M. mustelus</i>	1	1923	2024	1	1	0	0	Farrell <i>et al.</i> (2016b)
<i>M. punctulatus</i>	0	2023	2023	1	1	0	0	Ebert <i>et al.</i> (2021)
<i>G. galeus</i>	1	1923	2021	1	1	0	0	McCully <i>et al.</i> (2016)
<i>D. licha</i>	0	1991	1991	0	0	1	0	Walls & Guallart (2016)
<i>O. centrina</i>	1	1942	2024	1	0	1	0	Soldo & Guallart (2016)
<i>C. uyato</i>	0	1992	2019	1	0	1	1	Guallart & Walls (2015)
<i>S. acanthias</i>	1	1923	2023	1	1	1	0	Ellis <i>et al.</i> (2016c)
<i>S. blainville</i>	1	1923	2024	1	1	1	0	Soldo <i>et al.</i> (2016)
<i>E. brucus</i>	1	1923	2023	1	1	1	1	Ferretti & Buscher (2016)
<i>S. oculata</i>	0	1995	2018	1	1	1	0	Ferretti <i>et al.</i> (2016a)
<i>S. squatina</i>	1	1923	2023	1	1	0	0	Ferretti <i>et al.</i> (2016b)
<i>T. nobiliana</i>	?	?	?	?	1	1	0	Finucci <i>et al.</i> (2021)
<i>T. marmorata</i>	1	1923	2023	1	1	1	0	Notarbartolo di Sciarra <i>et al.</i> (2016)
<i>T. torpedo</i>	1	1923	2023	1	1	1	0	Serena <i>et al.</i> (2016a)
<i>D. batis</i>	1	1923	2023	1	1	1	0	Ellis <i>et al.</i> (2021)
<i>D. oxyrinchus</i>	1	1923	2024	1	0	1	1	Ellis <i>et al.</i> (2016d)
<i>L. naevus</i>	1	1996	1996	0	1	1	0	Ellis & Dulvy (2016)
<i>R. asterias</i>	0	1996	2009	1	1	0	0	Serena <i>et al.</i> (2016b)
<i>R. clavata</i>	1	1923	2023	1	1	0	0	Ellis <i>et al.</i> (2016e)
<i>R. miraletus</i>	1	1923	2014	1	1	0	0	Dulvy <i>et al.</i> (2020)
<i>R. montagui</i>	1	1996	1996	0	1	0	0	Ellis <i>et al.</i> (2016f)
<i>R. radula</i>	0	2006	2023	1	1	1	0	Mancusi <i>et al.</i> (2016)
<i>D. pastinaca</i>	1	1942	2023	1	1	0	0	Serena <i>et al.</i> (2016c)
<i>D. tortonesei</i>	0	2016	2016	1	1	0	0	Jabado & Derrick (2021)
<i>G. altavela</i>	0	1984	?	0	1	0	0	Walls <i>et al.</i> (2016)
<i>A. bovinus</i>	0	?	2019	1	1	0	0	Walls & Buscher (2016)
<i>M. aquila</i>	1	1923	2024	1	1	1	0	Serena <i>et al.</i> (2016d)

the SoM (Villagra *et al.*, 2022). This analysis tested the hypothesis that “demersal elasmobranch species with a wider bathymetric distribution range are more likely to have adapted to, and persisted in, the SoM”. The most recent species lists (Artüz & Fricke, 2024; Kabasakal & Karakulak, 2024; Karadurmuş & Sarı, 2024) were consulted to compile the life history parameters of the 32 demersal elasmobranch species reported from the SoM over the past century and currently categorised as ‘present’, ‘absent’ or ‘doubtful’ in this region. Information on the species’ first and last recorded occurrences (date of first record in the Sea of Marmara (DFRSOM) and date of last record in the Sea of Marmara (DLRSOM) in Table 1), as well as depth distribution in the SoM (continental shelf occurrence (SLF), shelf-upper slope occurrence (SUS) and deep bathyal water occurrence (DWT) in Table 1) were retrieved from the literature (Tab. 1), specifically from Ninni (1923), Rhasis Erazzi (1942), Geldiay (1969), Anonymous (1974), JICA (1993), Benli *et al.* (1993), Meriç (1995), Uysal *et al.* (1996), Zengin *et al.* (2004), Yaka and Yüce (2006), Torcu Koç *et al.* (2012), Yıldız *et al.* (2016), Bilecenoglu (2019), Kabasakal *et al.* (2023a, 2024a, b), and Karadurmuş and Sarı (2024). The sources for data on total length (TL), male and female sexual maturation lengths (L_m and L_f , respectively), generation length (GL), depth distribution, and habitat are provided in Table 1. These life history parameters, compiled from the literature, were first organised in a raw data table. A matrix was then formulated where these data were expressed as numerical values (maximum TL and sexual maturation length in cm; GL in years; trophic level) and binary values (0 for absent and 1 for present). The PCA analysis was based on this final data matrix (Villagra *et al.*, 2022), using PAST - Paleontological Statistics ver. 4.03 (Hammer *et al.*, 2001; Rincón-Díaz *et al.*, 2018).

RESULTS AND DISCUSSION

Updated species inventory and taxonomic issues

Sixteen elasmobranch species (nine sharks and seven batoids; Tab. 2) were sampled between 2014 and 2024 in the present study (Fig. 2). During the same period, an additional nine species (six sharks and three batoids; Table 2) were recorded in same area by Bilecenoglu (2019), Daban *et al.* (2021), Kabasakal and Türetken (2021), Karadurmuş and Sarı (2024), and Kabasakal *et al.* (2024a, b). As evidenced by the ichthyological inventories presented in Table 2, a total of 32 species of demersal elasmobranchs (16 sharks and 16 batoids) have been reported from the SoM over the past 100 years. However, using the criterion proposed by Bilecenoglu (2024) – including only species whose

presence has been confirmed within the past 10 years – sets the current count at 25 (15 sharks and 10 batoids; Table 2). This figure is remarkably lower than the totals reported by Bilecenoglu (2024; $n = 34$) and Artüz and Fricke (2024; $n = 38$), as these authors included in their checklists all shark and batoid species recorded in the SoM since 1923, regardless of whether any had been observed in the region within the past decade or not. Furthermore, the presence of the 25 species was assessed applying the criteria for “evidence-based confirmed or unconfirmed presence” proposed by Kovačić *et al.* (2020). Only 16 species have a verified presence under Criterion 1, which requires that at least one specimen from the SoM is stored in a scientific collection with a published reference. Six species were confirmed under Criterion 2, which applies where there are no stored specimens, but the species can be positively identified from a photo of the SoM locality provided in a published record. Three species were confirmed under Criterion 5; this is used when there are no stored specimens or published morphological/genetic evidence, but species citations can be traced back to an author – with expert knowledge of the taxon or fish guild of a particular habitat – who reported numerous findings (or, in a study on ecology or species biology, numerous specimens), making correct identification highly probable.

Although *Dalatias licha*, *Torpedo torpedo*, and *Leucoraja naevus* each have at least one positive historical record from the SoM and meet the criteria for verified presence (1, 5, and 2, respectively), they were not included in the updated inventory because none has been recorded in the region during the past decade. According to Bilecenoglu (2024), *Tetronarce nobiliana* and *Gymnura altavela* are also present in the SoM; however, their listing among elasmobranch fauna of the SoM is based on a historical ichthyological inventory (Bilecenoglu *et al.*, 2014). A critical retrospective review of ichthyological checklists for the seas of Turkey, published at roughly ten-year intervals (Bilecenoglu *et al.*, 2002, 2014; Bilecenoglu, 2024), reveals that all three studies cite the occurrence of *T. nobiliana* and *G. altavela* in the SoM based solely on Mater and Meriç (1996) – the first inventory of Turkish marine fishes. This original source, however, provides no information where the examined specimens were captured or deposited in a scientific collection for verification. Since the evidence was insufficient to confirm the contemporary occurrences of *T. nobiliana* and *G. altavela* in the SoM, the only applicable classification was exclusion Criterion 13 (Kovačić *et al.*, 2020). This applies when a species’ reported presence is based on a review publication citing an original reference that in itself contains no

Tab. 2: Chronological summary of demersal elasmobranch species ($n=32$) recorded in the SoM by different researchers between 1923 and 2024. The numbers in the top row refer to source references that reported shark and batoid records from the SoM (indicated by +). These references are as follows: (1) Ninni (1923), (2) Anonymous (1974), (3) JICA (1993), (4) Meriç (1995), (5) Mater and Meriç (1996), (6) Uysal et al. (1996), (7) Kabasakal (2003), (8) Zengin et al. (2004), (9) Yaka and Yüce (2006), (10) Torcu Koç et al. (2012), (11) Yıldız et al. (2016), (12) Bilecenoğlu (2019), (13) Kabasakal and Türetken (2021), (14) Daban et al. (2021), (15) Karadurmuş and Sarı (2024), (16) Kabasakal et al. (2023a), (17) Kabasakal et al. (2024a), (18) Kabasakal et al. (2024b), and (19) elasmobranch species captured during the 2014–2024 sampling period of the present study. Demersal elasmobranch species recorded in the SoM in the last 10 years are marked with a check (✓) in the 2025 column; these records constitute the updated list of elasmobranch species. C denotes the applicable criterion for confirmed or unconfirmed presence (Kovačić et al., 2020). Elasmobranch species not included in the updated inventory are shown in bold.

Tab. 2: Kronološki povzetek pridnenih vrst hrustančnic ($n=32$), ki so jih v Marmarskem morju zabeležili različni raziskovalci med letoma 1923 in 2024. Številke v zgornji vrstici se nanašajo na vire, ki so poročali o zapisih o morskih psih in skatih iz Marmarskega morja (označeno z +). Ti viri so naslednji: (1) Ninni (1923), (2) Anonymous (1974), (3) JICA (1993), (4) Meriç (1995), (5) Mater in Meriç (1996), (6) Uysal in sod. (1996), (7) Kabasakal (2003), (8) Zengin in sod. (2004), (9) Yaka in Yüce (2006), (10) Torcu Koç in sod. (2012), (11) Yıldız in sod. (2016), (12) Bilecenoğlu (2019), (13) Kabasakal in Türetken (2021), (14) Daban in sod. (2021), (15) Karadurmuş in Sarı (2024), (16) Kabasakal in sod. (2023a), (17) Kabasakal in sod. (2024a), (18) Kabasakal in sod. (2024b) in (19) vrste pridnenih hrustančnic, ujete v obdobju vzorčenja 2014–2024 v tej študiji. Pridnene vrste hrustančnic, zabeležene v Marmarskem morju v zadnjih 10 letih, so označene s kljukico (✓) v stolpcu 2025; ti zapisi predstavljajo posodobljen seznam vrst hrustančnic. C označuje veljavno merilo za potrjeno ali nepotrjeno prisotnost (Kovačić in sod., 2020). Vrste hrustančnic, ki niso vključene v posodobljeni popis, so prikazane s krepko pisavo.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	2025	C
<i>Hexanchus griseus</i>	+			+			+							+					+	✓	1
<i>Galeus melastomus</i>			+	+		+	+												+	✓	1
<i>Scyliorhinus canicula</i>	+	+	+	+				+		+				+	+				+	✓	1
<i>Scyliorhinus stellaris</i>	+	+					+	+							+					✓	2
<i>Mustelus asterias</i>	+		+	+			+							+	+					✓	1
<i>Mustelus mustelus</i>	+	+	+	+				+						+	+				+	✓	1
<i>Mustelus punctulatus</i>															+					✓	5
<i>Galeorhinus galeus</i>	+												+							✓	2
<i>Dalatias licha</i>				+																	1
<i>Oxynotus centrina</i>		+	+				+	+						+	+				+	✓	1
<i>Centrophorus uyato</i>			+	+												+			+	✓	1
<i>Squalus acanthias</i>	+	+	+	+			+	+						+	+				+	✓	1
<i>Squalus blainville</i>	+		+	+			+							+					+	✓	1
<i>Echinorhinus brucus</i>	+															+			+	✓	2
<i>Squatina oculata</i>																	+			✓	2
<i>Squatina squatina</i>	+	+	+				+							+	+					✓	1
<i>Tetronarce nobiliana</i>					+																13
<i>Torpedo marmorata</i>	+	+	+											+	+				+	✓	1
<i>Torpedo torpedo</i>	+	+								+											5
<i>Dipturus batis</i>	+	+						+							+					✓	5
<i>Dipturus oxyrinchus</i>	+					+												+		✓	2

<i>Leucoraja naevus</i>						+																2
<i>Raja asterias</i>					+							+										14
<i>Raja clavata</i>	+	+						+		+				+	+				+	✓		1
<i>Raja miraletus</i>	+	+													+				+	✓		1
<i>Raja montagui</i>					+																	14
<i>Raja radula</i>									+										+	✓		1
<i>Dasyatis pastinaca</i>		+						+						+	+				+	✓		1
<i>Dasyatis tortonesei</i>											+								+	✓		5
<i>Gymnura altavela</i>					+																	13
<i>Aetomylaeus bovinus</i>												+								✓		2
<i>Myliobatis aquila</i>	+	+												+	+				+	✓		1
Number of species per reference	17	13	10	9	4	3	8	8	1	3	1	2	1	12	14	2	1	1	16	25		

supporting data. Therefore, although listed in the most recent ichthyological checklist (Bilecenoğlu, 2024), *T. nobiliana* and *G. altavela* were excluded from our updated inventory.

The occurrence of *Raja asterias* in the SoM was first reported by Mater and Meriç (1996), again without information where the examined specimen was captured and without a voucher specimen stored for further inspection. In 2009, another purported specimen of *R. asterias* was recorded in the southwestern SoM (Torcu Koç *et al.*, 2012); although detailed collection data were provided (capture depth and coordinates, substratum type etc.), photographic evidence later revealed that the specimen was a misidentified juvenile *R. clavata* (Kabasakal *et al.*, 2025a). The occurrence record for *R. montagui* is also based on Mater and Meriç (1996) and does not provide the relevant information (e.g., original source reference) for this species either. Therefore, the only applicable criterion for SoM records of both *R. asterias* and *R. montagui* is exclusion Criterion 14, which is employed when a species' reported presence cited in a review publication is traceable to an original reference that, upon inspection, proves to have been based on synonymization or misidentification. Although *R. brachyura* was included in another recent elasmobranch inventory of the SoM (Artüz & Fricke, 2024), it does not appear in our updated list – not even as a 'questionable' or 'unconfirmed' species – due to its documented absence from Turkish seas (Kabasakal, 2002; Bilecenoğlu, 2024). Its inclusion in the aforementioned checklist may stem from misidentification of the polymorphic *R. clavata* (Criterion 14, Kovačić *et al.*, 2020), as emphasised by Kabasakal *et al.* (2025a).

Karadurmuş and Sarı (2024) recently reported *Mustelus punctulatus* from the SoM for the first time. However, their record was not supported by any morphological, genetic, or photographic evidence, aside from locality data. The principal diagnostic feature used to identify *M. punctulatus* is the presence of black spots on the body (Ebert *et al.*, 2021). Although Marino *et al.* (2018) note that black spots are almost exclusively associated with *M. punctulatus*, they also emphasise that their absence is not diagnostic of *M. mustelus*. Furthermore, Ebert and Stehmann (2013) pointed out that some *M. mustelus* specimens may also exhibit dark spotting. In this taxonomically challenging situation, a second diagnostic key feature for *M. punctulatus* is the presence of fringed posterior margins on the dorsal fins (Marino *et al.*, 2018). Bello *et al.* (2014) proposed a best-practice approach for avoiding unverified or unverifiable 'first records' in ichthyology, which entails the preparation and deposition of a voucher specimen in a curated collection, accompanied by photographs, meristic and morphometric data, and, where possible, a DNA sequence (barcode). This protocol was not followed for the *M. punctulatus* record reported by Karadurmuş and Sarı (2024). The same authors also reported a new occurrence of *Dipturus batis* in the SoM, based on a single specimen captured off the northern coast (Karadurmuş & Sarı, 2024). Aside from the locality data, no supporting material (e.g., morphometric data or photographs) was provided to substantiate this important record, contrary to the verification standards proposed by Bello *et al.* (2014). Although *D. batis* appears in the recent checklists of Bilecenoğlu (2024) and Artüz and Fricke (2024), its inclusion is based on two historical references (Deveciyan, 1915; Ninni, 1923). Notably, Kabasakal *et al.* (2024b) reported



Fig. 2: Examples of sharks and batoids occurring in the SoM: (a) *Hexanchus griseus*; (b) *Echinorhinus brucus*; (c) *Oxynotus centrina*; and (d) *Myliobatis aquila*. All specimens were held in a survival tank until examination and subsequently released alive (photos: Hakan Kabasakal & F. Saadet Karakulak).

Sl. 2: Primeri morskih psov in skatov, ki se pojavljajo v južnem delu Marmarskega morja: (a) *Hexanchus griseus*; (b) *Echinorhinus brucus*; (c) *Oxynotus centrina*; in (d) *Myliobatis aquila*. Vsi primerki so bili do pregleda shranjeni v rezervoarju za preživetje in nato živi izpušeni na prostost (fotografije: Hakan Kabasakal in F. Saadet Karakulak).

sympatric *D. oxyrinchus* almost at the same location where Karadurmuş and Sarı (2024) claimed *D. batis*. As *D. oxyrinchus* and *D. batis* are morphologically very similar (Ebert & Stehmann, 2013; Barone *et al.*, 2022), they may be confused by researchers not specialising in elasmobranchs. Because of ambiguous evidence, the presence of *M. punctulatus* and *D. oxyrinchus* in the SoM was classified as confirmed only under the lowest applicable criterion (Criterion 5; defined above). We hope that future studies will obtain additional specimens to clarify the occurrence of these elasmobranchs in the SoM, allowing their status to be assessed against more robust criteria.

Finally, a recent study showed that either a highly differentiated variety of *S. blainville* or an undescribed species within the *S. megalops* clade may occur in the SoM (Kabasakal *et al.*, 2024c). Given the uncertainties

surrounding several elasmobranch species – primarily arising from potential misidentifications by non-taxonomists or researchers without specialised expertise in elasmobranchology and species-specific polymorphism – a genetically informed revision of the elasmobranch fauna of the SoM is urgently needed. To summarise this section, as of 16 June 2025, the updated inventory of demersal elasmobranchs of the SoM consists of the following species: *Hexanchus griseus*, *Galeus melastomus*, *Scyliorhinus canicula*, *S. stellaris*, *Mustelus asterias*, *M. mustelus*, *M. punctulatus*, *Galeorhinus galeus*, *Oxynotus centrina*, *Centrophorus uyato*, *Squalus acanthias*, *S. blainville*, *Echinorhinus brucus*, *Squatina oculata*, *S. squatina*, *Torpedo marmorata*, *Dipturus batis*, *D. oxyrinchus*, *Raja clavata*, *R. miraletus*, *R. radula*, *Dasyatis pastinaca*, *D. tortonesei*, *Aetomylaeus bovinus*, and *Myliobatis aquila* (Tab. 2).

Dispersal/settlement success of demersal elasmobranch species

The PCA incorporating basic life history characteristics, first and last record dates, and bathymetric distribution (depth ranges) for the 32 demersal elasmobranch species recorded in the SoM (Tab. 2) explained 47.62% of the total variance (Fig. 3). In contrast, a second PCA considering only the species' first and last record dates and their common depth ranges in the SoM, accounted for a variance of approximately 61.34% (Fig. 3). While records from the SoM preceding 1990 show that elasmobranch

species were more common in the deep continental shelf (100–200 m depth) and in waters deeper than 200 m, since the 2000s, these species have mostly been recorded at depths of 0–100 m. The PCA model based solely on common depth ranges and recording dates showed a 13.72% difference in explained variance compared to the PCA including basic life history parameters – such as maximum size, size at sexual maturity, and generation length – of demersal elasmobranchs. This suggests that adaptability to shallower waters is a key factor influencing the spatial distribution of demersal elasmobranch species in the SoM.




Fig. 3: Plots of Principal Component Analysis (PCA) of demersal elasmobranch species of the SoM, based on (a) all available data, and (b) only the first and last record dates and the common depths at which individuals were recorded in the SoM. The three-letter abbreviations next to the red triangles represent FAO codes for cartilaginous fishes, shown alongside their corresponding species names: *Hexanchus griseus*, SBL; *Galeus melastomus*, SHO; *Scyliorhinus canicula*, SYC; *Scyliorhinus stellaris*, SYT; *Mustelus asterias*, SDS; *Mustelus mustelus*, SMD; *Mustelus punctulatus*, MPT; *Galeorhinus galeus*, GAG; *Dalatias licha*, SCK; *Oxynotus centrina*, OXY; *Centrophorus uyato*, CPU; *Squalus acanthias*, DGS; *Squalus blainville*, QUB; *Echinorhinus brucus*, SHB; *Squatina oculata*, SUT; *Squatina squatina*, AGN; *Tetronarce nobiliana*, TTO; *Torpedo marmorata*, TTR; *Torpedo torpedo*, TTV; *Dipturus batis*, RJB; *Dipturus oxyrinchus*, RJO; *Leucoraja naevus*, RJN; *Raja asterias*, JRS; *Raja clavata*, RJC; *Raja miraletus*, JAI; *Raja montagui*, RJM; *Raja radula*, JAR; *Dasyatis pastinaca*, JDP; *Dasyatis tortonesei*, JDO; *Gymnura altavela*, RGL; *Aetomylaeus bovinus*, MPO; *Myliobatis aquila*, MYL.

Sl. 3: Diagrami analize glavnih komponent (PCA) pridnenih vrst hrustančnic v Marmarskem morju, ki temeljijo na (a) vseh razpoložljivih podatkih in (b) le na datumu prvega in zadnjega zapisa ter skupnih globinah, na katerih so bili primerki osebkov zabeleženi v Marmarskem morju. Tri črkovne okrajšave poleg rdečih trikotnikov predstavljajo kode FAO za hrustančnice, prikazane poleg njihovih ustreznih imen vrst: *Hexanchus griseus*, SBL; *Galeus melastomus*, SHO; *Scyliorhinus canicula*, SYC; *Scyliorhinus stellaris*, SYT; *Mustelus asterias*, SDS; *Mustelus mustelus*, SMD; *Mustelus punctulatus*, MPT; *Galeorhinus galeus*, GAG; *Dalatias licha*, SCK; *Oxynotus centrina*, OXY; *Centrophorus uyato*, CPU; *Squalus acanthias*, DGS; *Squalus blainville*, QUB; *Echinorhinus brucus*, SHB; *Squatina oculata*, SUT; *Squatina squatina*, AGN; *Tetronarce nobiliana*, TTO; *Torpedo marmorata*, TTR; *Torpedo torpedo*, TTV; *Dipturus batis*, RJB; *Dipturus oxyrinchus*, RJO; *Leucoraja naevus*, RJN; *Raja asterias*, JRS; *Raja clavata*, RJC; *Raja miraletus*, JAI; *Raja montagui*, RJM; *Raja radula*, JAR; *Dasyatis pastinaca*, JDP; *Dasyatis tortonesei*, JDO; *Gymnura altavela*, RGL; *Aetomylaeus bovinus*, MPO; *Myliobatis aquila*, MYL.

Environmental implications

Table 3 compares the maximum recorded depths of demersal elasmobranchs in the SoM with records from other regions of the Mediterranean Basin. The table highlights notable differences in maximum recorded depths depending on the year of observation in SoM-specific studies. Specifically, in the early 1990s, elasmobranch species were recorded down to upper slope depths (≤ 500 m) throughout the SoM; however, since the mid-2000s, their bathymetric ranges have been restricted to shelf waters (see asterisked references in Tab. 3). The only exceptions to this narrowing of the bathymetry were observations of *E. brucus* and *G. melastomus* in the Tekirdağ Trench (northwestern SoM) at depths below 1,000 m. (Kabasakal *et al.*, 2005; Oral, 2010). Several elasmobranchs (marked with an “δ”) that elsewhere in the Mediterranean inhabit shelf waters as well as upper and lower slope depths (e.g., Follesa *et al.*, 2019; Ruiz-García *et al.*, 2023; Deval & Mutlu, 2024) have, in the SoM since the mid-2000s, had maximum recorded depths restricted to outer and even inner shelf waters (see asterisked references in Table 3). Daban *et al.* (2021) reported that the number of elasmobranch species in the SoM increases with depth, and that the highest catch per unit of effort (CPUE; kg/km²) for *H. griseus*, *M. asterias* and *S. acanthias* was recorded at depths between 100 and 200 m. This suggests a distribution largely restricted to outer shelf waters. They also observed increased CPUE for *M. mustelus*, *S. squatina*, and several batoids at depths of 20–50 m. Karadurmuş and Sarı (2024) reported a similar shelf-restricted bathymetric distribution for elasmobranchs in the region. A recent study by Kabasakal (2025) found that the spatial distribution and abundance (number/km²) of demersal elasmobranchs in the SoM are now restricted to the inner shelf at depths of 50–100 m. The author attributes this to vertical habitat compression, driven by deoxygenation and the emergence of dead zones in deep shelf and bathyal waters – a process particularly severe in the environmentally degraded eastern SoM.

Based on correlations between captures of *H. griseus* and environmental parameters, Kabasakal *et al.* (2024d) concluded that the recent increase in bluntnose sixgill shark captures in continental shelf waters is primarily associated with an annual rise in dissolved organic compounds (NO₂+NO₃ nitrogen and PO₄ phosphorus) in bathyal waters. They also found that deteriorating environmental conditions and progressive deoxygenation in the deep waters of the SoM coincide with a reduction in the capture depth of *H. griseus* on the continental shelf. Severe hypoxia in deep continental shelf waters (>100 m) and anoxia in the slope waters of the northeastern SoM drastically constrict the bathymetric and spatial

distribution of demersal elasmobranchs in the region. PCA shows that, since the 2000s, records of demersal elasmobranchs have been concentrated at depths shallower than 150 m, reflecting the ongoing mass migration of these species toward continental shelf areas. This suggests that ‘adaptability to shallows on the continental shelf’ is a key factor in determining the spatial distribution of these species in the SoM.

So what is the future of the demersal elasmobranchs still struggling to survive in the SoM today? According to Diaz and Rosenberg (2008), these species may recolonise their former habitats if dissolved oxygen (DO) levels and ecosystem functions improve, although they may not reach pre-hypoxia levels. Historical observations support the one-time suitability of deeper waters for demersal cartilaginous fish: *E. brucus* was recorded at 1,214 m in the Tekirdağ Trench in northwestern SoM in the early 2000s (Kabasakal *et al.*, 2005), and *G. melastomus* was sampled at similar depths in the same region in 2008 (Oral, 2010). Additionally, Gallo *et al.* (2019) noted that certain elasmobranch species, such as cat sharks (e.g., *Cephalurus cephalus*), can tolerate suboxic conditions and withstand hypoxia better than previously believed. The species list presented in the 1974 Environmental Impact Assessment (EIA) report prepared by the Hydrobiology Research Institute to provide biological information prior to the construction of a sewage system discharging into the Bosphorus Strait – the gateway between the SoM and the Black Sea – offers insight into the historical species richness of elasmobranchs in the SoM (Anonymous, 1974; Fig. 4). Although the taxonomy used in the report is no longer valid, it is still possible to identify many of the listed species, especially pelagic ones, that have been absent from the region for years.

The extreme rarity of large pelagic sharks in the SoM is both remarkable and alarming. Early ichthyological inventories (Ninni, 1923; Deveciyan, 1926; Anonymous, 1974) considered the following pelagic shark species to be seasonal visitors to or residents of the region: *Alopias vulpinus*, *Cetorhinus maximus*, *Carcharodon carcharias*, *Lamna nasus*, and *Prionace glauca*. In addition, *A. superciliosus* was first recorded in the SoM in 2007 (Kabasakal & Karhan, 2008), with a second specimen captured in 2011. Since then, no *A. superciliosus* specimens have been captured or sighted. Today, all pelagic sharks except *A. vulpinus* have been extirpated from the SoM, and even incidental captures or sightings of this remaining species are becoming increasingly rare. This decline coincides with the warming of the region: over the past 20 years, the annual average sea surface temperature (SST) in the seas of Türkiye has increased by 0.4–1.4 °C due to global warming and climate change, with the SoM being the second most affected after the Black Sea (Demircan, 2022).

Tab. 3: Comparison between maximum recorded depths for demersal elasmobranchs included in the updated SoM inventory and values recorded in various studies from other regions of the Mediterranean. Notes: [§] species observed at the surface, no maximum depth record available; ^μ record based on data from a commercial fisherman, no maximum depth record available; ^δ species with bathyal distribution in Mediterranean subregions; * study specific to SoM. Tab. 3: Primerjava med največjimi zabeleženimi globinami za pridnene hrustančnice, vključene v posodobljen popis vrst v Marmarskem morju, in vrednostmi, zabeleženimi v različnih študijah iz drugih regij Sredozemlja. Opombe: § - vrste, opažene na površini, ni na voljo zapisa o največji globini; μ - zapis temelji na podatkih komercialnega ribiča, ni na voljo zapisa o največji globini; δ - vrste z batialno razširjenostjo v sredozemskih podregijah; * študija, specifična za Marmarsko morje.

Species	Present Study	JICA (1993)*	Meriç (1995)*	Kabasakal et al. (2005)*	Oral (2010)*	Torcu Koç et al. (2012)*	Bilecenoğlu (2019)*	Follesa et al. (2019)	Daban et al. (2021)*	Ruiz-García et al. (2023)	Deval & Mutlu (2024)	Karadurmuş & Sari (2024)*
<i>Hexanchus griseus</i> ^δ	188	–	350	–	–	–	–	800	200	537	–	–
<i>Galeus melastomus</i> ^δ	140	500	350	–	1213	–	–	800	–	685	800	–
<i>Scyliorhinus canicula</i> ^δ	124	500	350	–	–	60	–	800	200	649	600	146
<i>Scyliorhinus stellaris</i>	–	–	–	–	–	–	–	200	200	–	–	146
<i>Mustelus asterias</i>	–	500	350	–	–	–	–	200	200	–	–	45
<i>Mustelus mustelus</i>	188	100	350	–	–	–	–	200	200	–	200	85
<i>Mustelus punctulatus</i>	–	–	–	–	–	–	–	200	–	–	–	146
<i>Galeorhinus galeus</i> ^δ	–	–	–	–	–	–	–	800	–	–	–	–
<i>Oxynotus centrina</i> ^δ	66	500	–	–	–	–	–	800	200	214	600	146
<i>Centrophorus uyato</i> ^δ	150	500	270	–	–	–	–	800	–	588	800	–
<i>Squalus acanthias</i> ^δ	140	200	350	–	–	–	–	800	200	–	–	146
<i>Squalus blainville</i> ^δ	140	500	350	–	–	–	–	800	200	–	600	–
<i>Echinorhinus brucus</i>	150	–	–	1214	–	–	–	–	–	–	–	–
<i>Squatina oculata</i> ^μ	–	–	–	–	–	–	–	–	–	–	–	–
<i>Squatina squatina</i>	–	–	–	–	–	–	–	–	50	–	–	47
<i>Torpedo marmorata</i>	116	200	–	–	–	45	–	200	200	257	300	85
<i>Dipturus batis</i>	–	–	–	–	–	–	–	–	–	–	–	68
<i>Dipturus oxyrinchus</i> ^δ	100	500	–	–	–	–	–	800	–	329	600	–
<i>Raja clavata</i> ^δ	188	500	–	–	–	45	–	800	50	329	500	146
<i>Raja miraletus</i>	95	–	–	–	–	–	–	200	–	–	200	85
<i>Raja radula</i>	25	–	–	–	–	–	–	200	–	–	–	–
<i>Dasyatis pastinaca</i>	84	200	–	–	–	–	–	200	50	133	200	146
<i>Dasyatis tortonesei</i>	67	–	–	–	–	–	–	–	–	–	–	–
<i>Aetomylaeus bovinus</i>	–	–	–	–	–	–	3	200	–	73	–	–
<i>Myliobatis aquila</i>	90	200	–	–	–	–	–	200	200	73	–	85

List of Fishes in the Project Area					
A. SELACHII	I	II	III	IV	Hab.
<i>Acanthias vulgaris</i> (Riss.)	+++	+	-	++	D
<i>Centrina vulpicula</i> (Bell.)	+	-	-	+	D
<i>Cetorhinus maximus</i> (L.)	-	-	-	+	P
<i>Chacharias glauca</i> (Gm.)	-	-	-	+	P
<i>Charcharodon charcharias</i> (L.)	-	-	-	+	P
<i>Charcharodon lamia</i> (Rond.)	-	-	-	+	P
<i>Dasyatis pastinaca</i> (L.)	+	-	-	+	D
<i>Mustelus mustelus</i> (L.)	+++	+	-	+++	D
<i>Mustelus vulgaris</i> (Mull-Hen)	+++	+	-	+++	D
<i>Myliobatis aquila</i> (L.)	++	+	-	++	D
<i>Raja batis</i> (L.)	+	+	-	+	D
<i>Raja clavata</i> (L.)	+++	+++	-	+++	D
<i>Raja mirelatus</i> (L.)	+	+	-	+	D
<i>Raja oculata</i> (Risso.)	+	+	-	+	D
<i>Raja punctata</i> (Risso.)	+	+	-	+	D
<i>Scyllium canicula</i> (L.)	+	+	-	+	D
<i>Scyllium stellare</i> (L.)	+	+	-	+	D
<i>Squatina squatina</i> (L.)	+	+	-	++	D
<i>Torpedo marmorata</i> (Risso.)	+	+	-	+	D
<i>Torpedo narke</i> (Risso.)	+	+	-	++	D
<i>Trygon pastinaca</i> (Cuv.)	+	-	-	-	D
- not present					
+ present but rare					
++ found frequently					
+++ abundant					

Fig 4: Pelagic and demersal elasmobranch species reported from the southern entrance of the Bosphorus Strait (northeastern SoM) in the Environmental Impact Assessment (EIA) report compiled by the Hydrobiology Research Institute (adapted from Anonymous, 1974).

Sl. 4: Pelaške in pridnene vrste hrustančnic, o katerih so poročali pri južnem vходу v Bosporsko ožino (severovzhodni del Marmarskega morja) v poročilu o presoji vplivov na okolje (EIA), ki ga je sestavil Raziskovalni inštitut za hidrobiologijo (prirejeno po Anonymous, 1974).

This rise in seawater temperatures has reduced the solubility of DO, causing elasmobranch species to become less tolerant of hypoxia, as their metabolic rates increase in parallel with warming (Breitburg *et al.*, 2018; Waller *et al.*, 2024). Vedor *et al.* (2021) suggested that the expanding oxygen minimum zones (OMZs) driven by climate change – (through shoaling of low-DO waters and rising SST) – will compress the habitat of pelagic sharks, such as the blue shark, *P. glauca*, forcing them into surface waters above the OMZs and thereby reducing their overall available habitat volume. Such dynamic is indeed described by the OMZ ‘habitat trap’ hypothesis. According to Sims (2019), the high oxygen demands of pelagic sharks compel them to retreat to normoxic zones when encountering oxygen-depleted waters of the open ocean (like those in the SoM). Consequently, the apparent absence of most pelagic shark species (except *A. vulpinus*) from the SoM today is likely due to regional deoxygenation and an increased rate of their incidental capture in the resulting OMZ habitat trap. It seems that decades of environmental degradation, marine pollution, and deoxygenation in the SoM have reduced the region’s elasmobranch fauna largely to demersal species. This highlights the potentially devastating impact of environmental degradation on marine fauna and biodiversity. A comparable depletion of large mesopredators such as *G. galeus* and *Carcharhinus plumbeus*, and top predators such as *A. vulpinus*, *L. nasus*, and *P. glauca*, has been recorded in the Adriatic Sea, where environmental variability, in addition to fisheries, has been suggested to impact elasmobranch abundance (Barausse *et al.*, 2014).

In conclusion, while overfishing was once considered the primary threat to chondrichthyans (sharks, rays, and chimaeras) (Stevens *et al.*, 2000), habitat degradation, pollution, and climate change are now recognised as additional major drivers of species loss in this group (Dulvy *et al.*, 2021). Based on the assessment of fish communities across several aquatic ecosystems worldwide, Moyle and Leidy (1992) concluded that most faunas are in serious decline, primarily due to commercial exploitation, habitat alteration, and pollution, and therefore require immediate protection. Pollom *et al.* (2024) reported of two South African cat shark species (*Haploblepharus*

fuscus and *H. kistnasamyi*) threatened not only by overfishing but also by coastal urbanisation, industrialisation, and increasing marine pollution. The combined pressures have led to both species being categorised as ‘Vulnerable’ on the Red List. The findings of Pollom *et al.* (2024) align with numerous other studies demonstrating the drastic effects of human activities on regional fish faunas, as also documented by Bin Aziz *et al.* (2021) and Yang *et al.* (2024). The assumption that shallow continental shelf areas may become a habitat trap for demersal elasmobranchs – forcing these out of deep waters and exposing them to higher fishing pressure – has now materialised in the SoM, as pointed out by Kabasakal *et al.* (2023b) and Waller *et al.* (2024). According to Akoğlu *et al.* (2024), the pattern traditionally observed in demersal fishery – years of overfishing followed by stock collapse and gradual recovery over decades – no longer holds true for demersal elasmobranchs in the SoM. Although Important Shark and Ray Areas (ISRAs) have recently been designated in the SoM (Jabado *et al.*, 2023) – that confirm the presence of 25 demersal elasmobranch species and encompass potential breeding grounds for various taxa (e.g., *S. canicula*, *O. centrina*, *S. blainvillei* and *M. aquila*) distributed across the region (Daban *et al.*, 2021; Kabasakal *et al.* 2024e, 2025b) – we propose that the entire SoM be declared an ISRA to protect the taxonomic diversity of these sensitive species in an area undergoing severe environmental degradation.

ACKNOWLEDGMENTS

The authors thank the crew of *R/V Yunus-S* for their efforts during the field work. This study is part of the following research projects: (1) Stock Assessment of Demersal Fish Species in the Eastern Marmara Sea”, which is funded by the Scientific Research Projects Coordination Unit of Istanbul University (Project No. 2714/51922); and (2) Integrated Marine Pollution Monitoring (DEN-İZ) 2023-2025 Programme” carried out by the Ministry of Environment, Urbanization and Climate Change / General Directorate of Environmental Impact Assessment, Permit and Inspection and coordinated by TUBITAK Marmara Research Center.

PRIDNE NE HRUSTANČNICE V MARMARSKEM MORJU: POSODOBLJENA INVENTARIZACIJA, TAKSONOMSKA VPRAŠANJA IN OKOLJSKE POSLEDICE

Hakan KABASAKAL

Istanbul University, Institute of Graduate Studies in Sciences, Fisheries Technologies and Management Program, Istanbul, Türkiye
WWF Türkiye, İstanbul, Türkiye
e-mail: kbasakal.hakan@gmail.com

F. Saadet KARAKULAK

Department of Fisheries Technologies and Management, Faculty of Aquatic Sciences, İstanbul University, İstanbul, Türkiye

POVZETEK

Pričujoča raziskava temelji na vzorcih 16 vrst hrustančnic (devet vrst morskih psov in sedem vrst skatov), pridobljenih med letoma 2014 in 2024 v Marmarskem morju (SoM). V istem obdobju so drugi raziskovalci na istem območju zabeležili še dodatnih devet vrst (šest vrst morskih psov in tri vrste skatov). V zadnjih 100 letih je bilo na območju SoM tako zabeleženih 32 vrst pridnenih hrustančnic (16 vrst morskih psov in 16 vrst skatov), vendar uporaba meril za „potrjeno ali nepotrjeno prisotnost na podlagi dokazov“ trenutno število zmanjša na 25 (15 vrst morskih psov, 10 vrst skatov). Analiza glavnih komponent (PCA) kaže, da je prilagodljivost plitvejšim vodam zdaj ključnega pomena za prostorsko porazdelitev pridnenih hrustančnic v morskem okolju. Ta premik jasno kaže, kako dramatično so degradacija okolja, onesnaževanje morja in deoksigenacija v zadnjih 40 letih zmanjšali favno hrustančnic v regiji, predvsem pridnenih vrst, kar poudarja potencialno uničujoč vpliv takšne degradacije na biotsko raznovrstnost.

Ključne besede: Elasmobranchii, preživetje, ohranjanje, inventarizacija, Marmarsko morje, Turčija

REFERENCES

- Akoğlu, E., İ. Saygu & N. Demirel (2024):** Decadal changes in the Sea of Marmara indicate degraded ecosystem conditions and unsustainable fisheries. *Front. mar. sci.*, 11 (1412656). <https://doi.org/10.3389/fmars.2024.1412656>.
- Anonymous (1974):** Biological information for sewage disposal in the Bosphorus, Hydrobiological Research Institute, Project I/1974.
- Anonymous (2017):** MEDITS-Handbook, Version n. 9, MEDITS Working Group, 106 pp.
- Artüz, M.L. & R. Fricke (2024):** The marine cartilaginous fishes and sturgeons of the Sea of Marmara: an updated and annotated checklist. *Zootaxa*, 5501, 531–541. <https://doi.org/10.11646/zootaxa.5501.4.3>.
- Barausse, A., V. Corrales, A. Curkovic, L. Finotto, E. Riginella, E. Visentin & C. Mazzoldi (2014):** The role of fisheries and the environment in driving the decline of elasmobranchs in the northern Adriatic Sea. *ICES J. Mar. Sci.*, 71, 1593–1603. doi:10.1093/icesjms/fst222.
- Barone, M., C. Mazzoldi & F. Serena (2022):** Sharks, rays and chimaeras in Mediterranean and Black Seas – Key to identification. FAO, Rome, 87 pp. <https://doi.org/10.4060/cc0830en>.
- Bello, G., R. Causse, L. Lipej & J. Dulčić (2014):** A proposed best practice approach to overcome unverified and unverifiable “first records” in ichthyology. *Cybiu*, 38, 9–14.
- Benli, H.A., B. Cihangir & K.C. Bizsel (1993):** A new record for the Sea of Marmara; (Family: Squalidae) *Centrophorus granulosus* (Bloch & Schneider, 1801). *Turk. J. Zool.*, 17, 133–135.
- Beşiktepe, Ş.T., H.İ. Sur, E. Özsoy, M.A. Latif, T. Oğuz & Ü. Ünlüata (1994):** The circulation and hydrography of the Marmara Sea. *Prog. Oceanogr.*, 34, 285–334. [https://doi.org/10.1016/0079-6611\(94\)90018-3](https://doi.org/10.1016/0079-6611(94)90018-3).
- Bilecenoğlu, M. (2019):** First record of *Aetomylaeus bovinus* (Geoffroy St. Hilaire, 1817) Elasmobranchii: Myliobatidae, from the Sea of Marmara. *J. Black Sea/Medit.*, 25, 182–187.
- Bilecenoğlu, M. (2024):** Diversity of fishes along the coast of Türkiye. *Turk. J. Zool.*, 48, 589–616. <https://doi.org/10.55730/1300-0179.3197>.
- Bilecenoğlu, M., E. Taşkavak, S. Mater & M. Kaya (2002):** Checklist of the marine fishes of Turkey. *Zootaxa*, 113, 1–194.
- Bilecenoğlu, M., M. Kaya, B. Cihangir & E. Çiçek (2014):** An updated checklist of the marine fishes of Turkey. *Turk. J. Zool.*, 38, 901–929. <https://doi.org/10.3906/zoo-1405-60>.
- Bin Aziz, S., N.A. Hasan, M.R. Mondol, M. Alam & M.M. Haque (2021):** Decline in fish species diversity due to climatic and anthropogenic factors in Hakaluki Haor, an ecologically critical wetland in northeast Bangladesh. *Heliyon*, 7, e05861. <https://doi.org/10.1016/j.heliyon.2020.e05861>.
- Breitbart, D., L.A. Levin, A. Oschlies, M. Grégoire, F.P. Chavez, D.J. Conley, V. Garçon, D. Gilbert, G. Gutiérrez, K. Isensee, G.S. Jacinto K.E. Limburg, I. Montes, S.W.A. Naqvi, G.C. Pitcher, N.N. Rabalais, M.R. Roman, K.A. Rose, B.A. Seibel, M. Telszewski, M. Yasuhara & J. Zhang (2018):** Declining oxygen in the global ocean and coastal waters. *Science* 359, eaam7240. doi: 10.1126/science.aam7240.
- ÇŞİDB, TÜBİTAK-MAM (2021):** Denizlerde Bütünleşik Kirlilik İzleme Programı 2014-2019 Marmara Denizi Özet Raporu, TÜBİTAK-MAM Matbaası, Kocaeli. [in Turkish].
- Daban, İ.B., A. İşmen, M. Şirin, C.Ç. Yığın & M. Arslan İhsanoğlu (2021):** Analysis of Demersal Fish Fauna off the Sea of Marmara, Turkey. *COMU-JMSF.*, 4, 20–31. <https://doi.org/10.46384/jmsf.912403>.
- Demircan, M. (2022):** Climate Change's Impact on Turkish Seas' Temperature and Aquaculture. *JENAS.*, 4, 96–108. <https://doi.org/10.53472/jenas.1096917>.
- Demirel, N., M. Zengin & A. Ulman (2020):** First large-scale eastern Mediterranean and Black Sea stock assessment reveals a dramatic decline. *Front. mar. sci.*, 7, 103. <https://doi.org/10.3389/fmars.2020.00103>.
- Deval, M.C. & E. Mutlu (2024):** Spatio-temporal density of the demersal chondrichthyes assemblage in an upper bathyal of the eastern Mediterranean Sea. *Mar. Biodivers.*, 54, 31. <https://doi.org/10.1007/s12526-024-01423-x>.
- Devedjian, K. (1915):** Balık ve Balıkçılık. İstanbul: Düyun-u Umumiye-i Osmaniye Varidat-ı Mahsusa İdare-i Merkeziyesi Matbaası. [in Turkish].
- Deviciyan, K. (1926):** Pêche et Pêcheries en Turquie. Imprimerie de l'Administration de la Dette Publique Ottomane, Istanbul, 480 pp.
- Diaz, R.J. (2016):** Anoxia, hypoxia, and dead zones. In: M.J. Kennish (Ed.). *Encyclopedia of Estuaries*, Springer Science+Business Media, Dordrecht, pp. 19–29. <http://dx.doi.org/10.1007/978-94-017-8801-4>.
- Diaz, R.J. & R. Rosenberg (2008):** Spreading dead zones and consequences for marine ecosystems. *Science*, 321, 926–929. <http://dx.doi.org/10.1126/science.1156401>.
- Dulvy N.K., N. Pacoureau, C.L. Rigby, R. A. Pollock, R.W. Jabado, D.A. Ebert, B. Finucci, C.M. Pollock, J. Cheok, D.H. Derrick, K.B. Herman, C.S. Sherman, W.J. VanderWight, J.M. Lawson, R.H.L. Walls, J.K. Carlson, P. Charvet, K.K. Bineesh, D. Fernando, G.M. Ralph, J.H. Matsushiba, C. Hilton-Taylor, S.V. Fordham & C.A. Simpfendorfer (2021):** Overfishing drives over one-third of all sharks and rays toward a global extinction crisis. *Curr. Biol.*, 31, 1–15. <https://doi.org/10.1016/j.cub.2021.08.062>.

- Ebert, D.A. & M.F.W. Stehmann (2013):** Sharks, batoids, and chimaeras of the North Atlantic, FAO Species Catalogue for Fishery Purposes. No. 7. FAO, Rome, 523 pp.
- Ebert, D.A., M. Dando & S. Fowler (2021):** Sharks of the World: A Complete Guide. Princeton University Press, Princeton, 608 pp.
- Ellis, J.R. & N.K. Dulvy (2016):** *Leucoraja naevus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T161626A16527841.
- Ellis, J.R., S.R. McCully Phillips & F. Poisson (2016a):** A review of capture and post-release mortality of elasmobranchs. J. Fish Biol., 90, 653–722. <https://doi.org/10.1111/jfb.13197>.
- Ellis, J.R., F. Serena, C. Mancusi, F. Haka, G. Morey, J. Guallart & T. Schembri (2016b):** *Scyliorhinus stellaris* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T161484A16527873.
- Ellis, J.R., A. Soldo, M. Dureuil & S. Fordham (2016c):** *Squalus acanthias* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T91209505A16527761.
- Ellis, J.R., A. Abella, F. Serena & M.F.W. Stehmann (2016d):** *Dipturus oxyrinchus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T63100A16527733.
- Ellis, J.R., N.K. Dulvy & F. Serena (2016e):** *Raja clavata* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T39399A103113598.
- Ellis, J.R., F. Serena & N. Dulvy (2016f):** *Raja montagui* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T63146A16564906.
- Ellis, J., S.R. McCully-Phillips, D. Sims, D. Derrick, J. Cheok & N.K. Dulvy (2021):** *Dipturus batis*. The IUCN Red List of Threatened Species 2021: e.T203364219A203375487. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T203364219A203375487.en>.
- Eryılmaz, L. & N. Meriç (2005).** Review of fish fauna of the Sea of Marmara. J. Black Sea/Medit., 11, 153–178.
- FAO, ACCOBAMS (2018):** Good practice guide for the handling of sharks and rays caught incidentally in Mediterranean pelagic longline fisheries. <http://www.fao.org/3/i9152en/I9152EN.pdf>.
- Farrell, E.D., S. McCully, N.K. Dulvy, C. Mancusi & J.R. Ellis (2016a):** *Mustelus asterias* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T39357A16527992.
- Farrell, E.D. & N.K. Dulvy (2016b):** *Mustelus mustelus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T39358A16527988.
- Ferretti, F. & E. Buscher (2016):** *Echinorhinus brucus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T41801A90728641.
- Ferretti, F., G. Morey, F. Serena, C. Mancusi, R.P. Coelho, M. Seisay, F. Litvinov & E. Buscher (2016a):** *Squatina oculata* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T61418A16570000.
- Ferretti, F., G. Morey, F. Serena, C. Mancusi, S.L. Fowler, F. Dipper & J.R. Ellis (2016b):** *Squatina squatina* (Mediterranean assessment) (errata version published in 2016). The IUCN Red List of Threatened Species 2016: e.T39332A101695971.
- Finucci, B., A. Barnett, K.K. Bineesh, J. Cheok, C.F. Cotton, Dharmadi, K.J. Graham, D.W. Kulka, F.C. Neat, N. Pacoureaux, C.L. Rigby, S. Tanaka & T.I. Walker (2020):** *Hexanchus griseus*. The IUCN Red List of Threatened Species 2020: e.T10030A495630. <https://dx.doi.org/10.2305/IUCN.UK.2020-3.RLTS.T10030A495630.en>.
- Finucci, B., D. Derrick, J. Dossa & A.B. Williams (2021):** *Tetronarce nobiliana*. The IUCN Red List of Threatened Species 2021: e.T116861529A116861706. <https://dx.doi.org/10.2305/IUCN.UK.2021-2.RLTS.T116861529A116861706.en>.
- Follesa, M.C., M.F. Marongiu, W. Zupa, A. Bel-lodi, A. Cau, R. Cannas, F. Colloca, M. Djurović, I. Isajlović, A. Jadaud, C. Manfredi, A. Mulas, P. Peristeraki, C. Porcu, S. Ramirez-Amaro, F. S. Jiménez, F. Serena, L. Sion, I. Thasitis, A. Cau & P. Carbonara (2019):** Spatial variability of Chondrichthyes in the northern Mediterranean. In: Mediterranean demersal resources and ecosystems: 25 years of MEDITS trawl surveys (M.T. Spedicato, G. Tserpes, B. Mérigot & E. Massutí, eds.) Sci. Mar., 83S1, December 2019, 81–100. ISSN-L 0214-8358 <https://doi.org/10.3989/scimar.04998.23A>.
- Gallo, N.D., L.A. Levin, M. Beckwith & J.P. Barry (2019):** Home sweet suboxic home: remarkable hypoxia tolerance in two demersal fish species in the Gulf of California. Ecology, 100 (3), e02539. <https://doi.org/10.1002/ecy.2539>.
- Geldiay, R. (1969):** İzmir Körfezinin Başlıca Balıkları ve Muhtemel İnvasyonları. Ege Üniversitesi Matbaası, İzmir. [in Turkish].
- GFCM (2018):** GFCM Data Collection Reference Framework (DCRF). Version: 23.2., FAO, 190 p.
- Guallart, J. & R. Walls (2015):** *Centrophorus uyato* (Europe assessment). The IUCN Red List of Threatened Species 2015: e.T41745A72821789.
- Gül, G. & N. Demirel (2021):** Evaluation of the comprehensive feeding strategy and trophic role of overexploited mesopredator species in the Sea of Marmara (northeastern Mediterranean). Estuar. Coast. Shelf Sci., 259, 107448. <https://doi.org/10.1016/j.ecss.2021.107448>.

- Hammer, Ø., D.A.T. Harper & P.D. Ryan (2001):** Past: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontol. electronica*, 4, 9 pp. http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- Jabado, R.W. & D. Derrick (2021):** *Dasyatis tortonesei*. The IUCN Red List of Threatened Species 2021: e.T18907453A68783115.
- Jabado, R.W., E. García-Rodríguez, P.M. Kyne, R. Charles, A.H. Armstrong, J. Bortoluzzi, T.L. Mouton, A. GonzalezPestana, A. Battle-Morera, C. Rohner & G. Notarbartolo di Sciara (2023):** Mediterranean and Black Seas: A regional compendium of Important Shark and Ray Areas, IUCN SSC Shark Specialist Group, Dubai.
- JICA (1993):** Marmara, Ege ve Akdenizde demersal balıkçılık kaynakları sörvey raporu. Japonya Uluslararası İşbirliği Ajansı (JICA), T.C. Tarım ve Köyişleri Bakanlığı Tarımsal Üretim ve Geliştirme Genel Müdürlüğü, Ankara, 579 p. [in Turkish].
- Kabasakal, H. (2002):** Elasmobranch species of the seas of Turkey. *Ann. Ser. Hist. Nat.*, 12, 15-22.
- Kabasakal, H. (2016):** Status of Cartilaginous Fishes in the Sea of Marmara. In: E. Özsoy, M.N. Çağatay, N. Balkis & B. Öztürk (Eds.). *The Sea of Marmara - Marine Biodiversity, Fisheries, Conservation and Governance*. İstanbul, Turkish Marine Research Foundation No: 42, pp. 903-918.
- Kabasakal, H. (2022):** Marmara Denizi'nde köpekbalıkları: İçdenizde yaşayan türlere güncel bilgiler ışığında bir bakış. In: B. Öztürk, H.A. Ergül, A.C. Yalçiner, H. Öztürk & B. Salihoğlu (Eds.). *Marmara Denizi 2022 Sempozyumu Bildiriler Kitabı, Türk Deniz Araştırmaları Vakfı*, Publication No: 63, 8-9 Ocak 2022 İstanbul, 337-343. [in Turkish].
- Kabasakal, H. (2025):** Effects of dissolved oxygen on distribution of chondrichthyes in eastern Sea of Marmara. PhD Thesis, İstanbul University, Institute of Graduate Studies in Sciences, Fisheries Technologies and Management Program, İstanbul, 165 pp.
- Kabasakal, H. & S.Ü. Karhan (2008):** On the occurrence of the bigeye thresher shark, *Alopias superciliosus* (Chondrichthyes: Alopiidae), in Turkish waters. *Mar. Biodivers. Rec.*, 1, e69, 1-3. <https://doi.org/10.1017/S1755267207007452>.
- Kabasakal, H. & K.F. Türetken (2021):** Photographic documentation of tope shark, *Galeorhinus galeus* (Linnaeus, 1758), in İstanbul Strait: Reoccurrence of a rare shark in troubled waters. *J. Black Sea/Medit.*, 27, 372-378.
- Kabasakal, H. & F.S. Karakulak (2024):** A Review of Sharks and Batoids of the Sea of Marmara: Current Inventory, Conservation Issues and Future Projections. In: M. İşinibilir, A.E. Kideyş & A. Malej (Eds.). *Ecological Changes in the Sea of Marmara*, İstanbul University Press, E-ISBN 978-605-07-1633-7. pp. 819-846. doi: 10.26650/B/LSB21LSB37.2024.023.025.
- Kabasakal, H., M.İ. Öz, S.Ü. Karhan, Z. Çaylarbaşı & U. Tural (2005):** Photographic evidence of the occurrence of bramble shark, *Echinorhinus brucus* (Bonnaterre, 1788) (Squaliformes: Echinorhinidae) from the Sea of Marmara. *Annales, Ser. Hist. Nat.*, 15, 51-56.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2023a):** Occurrence of deep-sea Squaliform sharks, *Echinorhinus brucus* (Echinorhinidae) and *Centrophorus uyato* (Centrophoridae), in Marmara shelf waters. *Annales, Ser. Hist. Nat.*, 33, 27-36. <https://doi.org/10.19233/ASHN.2023.05>.
- Kabasakal, H., S. Sakınan, L. Lipej & D. Ivajnsiç (2023b):** A preliminary life history traits analysis of sharks in the Sea of Marmara (Türkiye), where deoxygenation and habitat deterioration are raising concerns. *Aquat. Res.*, 6, 72-82. <https://doi.org/10.3153/AR23008>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2024a):** Where have you been hiding? Recurrence of *Squatina oculata* in the Sea of Marmara, with a review of the captures in Turkish seas. *Aquat. Sci. Eng.*, 39, 200-205. <https://doi.org/10.26650/ASE20241406066>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2024b):** Occurrence of longnosed skate, *Dipturus oxyrinchus*, in the Sea of Marmara. *Annales, Ser. Hist. Nat.*, 34, 221-228. <https://doi.org/10.19233/ASHN.2024.27>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2024c):** Is there a third species of the genus *Squalus* (Squalidae) present in the Marmara Sea, or just an extreme case of intraspecific variability of *Squalus blainville*? *J. Ichthyol.*, 64, 727-736. <https://doi.org/10.1134/S0032945224700462>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2024d):** Impact of fishing capacity and environmental parameters on landings of *Hexanchus griseus* in the Sea of Marmara, *Annales, Ser. Hist. Nat.*, 34, 257-272. <https://doi.org/10.19233/ASHN.2024.31>.
- Kabasakal, H., U. Uzer, O. Gönülal & F.S. Karakulak (2024e):** Bioecological lessons learned from the neonate longnose spurdogs *Squalus blainville* (Squaliformes: Squalidae) suggest a potential nursery ground in the Marmara Sea. *Acta Adriatica*, 65. <https://doi.org/10.32582/aa.65.2.2>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2025a):** Did *Raja asterias* ever occur in the Sea of Marmara or were previous records just the cases of misidentified *Raja clavata*? *Turk. J. Zool.*, 49, 92-102. <https://doi.org/10.55730/1300-0179.3215>.
- Kabasakal, H., U. Uzer & F.S. Karakulak (2025b):** Captures of maturing and gravid female angular roughsharks, *Oxynotus centrina* (Oxynotidae), in the Sea of Marmara. *J. Ichthyol.*, 65, 282-290. <https://doi.org/10.1134/S0032945224701029>.

- Karadurmuş, U. & M. Sarı (2024):** Distribution and diversity of elasmobranchs in the Sea of Marmara: a 2023 status report. *Aquat. Conserv.: Mar. Freshw.*, 34, e4121. <https://doi.org/10.1002/aqc.4121>.
- Kocataş, A., T. Koray, M. Kaya, & Ö.F. Kara (1993)** Review of the fishery resources and their environment in the Sea of Marmara. *Studies and Reviews No: 64, Fisheries and Environmental Studies in the Black Sea System*, FAO, Rome, pp. 87-143.
- Kovačić, M., L. Lipej & J. Dulčić (2020):** Evidence approach to checklists: critical revision of the checklist of the Adriatic Sea fishes. *Zootaxa*, 4767, 001-055. <https://doi.org/10.11646/zootaxa.4767.1.1>.
- Last P.R., B. Séret, M.F.W. Stehmann & S. Weigmann (2016):** Skates - Family Rajidae, In: (W.T. White, M.R. de Carvalho, B. Séret, M.F.W. Stehmann, G.J.P. Naylor & L. Marshall) (Eds.). *Rays of the world*, CSIRO, Clayton South, pp. 204-363.
- Mancusi, C., G. Morey & F. Serena (2016):** *Raja radula*. The IUCN Red List of Threatened Species 2016: e.T161339A16527984.
- Mantıkçı, M., H. Örek, M. Yücel, Z. Uysal, S. Arkin & B. Salihoğlu (2022):** Marmara Denizi'nin güncel oksijen durumu ve müsilajın etkisi, In: B. Öztürk, H.A. Ergül, A.C. Yalçiner, H. Öztürk & B. Salihoğlu (Eds.). *Marmara Denizi 2022 Sempozyumu Bildiriler Kitabı*, Türk Deniz Araştırmaları Vakfı, Publication No: 63, 8-9 Ocak 2022 İstanbul, pp. 18-24. [in Turkish].
- Marino, I.A.M., L. Finotto, F. Colloca, M. Di Lorenzo, M. Gristina, E.D. Farrell, L. Zane & C. Mazzoldi (2018):** Resolving the ambiguities in the identification of two smooth-hound sharks (*Mustelus mustelus* and *Mustelus punctulatus*) using genetics and morphology. *Mar. Biodiv.*, 48, 1551-1562. <https://doi.org/10.1007/s12526-017-0701-8>.
- Mater, S. & N. Meriç (1996):** Deniz balıkları - Pisces. In: Türkiye Omurgalıları Tür Listesi (A. Kence & C.C. Bilgin, eds.). TÜBİTAK, Ankara, 133-172. [in Turkish].
- McCully, S., M. Dureuil & E.D. Farrell (2016):** *Galeorhinus galeus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T39352A16527949.
- Meriç, N. (1995):** A study on existence of some fishes on the continental slope of the Sea of Marmara. *Turk. J. Zool.*, 19, 191-198.
- Moyle, P.B. & R.A. Leidy (1992):** Loss of biodiversity in aquatic ecosystems: evidence from fish faunas. In: P.L. Fiedler & S.K. Jain (Eds.). *Conserv. Biol.* Springer, Boston, MA, pp. 127-169. https://doi.org/10.1007/978-1-4684-6426-9_6.
- Ninni, E. (1923):** Primo contributo allo studio dei pesci e della pesca nelle acque dell'Impero Ottomano. *Premiate Officine Grafiche Carlo Ferrari*, Venezia.
- Notarbartolo di Sciara, G., F. Serena, N. Ungaro, F. Ferretti, S. Pheeha, B. Human, S. McCully & E. Buscher (2016):** *Torpedo marmorata* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T161328A16527917.
- Oral, M. (2010):** A preliminary study on feeding habits of blackmouth catshark *Galeus melastomus*, Rafinesque, 1810, sampled from the Sea of Marmara. p. 312-316. In: B. Öztürk (Ed.). *Marmara Denizi 2010 Sempozyumu*, Bakırköy Yunus Emre Kültür Merkezi, İstanbul, 25-26 September 2010. *Bildiriler Kitabı*. Türk Deniz Araştırmaları Vakfı İstanbul. [in Turkish].
- Pollom, R.A., J. Cheok, N. Pacoureaux, K.S. Gledhill, P.M. Kyne, D.A. Ebert, R.W. Jabado, K.B. Herman, R.H. Bennett, C. da Silva, S. Fernando, B. Kuguru, R.W. Leslie, M.E. McCord, M. Samoilys, H. Winker, S.T. Fennessy, C.M. Pollock, C.L. Rigby & N.K. Dulvy, N.K. (2024):** Overfishing and climate change elevate extinction risk of endemic sharks and rays in the southwest Indian Ocean hotspot. *PLoS ONE*, 19, e0306813. <https://doi.org/10.1371/journal.pone.0306813>.
- Rhesis Erazi, R.A. (1942):** Marine fishes found in the Sea of Marmara and in the Bosphorus. *İstanbul Üniversitesi Fen Fakültesi Mecmuası, Seri B*, 7, 103-115.
- Rincón-Díaz, M.P., S.J. Pittman, I. Arismendi & S.S. Heppell (2018):** Functional diversity metrics detect spatio-temporal changes in the fish communities of a Caribbean marine protected area. *Ecosphere*, 9, e02433. <https://doi.org/10.1002/ecs2.2433>.
- Ruiz-García, D., J.A. Raga, D. March, A.I. Colmenero, F. Quattrocchi, J.B. Company, L. Recasens & C. Barría (2023):** Spatial distribution of the demersal chondrichthyan community from the western Mediterranean trawl bycatch. *Front. mar. sci.*, 10:1145176. <https://doi.org/10.3389/fmars.2023.1145176>.
- Salihoğlu, B., B.F. Salihoğlu, D. Tezcan, S. Tuğrul, E. Akoğlu, K. Özkan & S. Arkin (2022):** Sea of Marmara ecosystem under multiple stressors and rehabilitation plans, In: B. Öztürk, H.A. Ergül, A.C. Yalçiner, H. Öztürk & B. Salihoğlu (Eds.). *Marmara Denizi 2022 Sempozyumu Bildiriler Kitabı*, Türk Deniz Araştırmaları Vakfı, Yayın No: 63, 8-9 Ocak 2022 İstanbul, pp. 49-53. [in Turkish].
- Saygu, İ. & M.C. Deval (2014):** The post-release survival of two skate species discarded by bottom trawl fisheries in Antalya Bay, Eastern Mediterranean. *TrJFAS*, 14, 947-953. https://doi.org/10.4194/1303-2712-v14_4_14.
- Saygu, İ., E. Akoğlu, G. Gül, D. Bedikoğlu & N. Demirel (2023):** Fisheries impact on the Sea of Marmara ecosystem structure and functioning during the last three decades. *Front. mar. sci.*, 9, 1076399. <https://doi.org/10.3389/fmars.2022.1076399>.

- Serena, F. (2005):** Field identification guide to the sharks and rays of the Mediterranean and Black Sea. FAO Species Identification Guide for Fishery Purposes. FAO, Roma.
- Serena, F., J. Ellis, A. Abella, C. Mancusi, F. Haka, J. Guallart, N. Ungaro, R.P. Coelho, T. Schembri & M. Kirsteen (2015):** *Scyliorhinus canicula* (Europe assessment). The IUCN Red List of Threatened Species 2015: e.T161307554A201955962. <https://dx.doi.org/10.2305/IUCN.UK.2015-1.RLTS.T161307554A201955962.en>.
- Serena, F., G. Notarbartolo di Sciara, & N. Ungaro (2016a):** *Torpedo torpedo* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T161397A16527797.
- Serena, F., A. Abella, R.H.L. Walls & N.K. Dulvy (2016b):** *Raja asterias* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T63120A16527773.
- Serena, F., C. Mancusi, G. Morey & J.R. Ellis (2016c):** *Dasyatis pastinaca* (Mediterranean assessment) (errata version published in 2016). The IUCN Red List of Threatened Species 2016: e.T161453A97841681.
- Serena, F., J. Holtzhausen, D.A. Ebert & C. Mancusi (2016d):** *Myliobatis aquila* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T161569A16527996.
- Silveira, L.F., D. de Mello Beisiegel, F.F. Curcio, P.H. Valdujo, M. Dixo, V.K. Verdade, G. M.T. Mattox & P.T.M. Cunningham (2010):** What use do fauna inventories serve? *Estud. Av.*, 24, 173–207.
- Sims, D.W. (2019):** The significance of ocean deoxygenation for elasmobranchs. In: Laffoley, D. and Baxter, J.M. (Eds.). *Ocean deoxygenation: everyone's problem: causes, impacts, consequences and solutions*. IUCN Global Marine and Polar Programme. pp. 431–448. <https://doi.org/10.2305/IUCN.CH.2019.13.en>.
- Soldo, A. & J. Guallart (2016):** *Oxynotus centrina* (Mediterranean assessment) (errata version published in 2016). The IUCN Red List of Threatened Species 2016: e.T63141A97834254.
- Soldo, A., M.N. Bradai, E. Buscher, D.A. Ebert, F. Serena & C. Mancusi (2016):** *Squalus blainville* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T169229923A200982471.
- Stauffer Jr., J.R. & P.M. Kocovsky (2007):** Exploring links between systematics and fisheries management. *Trans. Am. Fish. Soc.*, 136, 1122–1125. <https://doi.org/10.1577/T06-166.1>.
- Stevens, J.D., R. Bonfil, N.K. Dulvy & P.A. Walker (2000):** The effects of fishing on sharks, rays, and chimaeras (chondrichthyans), and the implications for marine ecosystems. *ICES Journal of Marine Science*, 57(3), 476–494. <https://doi.org/10.1006/jmsc.2000.0724>.
- Torcu Koç, H., F. Üstün, Z. Erdoğan & L. Artüz (2012):** Species composition of benthic fish fauna in the Sea of Marmara, Turkey. *J. Appl. Ichthyol.*, 37, 303–307. <https://doi.org/10.1111/j.1439-0426.2012.02037.x>.
- Uysal, A., A. Yüksek & E. Okuş (1996):** Marmara Denizi'nde ender rastlanan üç elasmobranchii türü ve dağılımları: (*Raja naevus* Müller & Henle, 1841; *Raja oxyrinchus* Linnaeus, 1758; *Galeus melastomus* Rafinesque, 1809), Abstracts of 13th National Congress of Biology, 17–20 September 1996, 28. [in Turkish].
- Ünlüata, Ü. & E. Özsoy (1988):** Deep water renewals and oxygen deficiency in the Sea of Marmara. *Rapp. Comm. int. Mer Medit.*, 31, 193.
- Vaquier-Sunyer, R. & C.M. Duarte (2008):** Thresholds of hypoxia for marine biodiversity. *Proc. Natl. Acad. Sci. U. S. A.*, 105, 15452–15457. <https://doi.org/10.1073/pnas.0803833105>.
- Vedor, M., N. Queiroz, G. Mucientes, A. Couto, I. da Costa, A. dos Santos, F. Vandeperre, J. Fontes, P. Afonso, R. Rosa, N.E. Humphries & D.W. Sims (2021):** Climate-driven deoxygenation elevates fishing vulnerability for the ocean's widest ranging shark. *eLife*, 10, e62508. <https://doi.org/10.7554/eLife.62508>.
- Villagra, D., N. Van Bogaert, B. Ampe, P. Walker & S.S. Uhlmann (2022):** Life-history traits of batoids (Superorder Batoidea) in the northeast Atlantic and the Mediterranean. *Rev. Fish Biol. Fish.*, 32(2), 473–495. <https://doi.org/10.1007/s11160-021-09695-3>.
- Waller, M.J., N.E. Humphries, F.C. Womersley, A. Loveridge, A.L. Jeffries, Y. Watanabe, N. Payne, J. Semmens, N. Queiroz, E.J. Southall & D.W. Sims (2024):** The vulnerability of sharks, skates, and rays to ocean deoxygenation: physiological mechanisms, behavioral responses, and ecological impacts. *J. Fish Biol.*, 105, 482–511. <https://doi.org/10.1111/jfb.15830>.
- Walls, R.H.L. & E. Buscher (2016):** *Aetomylaeus bovinus* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T60127A81163810.
- Walls, R.H.L. & J. Guallart (2016):** *Dalatias licha* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T6229A16527825.
- Walls, R.H.L., M. Vacchi, G. Notarbartolo di Sciara, F. Serena & N.K. Dulvy (2016):** *Gymnura altavela* (Mediterranean assessment). The IUCN Red List of Threatened Species 2016: e.T63153A16527909.
- WoRMS Editorial Board (2025):** World Register of Marine Species. Available from <https://www.marinespecies.org> at VLIZ. Accessed 2025-06-11. doi:10.14284/170.

Yaka, U. & R. Yüce (2006): The rough ray, *Raja radula* Delaroche, 1809 (Rajidae), new to the Sea of Marmara, Turkey. Zool. Middle East, 39, 112-114. <http://dx.doi.org/10.1080/09397140.2006.10638193>.

Yang, B., X. Qu, H. Liu, M. Yang, W. Xin, W. Wang & Y. Chen (2024): Urbanization reduces fish taxonomic and functional diversity while increases phylogenetic diversity in subtropical rivers. Sci. Total Environ., 908, 168178. <https://doi.org/10.1016/j.scitotenv.2023.168178>.

Yıldız, T., E. Yemişken, F.S. Karakulak, U. Uzer, C. Dalyan & I.K. Oray (2016): A new record of dasyatid fish from the Sea of Marmara: Tortonese's stingray, *Dasyatis tortonesei* Capapé, 1975 (Dasyatidae). J. Appl. Ichthyol., 32, 721-723. <https://doi.org/10.1111/jai.13087>.

Yılmaz, A., Ö. Baştürk, S. Tuğrul, C. Saydam & Ü. Ünlüata (1990): Marmara Denizi ve boğazlar sisteminin su dinamiği ve çevresel karakteristiği. 7 Bölümde İstanbul'un Çevre Sorunları ve Çözümleri Sempozyumu, 9-13 Nisan 1990, Türkiye Çevre Koruma ve Yeşillendirme Kurumu, pp. 231-261. [in Turkish].

Zengin, M., H. Polat, S. Kutlu, A.C. Dinçer, H. Güngör, M. Aksoy, C. Özgündüz, E. Karaaslan & Ş. Firidin (2004): Marmara Denizi'ndeki derin su pembe karidesi (*Parapenaeus longirostris* Lucas, 1846) balıkçılığın geliştirilmesi üzerine bir araştırma, Sonuç Raporu (TAGEM/HAYSUD/2001/09/2004), T.C. Tarım ve Köyişleri Bakanlığı Tarımsal Araştırmalar Genel Müdürlüğü. [in Turkish].