

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 35, 2025, 2





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 35, 2025, 2

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 2

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednica/Redattrice/Editor:

Martina Orlando-Bonaca

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Založnika/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 6. 12. 2025.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / *CONTENTS*SREDOZEMSKA HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
*MEDITERRANEAN SHARKS AND RAYS***Hakan KABASAKAL**

Analysis of Confirmed Shark Attacks
in the Eastern Mediterranean Sea and
the Sea of Marmara (1827–2025) 169
*Analiza potrjenih napadov morskih psov
v vzhodnem Sredozemskem morju
in Marmarskem morju (1827–2025)*

Alen SOLDO

Observations of a Juvenile Basking Shark
Cetorhinus maximus in the Adriatic Sea
Support the Hypothesis of a Distinct
Mediterranean Population 187
*Opazovanja mladega morskega psa
orjaka (Cetorhinus maximus)
v Jadranskem morju podpirajo
hipotezo o ločeni sredozemski
populaciji*

**Jacopo BERNARDI, Pero UGARKOVIĆ &
Ilija ČETKOVIĆ**

An Unusual Encounter with a Juvenile Kitefin
Shark, *Dalatias licha*, in Shallow Coastal
Waters of Montenegro (Adriatic Sea) 197
*Nenavadno srečanje z mladim primerkom
klinoplavutega morskega psa, Dalatias licha,
v plitvih obalnih vodah Črne gore
(Jadransko morje)*

**Cemal TURAN, Alen SOLDO, Servet A. DOĞDU,
Funda TURAN, Aysegül ERGENLER & Ali UYAN**

Identification of a Potential Nursery Ground
of the Spiny Butterfly Ray, *Gymnura altavela*,
in the Northeastern Mediterranean Sea, Türkiye 205
*Identifikacija potencialnih jaslčic za
metuljastega skata, Gymnura
altavela, v severovzhodnem
Sredozemskem morju,
Turčija*

**Alen SOLDO, Eleonora DE SABATA &
Simona CLO**

Recent Records of Critically Endangered
Common Guitarfish *Rhinobatos rhinobatos*
(Linnaeus, 1758) in the Northern
Mediterranean 213
*Nedavni zapiski o pojavljanju kritično
ogroženega navadnega goslaša
Rhinobatos rhinobatos
(Linnaeus, 1758) v
severnem Sredozemlju*

Hakan KABASAKAL & F. Saadet KARAKULAK

Demersal Elasmobranchs of the Sea of Marmara:
Updated Inventory, Taxonomic Issues and
Environmental Implications 221
*Pridnene hrustančnice v Marmarskem morju:
posodobljena inventarizacija, taksonomska
vprašanja in okoljske posledice*

IHTIOLOGIJA**ITTILOGIA****ICHTHYOLOGY****Jamila RIZGALLA, Amani FITORI, Francesco
TIRALONGO & Abdalrh BEN ABDALAH**

First Records of Blennies (Suborder
Blennioidea) off the Coast of Libya 245
*Prvi zapisi o pojavljanju babic
(podred Blennioidea) ob
libijski obali*

**Farid HEMIDA, Sara LADOUL, Christian
REYNAUD & Christian CAPAPÉ**

Confirmed Occurrence of the Mediterranean
Spearfish *Tetrapturus belone* (Osteichthyes:
Istiophoridae) from the Algerian Coast
(Southwestern Mediterranean Sea) 257
*Potrjeno pojavljanje sredozemske
jadrovnice, Tetrapturus belone
(Osteichthyes: Istiophoridae) iz
alžirske obale (jugovzhodno
Sredozemsko morje)*

Lana KHREMA, Adib SAAD & Christian CAPAPÉ

First Substantiated Record of Blackfish
Centrolophus niger (Centrolophidae)
from the Syrian Coast (Eastern
Mediterranean Sea) 263
Prvi potrjen zapis črnuha, Centrolophus niger
(*Centrolophidae*), s sirske obale
(vzhodno Sredozemsko morje)

Amir IBRAHIM, Chirine HUSSEIN, Firas ALSHAWY & Alaa ALCHEIKH AHMAD

Marine Fishes (Teleostei / Osteichthyes)
of Syria (Eastern Mediterranean):
an Updated Checklist 269
Morske ribe (Teleostei / Osteichthyes)
Sirije (vzhodno Sredozemsko morje):
posodobljeni seznam vrst

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO, Enrico LA SPINA, Daniele TIBULLO & Francesco TIRALONGO

Cassiopea andromeda at the Southernmost
Tip of Italy: a Recent Arrival or an
Overlooked Resident? 287
Tujerodni klobučnjak Cassiopea andromeda
na najjužnejši konici Italije: nedavni
prišlek ali spregledan prebivalec?

Mourad CHÉRIF, Rimel BENMESSAOUD, Sihem RAFRAFI-NOUIRA, Mohamed Nouri AYADI & Christian CAPAPÉ

First Substantiated Record of the Golden-Banded
Goatfish *Upeneus moluccensis* (Osteichthyes:
Mullidae) from the Coast of Tunisia
(Central Mediterranean Sea) 295
Prvi potrjen zapis o zlatoprogem bradaču
Upeneus moluccensis (Osteichthyes: Mullidae)
z obale Tunizije (osrednje Sredozemsko morje)

Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ČETKOVIĆ, Borut MAVRIČ & Lovrenc LIPEJ

New Record of the Long-Jawed Squirrelfish,
Holocentrus adscensionis (Osbeck, 1765),
in the Adriatic Sea 303
Novi zapis o pojavljanju veveričjaka vrste
Holocentrus adscensionis (Osbeck, 1765)
v Jadranskem morju

MORSKA FAVNA**FAUNA MARINA****MARINE FAUNA****Nicola BETTOSO, Lisa FARESI, Saul CIRIACO, Marco FANTIN & Marco SEGARICH**

New Findings of the Cup-Shaped Demosponge
Calyx nicaeensis (Risso, 1826) on the Rocky
Outcrops in the Gulf of Trieste
(Northern Adriatic Sea) 311
Nove najdbe morskega keliha Calyx nicaeensis
(*Risso, 1826*) *na skalnih osamelcih v Tržaškem*
zalivu (severno Jadransko morje)

Aleksandra V. BORODINA & Yuri O. VELIAEV

Features of Lipid Accumulation in Striped Venus
Clam *Chamelea gallina* in the Sublittoral
Zone of the Crimean Coast (Black Sea) 319
Značilnosti kopičenja lipidov v navadni
venerici (Chamelea gallina) v obrežnem
pasu krimske obale (Črno morje)

Okan AKYOL & Oğuzhan TAKICAK

Recent Observations on *Monachus monachus*
(Phocidae) at Sea-Cage Fish Farms in Izmir
(Turkish Aegean Sea) 329
Nedavna opažanja primerkov Monachus monachus
(*Phocidae*) *v ribogojnicah z morskimi kletkami v*
Izmirju (turško Egejsko morje)

FLORA**FLORA****FLORA****Amelio PEZZETTA & Marco PAOLUCCI**

La Flora di Palena (Parco Nazionale della
Majella): Aggiornamento Floristico 337
Flora Palene (Nacionalni park Majella):
floristična posodobitev

DELO NAŠIH ZAVODOV IN DRUŠTEV
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ
ACTIVITIES BY OUR INSTITUTIONS AND
ASSOCIATIONS

Bojana LIPEJ

Reviving Landscapes, Connecting Species:
Lessons from the ReCo Project 383

IN MEMORIAM**Tom TURK**

V spomin Marjanu Richterju (1935–2025) 389

Kazalo k slikam na ovitku 391

Index to images on the cover 391

received: 2025-07-02

DOI 10.19233/ASHN.2025.35

RECENT OBSERVATIONS ON *MONACHUS MONACHUS* (PHOCIDAE) AT SEA-CAGE FISH FARMS IN IZMIR (TURKISH AEGEAN SEA)

Okan AKYOL

Ege University Faculty of Fisheries, 35440 Urla, Izmir, Türkiye
e-mail: okan.akyol@ege.edu.tr

Oğuzhan TAKİCAK

Ege University Urla Maritime Vocational School, 35440 Urla, Izmir, Türkiye

ABSTRACT

*On 8 February, 3 May, and 9 May 2025, three specimens of *Monachus monachus* were photographed by a staff diver at two fish farms rearing sea bass and sea bream in Gerence Bay, Izmir, in the Aegean Sea. A comprehensive set of behavioural data on Mediterranean monk seals was obtained from the diver via an interview. The findings confirm the near-permanent presence of monk seals in the study area. The paper also suggests that excessive dependence on fish farms could alter the monk seals' feeding behaviour and potentially even lead to their domestication.*

Key words: Mediterranean monk seal, feeding behaviour, interaction, Mediterranean Sea

OSSERVAZIONI RECENTI SU *MONACHUS MONACHUS* (PHOCIDAE) PRESSO IMPIANTI DI ACQUACOLTURA IN GABBIE A MARE NELL'AREA DI IZMIR (MAR EGEO TURCO)

SINTESI

*L'8 febbraio, il 3 maggio e il 9 maggio 2025, tre esemplari di *Monachus monachus* sono stati fotografati da un subacqueo del personale presso due impianti di allevamento di spigole e orate situati nella baia di Gerence, a Izmir, nel Mar Egeo. Un ampio insieme di dati comportamentali sulle foche monache del Mediterraneo è stato raccolto dal subacqueo tramite un'intervista. I risultati confermano la presenza quasi permanente di foche monache nell'area di studio. L'articolo suggerisce inoltre che una dipendenza eccessiva dagli impianti di acquacoltura potrebbe modificare il comportamento alimentare delle foche e persino portare, potenzialmente, a una loro domesticazione.*

Parole chiave: foca monaca, comportamento alimentare, interazione, Mediterraneo

INTRODUCTION

The Mediterranean monk seal (*Monachus monachus*) is regarded to be one of the rarest marine mammals in the world, with a current global population estimated at no more than 800 individuals (Dendrinis *et al.*, 2022). The species has registered a marked recovery over the past decade and is now classified as Vulnerable by the International Union for Conservation of Nature (IUCN). The largest subpopulation – estimated at 444–600 mature individuals – is currently found in the eastern Mediterranean Sea (Karamanlidis *et al.*, 2023; Karamanlidis, 2024). Other groups in the eastern Mediterranean consist of small, loosely structured aggregations, usually comprising fewer than 20 individuals. A recent study reported thirty-four sightings along the Syrian coast between 2001 and 2023 (Ibrahim *et al.*, 2024).

The Mediterranean monk seal is the sole resident pinniped species in the Mediterranean Sea (Karamanlidis, 2024). Sporadic sightings have been recorded primarily in the eastern part of the basin, including the islands of the Ionian and Aegean Seas, the mainland coast of Greece, the western and southern coasts of Türkiye, and the coasts of Cyprus (Dendrinis *et al.*, 2022). A recent study reported a total of 361 monk seal sightings in Cyprus between 2009 and 2018, with the vast majority (95%) involving juvenile and adult individuals, and only 18 sightings being of newborn pups (Nicolaou *et al.*, 2021). Additionally, Mediterranean monk seals were monitored in caves along the northern coast of Cyprus between November 2016 and May 2019, and seven individuals were confirmed: three pups and four juvenile-subadult-adult seals (Beton *et al.*, 2021).

The extirpation of the Mediterranean monk seal from the Black Sea is believed to have occurred as recently as 1997, though a small population persists in the Sea of Marmara (Dendrinis *et al.*, 2022), including two individuals recorded near Karabiga (Inanmaz *et al.*, 2014). Despite sporadic observations reported from this region, the only known active breeding populations are located in the Aegean Sea, the northeastern Mediterranean Sea, the Greek Ionian Sea, Madeira, and off the coast of Mauritania (Bundone *et al.*, 2019; Panou *et al.*, 2023).

The monk seal is strictly protected under Turkish law, European Directives, and International Conventions. In Türkiye, a national strategy for the conservation of the species was adopted in 1991, followed by the establishment of a national seal committee to coordinate conservation activities (Güçlüsoy *et al.*, 2004). In spite of these measures, the primary threats to the species in

the Mediterranean persist and can be categorised as follows: terrestrial and marine habitat loss and degradation resulting from increased human pressure, including tourism and pollution; negative interactions with fisheries and aquaculture, which occur even in countries and regions where the species is legally protected; and other unpredictable threats (Karamanlidis, 2024).

The Mediterranean monk seal spends most of its life at sea, primarily foraging for food (Dendrinis *et al.*, 2022). It is widely considered an opportunistic predator that feeds mostly on the continental shelf, with a diet dominated by fish, crustaceans, and cephalopods (Karamanlidis, 2024). Recent studies have confirmed that Mediterranean monk seals can successfully forage independently in the wild from as early as five months of age (Kıraç & Ok, 2019). Since the 1980s, the widespread expansion of mariculture in the Mediterranean has provided the species with a new and abundant food source, rivalling the availability of wild fish (Akyol & Ceyhan, 2020). This has led to monk seal attacks on sea-cages at fish farms, which have been documented. Güçlüsoy & Savaş (2003), for instance, reported 40 such attacks at 11 fish farms in the Turkish Aegean Sea between 1992 and 2000, which caused cage net damage and the escape of farmed fish. Similarly, Gerovasileiou *et al.* (2017) captured photographic evidence of seven Mediterranean monk seals at four different fish farms in the Aegean Sea.

This study provides further photographic evidence of *M. monachus* at two sea-cage fish farms in the Aegean Sea, thereby expanding our knowledge on the adapting feeding behaviour of this rare species.

MATERIAL AND METHODS

On 8 February, 3 May, and 9 May 2025, three *M. monachus* individuals (Fig. 1) were photographed by a staff diver using GoPro Hero 8 at two fish farms rearing sea bass and sea bream in Gerence Bay, İzmir (1st farm: 38.440312°N, 26.481155°E; 2nd farm: 38.449723°N, 26.417067°E, Fig. 2). The diver works at both fish farms, which are owned by the same company. The fish farms were deployed at depths of 70 m and 89 m, respectively, and are both located approximately 1 km from the mainland. They lie within the Chios and Turkish Coast IMMA (Important Marine Mammal Area) in the central eastern Aegean Sea, which encompasses an area bounded by Chios, Psarra, Çeşme, the Karaburun Peninsula, the Gulf of İzmir, and Foça, extending offshore towards the 200 m isobath (IUCN-MMPATF, 2017).




Fig. 1: Recent observations of *Monachus monachus* at sea-cage fish farms in Gerence Bay (i.e., Chios and the Turkish coast of the IMMA) include the following: (a) an adult female with a fish in its mouth, recorded on 8 February 2025 at the first fish farm; (b) an underwater view of a subadult specimen captured on 3 May 2025 at the second fish farm; and (c) an adult female observed eating fish at the sea surface on 9 May 2025 at the second fish farm (all courtesy of Davut Aydın).

Sl. 1: Nedavna opažanja vrste *Monachus monachus* v ribogojnicah v kletkah v zalivu Gerence (tj. Hios in turška obala IMMA) vključujejo naslednje: (a) odrasla samica z ribo v ustih, posneta 8. februarja 2025 v prvi ribogojnici; (b) podvodni pogled na mladostni primerek, posnet 3. maja 2025 v drugi ribogojnici; in (c) odrasla samica, opažena pri prehranjevanju z ribami na morski gladini 9. maja 2025 v drugi ribogojnici (vse z dovoljenjem Davuta Aydın).

The sex of Mediterranean monk seals was estimated based on their coloration across different life stages (Samaranch & González, 2000; Quintana MartínMontalvo & Muñoz Cañas, 2025).

RESULTS AND DISCUSSION

According to the diver we interviewed, the Mediterranean monk seals appeared occasionally during the harvest period, especially in summer. This occurred when a fish transfer bridge – made of a net and ropes

– was installed between a large cage containing reared fish (50 m in diameter) and a small, empty cage (20 m in diameter). The seals were observed entering the open-top net and stealing fish (Fig. 1a). In general, they would be seen reposing on the cage floats, exhibiting a curious but timid attitude, often observing people from a distance. Sightings increased in frequency in the spring and summer months, typically during daylight hours. The diver also reported that the seals did not display any aggressive behaviour and were never seen entering the cages themselves (D. Aydın, pers. comm.).




Fig. 2: Map showing the locations of the sea-cage fish farms (1: first fish farm, 2: second fish farm) in the Aegean Sea.

Sl. 2: Zemljevid obravnavanega območja z ribogojnicami s kletkami (1: prva ribogojnica, 2: druga ribogojnica) v Egejskem morju.

Gerovasiliou *et al.* (2017) documented an individual of *M. monachus* attacking a sea-cage fish farm in Izmir on 15 December 2016, further confirming the behaviour previously observed by Güçlüsoy & Savaş (2003). It is evident that Mediterranean monk seals are able to locate ample sustenance in the vicinity of fish farms, which act as Fish Aggregating Devices (FADs), and in the fish bridges formed during harvesting operations. This trend is problematic as it fosters negative interactions between the seals and man-made structures. Potential adverse consequences include Mediterranean monk seals abandoning open-sea foraging, increased attacks on fish farms, or harm inflicted upon the seals by unscrupulous fishers. It is therefore imperative that fishers, including

fish farm employees, understand the behaviour exhibited by Mediterranean monk seals. To support the conservation of this species, human interaction, such as feeding, should be actively discouraged. Moreover, since Mediterranean monk seals represent an important component of our natural heritage, their populations should be continuously monitored.

ACKNOWLEDGEMENTS

We would like to express our gratitude to staff diver Davut Aydın, who kindly provided the photographic material, and to two anonymous reviewers, whose valuable contributions to an earlier version of this paper are gratefully acknowledged.

NEDAVNA OPAŽANJA PRIMERKOV *MONACHUS MONACHUS* (PHOCIDAE)
V RIBOGOJNICAH Z MORSKIMI KLETKAMI V IZMIRJU
(TURŠKO EGEJSKO MORJE)

Okan AKYOL

Ege University Faculty of Fisheries, 35440 Urla, Izmir, Türkiye
e-mail: okan.akyol@ege.edu.tr

Oğuzhan TAKİCAK

Ege University Urla Maritime Vocational School, 35440 Urla, Izmir, Türkiye

POVZETEK

*Potapljač je 8. februarja, 3. maja in 9. maja 2025 fotografiral tri primerke vrste *Monachus monachus* na dveh ribogojnicah, kjer gojijo brancine in orade v zalivu Gerence v Izmirju v Egejskem morju. S pomočjo intervjujev je pridobil obsežen nabor vedenjskih podatkov o sredozemskih medvedkah. Ugotovitve potrjujejo, da je na preučevanem območju sredozemska medvedka skoraj stalno prisotna. Poleg tega prispevek tudi nakazuje, da bi lahko pretirana odvisnost od ribogojnic spremenila prehranjevalne navade sredozemskih medvedk in morda celo privedla do njihove udomačitve.*

Ključne besede: sredozemska medvedka, prehranjevalno vedenje, interakcije, Sredozemsko morje

REFERENCES

- Akyol, O. & T. Ceyhan (2020):** Wild fish communities associated with sea-cage fish farms in the Aegean Sea. In: Çoban, D., M.D. Demircan, & D.D. Tosun (Eds.): Marine Aquaculture in Turkey: Advancements and Management. Turkish Marine Research Foundation (TUDAV) Publication No: 59, İstanbul. pp. 401-415.
- Beton, D., A.C. Broderick, B.J. Godley, E. Kolaç, M. Ok, & R.T.E. Snape (2021):** New monitoring confirms regular breeding of the Mediterranean monk seal in Northern Cyprus. *Oryx*, 55(4), 522-525.
- Bundone, L., A. Panou, & E. Molinaroli (2019):** On sightings of (vagrant?) monk seals, *Monachus monachus*, in the Mediterranean Basin and their importance for the conservation of the species. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 29(4), 554-563.
- Dendrinis, P., S. Adamantopoulou, E. Tounta, K. Koemtzopoulos, & A.A. Karamanlidis (2022):** The Mediterranean Monk Seal (*Monachus monachus*) in the Aegean Sea. "The Living Myth of the Sirens". In: Anagnostou, C.L., A.G. Kostianoy, I.D. Mariolakis, P. Panayotidis, M. Soilemezidou, & G. Tsaltas (Eds.): The Aegean Sea Environment. The Handbook of Environmental Chemistry, Vol. 129. Springer, Cham. pp. 211-233.
- Gerovasileiou, V., E.H.Kh. Akel, O. Akyol, G. Alongi, F. Azevedo, N. Babali, et al., (2017):** New Mediterranean Biodiversity Records (July 2017). *Med. Mar. Sci.*, 18(2), 355-384.
- Güçlüsoy, H. & Y. Savaş (2003):** Interaction between monk seal, *Monachus monachus* (Hermann, 1779) and marine fish farms in the Turkish Aegean and management of the problem. *Aquac. Res.*, 34, 777-783.
- Güçlüsoy, H., C.O. Kırac, N.O. Veryeri & Y. Savaş (2004):** Status of the Mediterranean monk seal, *Monachus monachus* (Hermann, 1779) in the coastal waters of Turkey. *Ege J. Fish. & Aquat. Sci.*, 2, 201-210.
- Ibrahim, A., C. Hussein, F. Alshawy, & A.A. Ahmad (2024):** Distribution and habitats of the Mediterranean Monk Seal (*Monachus monachus*): In the Syrian Coast (Eastern Mediterranean). *Species*, 25, e2s1624.
- Inanmaz, Ö.E., Ö. Degirmenci, & A.C. Gücü (2014):** A new sighting of the Mediterranean monk seal, *Monachus monachus* (Hermann, 1779), in the Marmara Sea (Turkey). *Zool. Middle East*, 60(3), 278-280.
- IUCN-MMPATF (2017):** Chios and Turkish Coast IMMA. Full Accounts of Mediterranean IMMA Factsheet. IUCN Joint SSC/WCPA Marine Mammal Protected Areas Task Force, 2017. Available at <https://www.marinemammalhabitat.org/portfolioitem/chios-and-turkish-coast/>.
- Karamanlidis, A.A., P. Dendrinis, P. Fernandez de Larrinoa, C.O. Kırac, H. Nicolaou & R. Pires (2023):** *Monachus monachus*. The IUCN Red List of Threatened Species 2023: e.T13653A238637039. <https://dx.doi.org/10.2305/IUCN.UK.2023-1.RLTS.T13653A238637039.en>. Accessed on 25 Aug. 2025.
- Karamanlidis, A.A. (2024):** Current status, biology, threats and conservation priorities of the Vulnerable Mediterranean monk seal. *Endanger. Species Res.*, 53, 341-361.
- Kırac, C.O. & M. Ok. (2019):** Diet of a Mediterranean monk seal *Monachus monachus* in a transitional post-weaning phase and its implications for the conservation of the species. *Endanger. Species Res.*, 39, 315-320.
- Nicolaou, H., P. Dendrinis, M. Marcou, S. Michaelides, & A.A. Karamanlidis (2021):** Re-establishment of the Mediterranean monk seal *Monachus monachus* in Cyprus: priorities for conservation. *Oryx*, 55(4), 526-528.
- Panou, A., M. Giannoulaki, D. Varda, L. Lazaj, G. Pojana, & L. Bundone (2023):** Towards a strategy for the recovering of the Mediterranean monk seal in the Adriatic-Ionian Basin. *Front. Mar. Sci.*, 10, 1034124.
- Quintana MartínMontalvo, B. & M. Muñoz Cañas (2025):** Mediterranean monk seal. A comprehensive set of monitoring and research techniques for the study and conservation of *Monachus monachus* in the Mediterranean Sea (1st ed.). Gland, Switzerland: IUCN., pp.1-77.
- Samaranch, R. & L.M. González (2000):** Changes in morphology with age in Mediterranean monk seals (*Monachus monachus*). *Marine Mammal Science*, 16(1), 141-157.