

# ANNALES

*Anali za istrske in mediteranske študije*  
*Annali di Studi istriani e mediterranee*  
*Annals for Istrian and Mediterranean Studies*  
*Series Historia Naturalis, 35, 2025, 2*





# ANNALES

**Anali za istrske in mediteranske študije**  
**Annali di Studi istriani e mediterraneei**  
**Annals for Istrian and Mediterranean Studies**

**Series Historia Naturalis, 35, 2025, 2**

ISSN 1408-533X  
e-ISSN 2591-1783

UDK 5

Letnik 35, leto 2025 številka 2

**UREDNIŠKI ODBOR/  
COMITATO DI REDAZIONE/  
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Serena Fonda Umani (IT), Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/  
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/  
Redattore responsabile per le scienze  
naturali/Natural Science Editor:**

Lovrenc Lipej

**Urednica/Redattrice/Editor:**

Martina Orlando-Bonaca

**Prevajalci/Traduttori/Translators:**

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/  
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

**Tisk/Stampa/Print:**

Založništvo PADRE d.o.o.

**Založnika/Editori/Published by:**Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®  
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/  
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran  
SI-6330 Piran /Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;  
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 6. 12. 2025.

**Sofinancirajo/Supporto finanziario/  
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

*Annales - Series Historia Naturalis* izhaja dvakrat letno.**Naklada/Tiratura/Circulation:**

300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>  
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>  
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / *CONTENTS*SREDOZEMSKÉ HRUSTANČNICE  
*SQUALI E RAZZE MEDITERRANEE*  
*MEDITERRANEAN SHARKS AND RAYS***Hakan KABASAKAL**

Analysis of Confirmed Shark Attacks  
in the Eastern Mediterranean Sea and  
the Sea of Marmara (1827–2025) ..... 169  
*Analiza potrjenih napadov morskih psov*  
*v vzhodnem Sredozemskem morju*  
*in Marmarskem morju (1827–2025)*

**Alen SOLDÓ**

Observations of a Juvenile Basking Shark  
*Cetorhinus maximus* in the Adriatic Sea  
Support the Hypothesis of a Distinct  
Mediterranean Population ..... 187  
*Opazovanja mladega morskega psa*  
*orjaka (Cetorhinus maximus)*  
*v Jadranskem morju podpirajo*  
*hipotezo o ločeni sredozemski*  
*populaciji*

**Jacopo BERNARDI, Pero UGARKOVIĆ &  
Ilija ČETKOVIĆ**

An Unusual Encounter with a Juvenile Kitefin  
Shark, *Dalatias licha*, in Shallow Coastal  
Waters of Montenegro (Adriatic Sea) ..... 197  
*Nenavadno srečanje z mladim primerkom*  
*klinoplavutega morskega psa, Dalatias licha,*  
*v plitvih obalnih vodah Črne gore*  
*(Jadransko morje)*

**Cemal TURAN, Alen SOLDÓ, Servet A. DOĞDU,  
Funda TURAN, Ayşegül ERGENLER & Ali UYAN**

Identification of a Potential Nursery Ground  
of the Spiny Butterfly Ray, *Gymnura altavela*,  
in the Northeastern Mediterranean Sea, Türkiye ..... 205  
*Identifikacija potencialnih jaslčic za*  
*metuljastega skata, Gymnura*  
*altavela, v severovzhodnem*  
*Sredozemskem morju,*  
*Turčija*

**Alen SOLDÓ, Eleonora DE SABATA &  
Simona CLO**

Recent Records of Critically Endangered  
Common Guitarfish *Rhinobatos rhinobatos*  
(Linnaeus, 1758) in the Northern  
Mediterranean ..... 213  
*Nedavni zapiski o pojavljanju kritično*  
*ogroženega navadnega goslaša*  
*Rhinobatos rhinobatos*  
*(Linnaeus, 1758) v*  
*severnem Sredozemlju*

**Hakan KABASAKAL & F. Saadet KARAKULAK**

Demersal Elasmobranchs of the Sea of Marmara:  
Updated Inventory, Taxonomic Issues and  
Environmental Implications ..... 221  
*Pridnene hrustančnice v Marmarskem morju:*  
*posodobljena inventarizacija, taksonomska*  
*vprišanja in okoljske posledice*

**IHTIOLOGIJA****ITTILOGIA****ICHTHYOLOGY****Jamila RIZGALLA, Amani FITORI, Francesco  
TIRALONGO & Abdalrh BEN ABDALAH**

First Records of Blennies (Suborder  
Blennioidea) off the Coast of Libya ..... 245  
*Prvi zapisi o pojavljanju babic*  
*(podred Blennioidea) ob*  
*libijski obali*

**Farid HEMIDA, Sara LADOUL, Christian  
REYNAUD & Christian CAPAPÉ**

Confirmed Occurrence of the Mediterranean  
Spearfish *Tetrapturus belone* (Osteichthyes:  
Istiophoridae) from the Algerian Coast  
(Southwestern Mediterranean Sea) ..... 257  
*Potrjeno pojavljanje sredozemske*  
*jadrovnice, Tetrapturus belone*  
*(Osteichthyes: Istiophoridae) iz*  
*alžirske obale (jugovzhodno*  
*Sredozemsko morje)*

**Lana KHREMA, Adib SAAD & Christian CAPAPÉ**

First Substantiated Record of Blackfish  
*Centrolophus niger* (Centrolophidae)  
from the Syrian Coast (Eastern  
Mediterranean Sea) ..... 263  
*Prvi potrjen zapis črnuha, Centrolophus niger*  
(*Centrolophidae*), s sirske obale  
(vzhodno Sredozemsko morje)

**Amir IBRAHIM, Chirine HUSSEIN, Firas ALSHAWY & Alaa ALCHEIKH AHMAD**

Marine Fishes (Teleostei / Osteichthyes)  
of Syria (Eastern Mediterranean):  
an Updated Checklist ..... 269  
*Morske ribe (Teleostei / Osteichthyes)*  
*Sirije (vzhodno Sredozemsko morje):*  
*posodobljeni seznam vrst*

BIOTSKA GLOBALIZACIJA  
GLOBALIZZAZIONE BIOTICA  
BIOTIC GLOBALIZATION

**Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO, Enrico LA SPINA, Daniele TIBULLO & Francesco TIRALONGO**

*Cassiopea andromeda* at the Southernmost  
Tip of Italy: a Recent Arrival or an  
Overlooked Resident? ..... 287  
*Tujerodni klobučnjak Cassiopea andromeda*  
*na najjužnejši konici Italije: nedavni*  
*prišlek ali spregledan prebivalec?*

**Mourad CHÉRIF, Rimel BENMESSAOUD, Sihem RAFRAFI-NOUIRA, Mohamed Nouri AYADI & Christian CAPAPÉ**

First Substantiated Record of the Golden-Banded  
Goatfish *Upeneus moluccensis* (Osteichthyes:  
Mullidae) from the Coast of Tunisia  
(Central Mediterranean Sea) ..... 295  
*Prvi potrjen zapis o zlatoprogem bradaču*  
*Upeneus moluccensis (Osteichthyes: Mullidae)*  
*z obale Tunizije (osrednje Sredozemsko morje)*

**Nikola DJORDJEVIĆ, Slavica PETOVIĆ, Ilija ČETKOVIĆ, Borut MAVRIČ & Lovrenc LIPEJ**

New Record of the Long-Jawed Squirrelfish,  
*Holocentrus adscensionis* (Osbeck, 1765),  
in the Adriatic Sea ..... 303  
*Novi zapis o pojavljanju veveričjaka vrste*  
*Holocentrus adscensionis (Osbeck, 1765)*  
*v Jadranskem morju*

**MORSKA FAVNA****FAUNA MARINA****MARINE FAUNA****Nicola BETTOSO, Lisa FARESI, Saul CIRIACO, Marco FANTIN & Marco SEGARICH**

New Findings of the Cup-Shaped Demosponge  
*Calyx nicaeensis* (Risso, 1826) on the Rocky  
Outcrops in the Gulf of Trieste  
(Northern Adriatic Sea) ..... 311  
*Nove najdbe morskega keliha Calyx nicaeensis*  
(*Risso, 1826*) *na skalnih osamelcih v Tržaškem*  
*zalivu (severno Jadransko morje)*

**Aleksandra V. BORODINA & Yuri O. VELIAEV**

Features of Lipid Accumulation in Striped Venus  
Clam *Chamelea gallina* in the Sublittoral  
Zone of the Crimean Coast (Black Sea) ..... 319  
*Značilnosti kopičenja lipidov v navadni*  
*venerici (Chamelea gallina) v obrežnem*  
*pasu krimske obale (Črno morje)*

**Okan AKYOL & Oğuzhan TAKICAK**

Recent Observations on *Monachus monachus*  
(Phocidae) at Sea-Cage Fish Farms in Izmir  
(Turkish Aegean Sea) ..... 329  
*Nedavna opažanja primerkov Monachus monachus*  
(*Phocidae*) *v ribogojnicah z morskimi kletkami v*  
*Izmirju (turško Egejsko morje)*

**FLORA****FLORA****FLORA****Amelio PEZZETTA & Marco PAOLUCCI**

La Flora di Palena (Parco Nazionale della  
Majella): Aggiornamento Floristico ..... 337  
*Flora Palene (Nacionalni park Majella):*  
*floristična posodobitev*

DELO NAŠIH ZAVODOV IN DRUŠTEV  
ATTIVITÀ DEI NOSTRI ISTITUTI E SOCIETÀ  
ACTIVITIES BY OUR INSTITUTIONS AND  
ASSOCIATIONS

**Bojana LIPEJ**

Reviving Landscapes, Connecting Species:  
Lessons from the ReCo Project ..... 383

**IN MEMORIAM****Tom TURK**

*V spomin Marjanu Richterju (1935–2025)* ..... 389

Kazalo k slikam na ovitku ..... 391

**Index to images on the cover** ..... 391

received: 2025-10-06

DOI 10.19233/ASHN.2025.30

## CASSIOPEA ANDROMEDA AT THE SOUTHERNMOST TIP OF ITALY: A RECENT ARRIVAL OR AN OVERLOOKED RESIDENT?

*Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO*

Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
e-mail: pleotta1990@gmail.com; rocco.cicciarella@hotmail.it; ruben.garrano@gmail.com

*Enrico LA SPINA*

Department of Biomedical and Biotechnological Sciences, University of Catania, 95123 Catania, Italy  
e-mail: enricolaspina@outlook.it

*Daniele TIBULLO*

Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
Department of Biomedical and Biotechnological Sciences, University of Catania, 95123 Catania, Italy  
e-mail: daniele.tibullo@unict.it

*Francesco TIRALONGO*

Department of Biological, Geological and Environmental Sciences, University of Catania, 95124 Catania, Italy  
Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
e-mail: francesco.tiralongo@unict.it

### ABSTRACT

*The Lessepsian jellyfish Cassiopea andromeda is an invasive scyphozoan originating from the Red Sea and Indo-Pacific region. It was first recorded in the Mediterranean at the beginning of the 20th century and has since expanded its distribution across the basin. The main goal of this contribution is to document the first and well-documented record of this jellyfish from the Portopalo di Capo Passero area, which represents the southernmost tip of Italy (Sicily, Ionian Sea). The observation, made in October 2025, provides new evidence of the species' expansion along the Sicilian Ionian coast. This finding raises questions about whether the species is a recent colonizer or an overlooked resident and highlights the biogeographical and ecological importance of this area as a potential hotspot for the monitoring of Lessepsian and thermophilic species.*

**Key words:** jellyfish, alien species, Ionian Sea, Mediterranean, Lessepsian migrant

## CASSIOPEA ANDROMEDA ALL'ESTREMO SUD D'ITALIA: UNA PRESENZA RECENTE O UNA SPECIE PASSATA INOSSERVATA?

### SINTESI

*La medusa lessepsiana Cassiopea andromeda è uno scifozoo invasivo originario del Mar Rosso e della regione indo-pacifica. È stata segnalata per la prima volta nel Mar Mediterraneo all'inizio del XX secolo e da allora ha progressivamente ampliato la propria distribuzione in tutto il bacino. Il principale obiettivo di questo studio è quello di documentare la prima segnalazione di questa specie nell'area di Portopalo di Capo Passero, estrema punta meridionale d'Italia (Sicilia, Mar Ionio). L'osservazione, effettuata nell'ottobre 2025, fornisce nuove evidenze dell'espansione della specie lungo la costa ionica siciliana. Questo ritrovamento solleva interrogativi sul fatto che si tratti di un colonizzatore recente o di una specie precedentemente passata inosservata, e sottolinea l'importanza biogeografica ed ecologica di quest'area come potenziale hotspot per specie lessepsiane e termofile.*

**Parole chiave:** medusa, specie aliena, Mar Ionio, Mediterraneo, migrante lessepsiano


## INTRODUCTION

The Mediterranean Sea, recognized as a global biodiversity hotspot, hosts an exceptionally rich marine fauna with over 17,000 recorded species, representing about 7% of the world's marine animals (Coll *et al.*, 2010). Nowadays, this biodiversity is undergoing profound transformations driven by human activities, including both direct and indirect species introductions (Stock *et al.*, 2018). The spread of non-indigenous species (NIS), also referred to as alien, exotic, or non-native species, has become a key indicator of ecological imbalance and biodiversity decline (Katsanevakis *et al.*, 2014; Tiralongo *et al.*, 2020). Invasive alien species are now regarded as one of the primary threats to the integrity and functioning of Mediterranean marine ecosystems (Pyšek *et al.*, 2020).

In this context, the “upside-down jellyfish” *Cassiopea andromeda* (Forskål, 1775) is a Lessepsian immigrant native to the Red Sea and is widely distributed throughout the Indo-Pacific region. It entered the Mediterranean Sea through the Suez Canal and was first recorded off Cyprus in 1903 (Maas,

1903). Since then, it has progressively expanded westwards, with confirmed records of the species in the Levantine, Aegean and central Mediterranean sub-basins (Galil *et al.*, 1990; Çevik *et al.*, 2006; Schembri *et al.*, 2010; Ramos-Pérez *et al.*, 2025). More recently, dense aggregations of the jellyfish have been reported in semi-enclosed eutrophic environments such as harbours and salt pans (Cillari *et al.*, 2018; Deidun *et al.*, 2018; Kleitou *et al.*, 2025), while the westernmost Mediterranean record of the species was documented in Spain (Marambio *et al.*, 2025).

This species is highly tolerant to environmental fluctuations and is capable of asexual reproduction, which facilitates its rapid establishment in shallow coastal habitats (Thé *et al.*, 2021). Species of the genus *Cassiopea* exhibit a mixotrophic lifestyle sustained by an intracellular symbiosis with dinoflagellates (Symbiodiniaceae), which are acquired from the environment during the polyp stage (Djeghri *et al.*, 2019). This mutualistic association significantly shapes the holobiont's biochemical composition, nutritional strategy and overall metabolism (De Domenico *et al.*, 2025). Moreover, given the high



**Fig. 1: Specimens of *C. andromeda* observed inside the harbour of Portopalo di Capo Passero (Sicily, Ionian Sea) on 19 October 2025: (A) during measurements; (B) sampled specimens; (C) in situ individuals.**

**Sl. 1: Primerki vrste *C. andromeda*, opaženi v pristanišču Portopalo di Capo Passero (Sicilija, Jonsko morje) 19. oktobra 2025: (A) med meritvami; (B) vzorčeni osebki; (C) osebki in situ.**

interspecific morphological similarity within the genus, accurate species identification of *Cassiopea* often requires an integrative taxonomic approach, combining both morphological and molecular data (Rowe et al., 2025). From a morphological perspective, diagnostic characters such as the number and distribution of large appendages as well as the oral arm branching pattern have proven useful for species discrimination within the genus (Rowe et al., 2025).

Following its introduction from the Red Sea into the Mediterranean through the Suez Canal, *Cassiopea andromeda* has expanded its range under the influence of climate change and consequently to increasing seawater temperatures, now occurring more widely than previously recorded (Holland et al., 2004; Fumarola et al., 2025).

Here, we report the first well-documented occurrence of *C. andromeda* from the Portopalo di Capo Passero area (southeastern Sicily, Ionian Sea), providing new evidence of its range expansion within southern Italian waters. Preliminary data on individual size composition and on local abundance were also collected.

## MATERIAL AND METHODS


On 19 October 2025, several individuals of *Cassiopea andromeda* were observed and photographed by the authors in a sheltered coastal area inside the fishery harbour of Portopalo di Capo Passero (36.67095N, 15.12699E), at a depth of about 0.7 m (Fig. 1). The site is characterized by sandy and muddy substrates with reduced water circulation and elevated summer temperatures. Moreover, during the survey, two alien invasive crab species, namely *Callinectes sapidus* and *Portunus segnis*, were detected. A total of 42 specimens of *C. andromeda* were measured, recording the diameter of the exumbrella using a flexible measuring tape with a 0.1 cm precision. A total of 10 specimens were sampled in order to verify the identity of the species, whose general morphology clearly matched that of *C. andromeda* (Ames et al., 2020; Karunaratne et al., 2020). After the analyses, the specimens were preserved in alcohol for potential future laboratory examinations (e.g., molecular analysis) and deposited in the zoological collection of the Ente Fauna Marina Mediterranea in Avola (Sicily) under the code #EFMM191025. A species' density (individuals/m<sup>2</sup>) estimation was carried out. Finally, an updated map (Fig. 2) based on our new record and on the data provided by Katsanevakis et al. (2020) was produced, using the recent model proposed for the Mediterranean by Ramos-Pérez et al. (2025).

## RESULTS AND DISCUSSION

Specimens displayed the typical morphology and general colour pattern of *Cassiopea andromeda*: a flattened umbrella, whitish to brownish coloration, conspicuous white blotches on the exumbrella, and eight branched oral arms. However, although the general morphology and colour pattern clearly point to the genus *Cassiopea*, the absence of molecular analyses prevents a definitive species-level identification (Gamero-Mora et al., 2022; Rowe et al., 2025). The bell diameters of specimens measured from 5 to 13 cm, with a mean value of  $8.6 \pm 2.4$  cm. This size variability reflects a population composed of both juvenile and adult individuals, suggesting that the species is able to complete different stages of its life cycle within the studied area. Individuals rested inverted on the seabed, with the oral arms oriented upward to maximize exposure of the zooxanthellae to light, an adaptive strategy associated with their mixotrophic feeding (Symbiodiniaceae symbiosis) (Arossa et al., 2021). The estimated mean jellyfish density in the area, where the species was unevenly distributed, was that of approximately 8 jellyfish individuals/m<sup>2</sup>.

This record represents the southernmost occurrence of *C. andromeda* in Italian waters, about 120 km south of the population recently documented in the Augusta salt pans (Kleitou et al., 2025). The environmental conditions of Portopalo's harbour, with warm, eutrophic, and semi-enclosed calm waters, closely resemble habitats where the species has established stable aggregations elsewhere in the Mediterranean (Schembri et al., 2010; Cillari et al., 2018; Deidun et al., 2018; Kleitou et al., 2025).

Given its broad thermal (6–39 °C, with an optimum at 35.7 °C; Fumarola et al., 2025) and moderate salinity tolerance (30–50, Aljbour & Agustí, 2025), *Cassiopea andromeda* is likely to be more widespread in the Mediterranean Sea than previously assumed. This suggests that transitional environments such as saltpans, coastal lagoons, and harbours are particularly suitable for the presence of the species and should therefore be carefully inspected and monitored. The finding also raises the possibility that the species had previously remained undetected due to a limited monitoring in shallow coastal lagoons and harbour areas. The location of Portopalo, near the confluence of Ionian and Strait of Sicily currents, could act as an ecological “gateway” for a further dispersal of the species towards western and northern Mediterranean waters. Several hypotheses can be considered regarding the putative introduction pathway responsible for bringing the species to Portopalo. The proximity of Portopalo to main



**Fig. 2:** Distribution map of *Cassiopea andromeda* records in the Mediterranean Sea. In red circles historical records, in light blue historical records from the Suez Canal (Red Sea), and the yellow one represents the new record from Portopalo di Capo Passero (modified from Ramos-Pérez et al., 2025 and updated with records in Katsanevakis et al., 2020).

**Sl. 2:** Zemljevid razširjenosti vrste *Cassiopea andromeda* na podlagi zapisov o pojavljanju v Sredozemskem morju. V rdečih krogih so zgodovinski zapisi o pojavljanju, v svetlo modrih krogih zgodovinski zapisi o pojavljanju iz Sueškega prekopa (Rdeče morje), rumeni pa predstavlja nov zapis o pojavljanju iz Portopalo di Capo Passero (spremenjen po Ramos-Pérez in sod., 2025 in posodobljen z zapisi v Katsanevakis in sod., 2020).

commercial shipping routes and its intense fishing and boating activities suggest that *C. andromeda* may have been introduced through shipping-related vectors, either via ballast water or as fouling - polyps attached to ship hulls. Alternatively, the species could have reached the area by natural dispersal or through recreational vessel traffic from already-established populations in Malta or along the eastern Sicilian coast, supported by favourable currents and rising sea temperatures that enhance larval survival and transport. The latter pathway is especially plausible given the high volumes of recreational vessel traffic between Malta and Sicily during the summer season. The ability to exploit multiple energy sources, namely photosynthates provided by symbiotic dinoflagellates and heterotrophic feeding, enables *C. andromeda* to thrive under a wide range of environmental conditions (Thé et al., 2023). *Cassiopea* genus can form

high-density blooms when environmental conditions are favourable and anthropogenic pressures intensive, particularly with rising sea temperatures linked to climate change and coastal development. Such blooms may exert competitive pressure on seagrass and other vegetal communities, primarily by limiting the availability of space, light, and food (Rowe et al., 2025). Moreover, the trophic flexibility of *Cassiopea* genus, linked to both heterotrophic feeding and photosynthetic symbionts, can influence primary productivity, nutrient cycling, and local food web dynamics, potentially generating cascading ecological effects, as observed in other symbiotic jellyfish species (Djehri et al., 2021).

Continuous monitoring with the possibility to involve citizen science networks, will be essential to evaluate whether this population is ephemeral or self-sustaining, as well as to assess its ecological effects on local benthic communities and

on dissolved oxygen dynamics. Considering the species' long history of westward expansion in the Mediterranean (Fumarola *et al.*, 2025), it remains uncertain whether its presence in Portopalo represents a recent colonization event favoured by climate warming or whether the species has remained undetected for years due to its occurrence in localised habitats and due to the lack of specific expert monitoring programs.

In the context of this study, the harbour of Portopalo di Capo Passero can be regarded as a hotspot of alien biodiversity, as evidenced by the

occurrence of several alien species, including the invasive portunid crabs *Callinectes sapidus* and *Portunus segnis*. The coexistence of these taxa highlights the ecological susceptibility of this semi-enclosed coastal system to biological invasions and its potential role as an entry point and establishment site for thermophilic and Lessepsian species in the central Mediterranean. Indeed, such environments, often characterized by anthropogenic disturbance and intense maritime traffic, may act as stepping stones for the secondary dispersal of Lessepsian and thermophilic taxa along the Sicilian coast.

## TUJERODNI KLOBUČNJAK CASSIOPEA ANDROMEDA NA NAJJUŽNEJŠI KONICI ITALIJE: NEDAVNI PRIŠLEK ALI SPREGLEDAN PREBIVALEC?

*Paola LEOTTA, Rocco CICCARELLA, Ruben GARRANO*

Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
e-mail: pleotta1990@gmail.com; rocco.cicciarella@hotmail.it; ruben.garrano@gmail.com

*Enrico LA SPINA*

Department of Biomedical and Biotechnological Sciences, University of Catania, 95123 Catania, Italy  
e-mail: enricolaspina@outlook.it

*Daniele TIBULLO*

Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
Department of Biomedical and Biotechnological Sciences, University of Catania, 95123 Catania, Italy  
e-mail: daniele.tibullo@unict.it

*Francesco TIRALONGO*

Department of Biological, Geological and Environmental Sciences, University of Catania, 95124 Catania, Italy  
Ente Fauna Marina Mediterranean, Scientific Organization for Research and Conservation of Marine Biodiversity, 96012 Avola, Italy  
e-mail: francesco.tiralongo@unict.it

### POVZETEK

*Lesepski klobučnjak Cassiopea andromeda je invazivna vrsta klobučnjakov, ki izvira iz Rdečega morja in indo-pacifiške regije. V Sredozemskem morju so jo prvič zabeležili na začetku 20. stoletja in od takrat se je razširila po celotnem bazenu. Glavni cilj tega prispevka je bil dokumentirati prvi in dobro dokumentiran zapis te klobučnjaške meduze z območja Portopalo di Capo Passero, ki predstavlja najjužnejšo konico Italije (Sicilija, Jonsko morje). Opazovanje primerkov te vrste v oktobru 2025 prinaša nove dokaze o širjenju vrste vzdolž sicilijansko-jonske obale. Ta ugotovitev sproža vprašanja o tem, ali je vrsta nedavni kolonizator ali spregledan prebivalec, in poudarja biogeografski in ekološki pomen tega območja kot potencialne žariščne točke za spremljanje lesepskih in toploljubnih vrst.*

**Ključne besede:** klobučnjaki, tujerodne vrste, Jonsko morje, Sredozemsko morje, lesepska selivka

## REFERENCES

- Aljbour, S.M. & S. Agustí (2025):** Salinity over source: Osmotic stress and metabolic plasticity in *Cassiopea andromeda* under desalination and runoff exposure. *Regional Studies in Marine Science*, 90, 104464.
- Ames, C.L., A.M.L. Klompen, K. Badhiwala, K. Muffett, A.J. Reft, M. Kumar, J.D. Janssen, J.N. Schultzhau, L.D. Field, M.E. Muroski, N. Bezio, J.T. Robinson, D.H. Leary, P. Cartwright, A.G. Collins & G.J. Vora (2020):** Cassiosomes are stinging-cell structures in the mucus of the upside-down jellyfish *Cassiopea xamachana*. *Communications Biology*, 3, 67.
- Arossa, S., A. Barozzi, M. Callegari, S.G. Klein, A.J. Parry, S.-H. Hung, A. Steckbauer, M. Aranda, D. Daffonchio & C.M. Duarte (2021):** The internal microenvironment of the symbiotic jellyfish *Cassiopea* sp. from the Red Sea. *Frontiers in Marine Science*, 8, 705915.
- Çevik, C., İ.L. Erkol & B. Toklu (2006):** A new record of an alien jellyfish from the Levantine coast of Turkey - *Cassiopea andromeda* (Forskål, 1775) [Cnidaria: Scyphozoa: Rhizostomea]. *Aquatic Invasions*, 1(3), 196-197.
- Cillari, T., F. Andaloro & L. Castriota (2018):** First documented record of *Cassiopea cf. andromeda* (Cnidaria: Scyphozoa) in Italian waters. *Cahiers de Biologie Marine*, 59(2), 193-195.
- Coll, M., C. Piroddi, J. Steenbeek, K. Kaschner, F. Ben Rais Lasram, J. Aguzzi, E. Ballesteros, C.N. Bianchi, J. Corbera, T. Dailianis, R. Danovaro, M. Estrada, C. Frogliia, B.S. Galil, J.M. Gasol, R. Gertwagen, J. Gil, F. Guilhaumon, K. Kesner-Reyes, M.-S. Kitsos, A. Koukouras, N. Lampadariou, E. Laxamana, C.M. López-Fé de la Cuadra, H.K. Lotze, D. Martin, D. Mouillot, D. Oro, S. Raicevich, J. Rius-Barile, J.I. Saiz-Salinas, C. San Vicente, S. Somot, J. Templado, X. Turon, D. Vafidis, R. Villanueva & E. Voultsiadou (2010):** The biodiversity of the Mediterranean Sea: estimates, patterns, and threats. *PLoS ONE*, 5(8), e11842.
- De Domenico, S., A. Toso, G. De Rinaldis, G. Tedesco, S. Mammone, A. Allegra, A. Spanò & F. Cammarata (2025):** Wild or reared? *Cassiopea andromeda* jellyfish as a potential biofactory. *Marine Drugs*, 23, 19-23.
- Deidun, A., A. Gauci, A. Sciberras & S. Piraino (2018):** Back with a bang – an unexpected massive bloom of *Cassiopea andromeda* (Forskål, 1775) in the Maltese Islands, nine years after its first appearance. *BiolInvasions Records*, 7(4), 399-404.
- Djehgri, N., P. Pondaven, H. Stibor & M.N. Dawson (2019):** Review of the diversity, traits, and ecology of zooxanthellate jellyfishes. *Marine Biology*, 166(11), 147.
- Djehgri, N., P. Pondaven, F. Le Grand, A. Bideau, N. Duquesne, M. Stockenreiter, S. Behl, J.Y.-T. Huang, T. Hansen, S. Patris, G. Ucharm & H. Stibor (2021):** High trophic plasticity in the mixotrophic *Mastigias papua*-*Symbiodiniaceae* holobiont. *Marine Ecology Progress Series*, 666, 73-88.
- Fumarola, L.M., V. Leoni, G. Marchessaux, G. Sarà, S. Piraino & M. Bosch-Belmar (2025):** Global warming and the spread of the introduced jellyfish *Cassiopea andromeda*: thermal niche and habitat suitability in the Mediterranean Sea. *Global Change Biology*, 31(10), e70548.
- Galil, B.S., E. Spanier & W.W. Ferguson (1990):** The Scyphomedusae of the Mediterranean coast of Israel, including two Lessepsian migrants new to the Mediterranean. *Zoologische Mededelingen*, 64, 95-105.
- Gamero-Mora, E., A.G. Collins, S.R. Boco, S.M. Geson III & A.C. Morandini (2022):** Revealing hidden diversity among upside-down jellyfishes (Cnidaria: Scyphozoa: Rhizostomeae: *Cassiopea*): distinct evidence allows the change of status of a neglected variety and the description of a new species. *Invertebrate Systematics*, 36, 63-89.
- Holland, B.S., M.N. Dawson, G.L. Crow & D.K. Hofmann (2004):** Global phylogeography of *Cassiopea* (Scyphozoa: Rhizostomeae): molecular evidence for cryptic species and multiple invasions of the Hawaiian Islands. *Marine Biology*, 145, 1119-1128.
- Karunaratne, K.D., S.M. Liyanarachchi & M.D.S.T. de Croos (2020):** First record of upside-down jellyfish *Cassiopea andromeda* (Forskål, 1775) (Cnidaria: Scyphozoa: Rhizostomeae: Cassiopeidae) from Sri Lanka. *Sri Lanka Journal of Aquatic Sciences*, 25(2), 57-65.
- Kleitou, P., D. Agius, S. Akalin, M. Albano, I.A. Ammar, C.M. Aydin, E. Azzurro, L. Bellomo, A. Bonifazi, G. Capillo, F. Crocetta, I. Cvitković, A. Fortič, S. Giakoumi, D. Grech, H.B. Hasan, F. Laspina, M. Lezzi, P. Lucic, E. Mancini, W.M. Mayya, M.T. Olgüner, P. Ovalis, L. Pisani, V. Pitacco, A. Salman, L. Sandri, P. Slavinec, F. Tiralongo, F. Turano, A. Yildiz, A. Zenetos, A.H. Zieni & A. Žuljević (2025):** New records of introduced species in the Mediterranean Sea (February 2025). *Mediterranean Marine Science*, 26(1), 175-198.
- Katsanevakis, S., M. Coll, C. Piroddi, J. Steenbeek, F. Ben Rais Lasram, A. Zenetos & A.C. Cardoso (2014):** Invading the Mediterranean Sea: biodiversity patterns shaped by human activities. *Frontiers in Marine Science*, 1, 32.

- Katsanevakis, S., D. Poursanidis, R. Hoffman, J. Rizzagalla, S.B.-S. Rothman, Y. Levitt-Barmats, L. Hadjioannou, D. Trkov, J.M. Garmendia, M. Rizzo, A.G. Bartolo, M. Bariche, F. Tomas, P. Kleitou, P.J. Schembri, D. Kletou, F. Tiralongo, C. Pergent, G. Pergent, E. Azzurro, M. Bilecenoglu, A. Lodola, E. Ballesteros, V. Gerovasileiou, M. Verlaque, A. Occhipinti-Ambrogi, E. Kytinou, T. Dailianis, J. Ferrario, F. Crocetta, C. Jimenez, J. Evans, M. Ragkousis, L. Lipej, J.A. Borg, C. Dimitriadis, G. Chatzigeorgiou, P.G. Albano, S. Kalogirou, H. Bazairi, F. Espinosa, J. Ben Souissi, K. Tsiamis, F. Badalamenti, J. Langeneck, P. Noel, A. Deidun, A. Marchini, G. Skouradakis, L. Royo, M. Sini, C.N. Bianchi, Y.-R. Sghaier, R. Ghanem, N. Doumpas, J. Zaouali, K. Tsirintanis, O. Papadakis, C. Morri, M.E. Çinar, J. Terrados, G. Insacco, B. Zava, E. Soufi-Kechaou, L. Piazzzi, C. Ounifi Ben Amor, E. Andriotis, M.C. Gambi, M.M. Ben Amor, J. Garrabou, C. Linares, A. Fortič, M. Diogenis, E. Cebrian, M. Fourt, M. Zotou, L. Castriota, V. Di Martino, A. Rosso, C. Pipitone, M. Falautano, M. García, R. Zakhama-Sraieb, F. Khamassi, A.M. Mannino, M.H. Ktari, I. Kosma, M. Rifi, P.K. Karachle, S. Yapıcı, A.R. Bos, P. Balistreri, A.A. Ramos Esplá, J. Tempesti, O. Inglese, I. Giovos, D. Damalas, S. Benhissoune, M.F. Huseyinoglu, W. Rjiba-Bahri, J. Santamaría, M. Orlando-Bonaca, A. Izquierdo, C. Stamouli, M. Montefalcone, H. Cerim, R. Golo, S. Tsioli, S. Orfanidis, N. Michailidis, M. Gaglioti, E. Taşkın, E. Mancuso, A. Žunec, I. Cvitković, H. Filiz, R. Sanfilippo, A. Siapatis, B. Mavrič, S. Karaa, A. Türker, F. Monniot, J. Verdura, N. El Ouamari, M. Selfati & A. Zenetos (2020): Unpublished Mediterranean records of marine alien and cryptogenic species. *BioInvasions Records*, 9(2), 165-182.
- Maas, O. (1903): Die Scyphomedusen der Siboga Expedition. Siboga Expedition 1901, 1-91.
- Marambio, M., M. Pascual-Torner, U. Tilves, A. Pérez, A. Ballesteros & J.-M. Gili (2025): The westernmost record of the Scyphomedusa *Cassiopea andromeda* (Forskål, 1775) in the Mediterranean: marine citizen science contributions to invasive species detection and monitoring. *Environmental Management*, 75, 3721-3735.
- Pyšek, P., P.E. Hulme, D. Simberloff, S. Bacher, T.M. Blackburn, J.T. Carlton, W. Dawson, F. Essl, L.C. Foxcroft, P. Genovesi, J.M. Jeschke et al. (2020): Scientists' warning on invasive alien species. *Biological Reviews*, 95, 1511-1534.
- Ramos-Pérez, P., J.I. Quispe-Becerra, J.A. Oliver, C. Marcos, A. Pérez-Ruzafa & A. Fernández-Alías (2025): Colonization of the Mediterranean Sea by the Lessepsian invasive jellyfish *Cassiopea andromeda* (Forskål, 1775) – a systematic review. *Mediterranean Marine Science*, 26(3), 714-730.
- Rowe, C.E., S.T. Ahyong, W.F. Figueira, I. Burghardt & S.J. Keable (2025): Identification of *Cassiopea* sp. in Lake Macquarie, Australia and revision of the taxonomic status of *Cassiopea maremetens* Gershwin, Zeidler & Davie, 2010 (Cnidaria: Scyphozoa: Cassiopeidae). *PeerJ*, 13, e19669.
- Schembri, P.J., A. Deidun & P.J. Vella (2010): First record of *Cassiopea andromeda* (Scyphozoa: Rhizostomeae: Cassiopeidae) from the central Mediterranean Sea. *Marine Biodiversity Records*, 3, e6.
- Stock, A., L.B. Crowder, B.S. Halpern & F. Micheli (2018): Uncertainty analysis and robust areas of high and low modeled human impact on the global oceans. *Conservation Biology*, 32(6), 1368-1379.
- Thé, J., E. Gamero-Mora, M.V. Chagas da Silva, A.C. Morandini, S. Rossi & M. de Oliveira Soares (2021): Non-indigenous upside-down jellyfish *Cassiopea andromeda* in shrimp farms (Brazil). *Aquaculture*, 532, 735999.
- Thé, J., M. Mammone, S. Piraino, A. Pennetta, G.E. De Benedetto, T.M. Garcia, M. de Oliveira Soares & S. Rossi (2023): Understanding *Cassiopea andromeda* (Scyphozoa) invasiveness in different habitats: a multiple biomarker comparison. *Water*, 15, 2599.
- Tiralongo, F., F. Crocetta, E. Riginella, A.O. Lillo, E. Tondo, A. Macali, E. Mancini, F. Russo, S. Coco, G. Paolillo & E. Azzurro (2020): Snapshot of rare, exotic and overlooked fish species in the Italian seas: a citizen science survey. *Journal of Sea Research*, 164, 101930.