

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 36, 2026, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 36, 2026, 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Ana Fortič, Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, James Pierson (USA), Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednici/Redattrici/Editors:

Martina Orlando-Bonaca, Ana Fortič

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*[®]
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment* / *Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*[®]

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano* / *National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2026.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base* / *Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza* / *This work is licensed under a Creative Commons BY 4.0.*



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / *INDICE GENERALE* / *CONTENTS*

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Alen SOLDO, Rigers BAKIU & Sherif DURMISHAJ
The First Record of Bastard Grunt *Pomadasys incisus* (Bowdich, 1825) in Albanian Waters (Adriatic Sea) 1
Prvi zapis o pojavljanju neprave prašičevke Pomadasys incisus (Bowdich, 1825) v albanskih vodah (Jadransko morje)

Pero UGARKOVIĆ, Ilija ČETKOVIĆ, Olivera MARKOVIĆ, Aleksandar JOKSIMOVIĆ, Nikola ĐORĐEVIĆ & Jakov DULČIĆ Additional Records of the Bastard Grunt, *Pomadasys incisus* (Bowdich, 1825), from the Eastern Adriatic Sea 7
Dodatne najdbe neprave prašičevke, Pomadasys incisus (Bowdich, 1825), iz vzhodnega Jadranskega morja

Rüştü KIRMAN & Murat BİLECENOĞLU Significant Range Expansion of *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) in the Aegean Sea Based on Scuba and Photographic Observations 15
Znatno razširjanje areala vrste Sepioteuthis lessoniana (Cephalopoda: Loliginidae), v Egejskem morju na podlagi podvodnih opazovanj in fotografiranja

Adrian BRAJKOVIĆ, Iris MATULJA & Neven IVEŠA
Conquering the North: New Occurrence of the Common Lionfish, *Pterois miles* (Bennett, 1828), in the Adriatic Sea 21
Osvajanje severa: novi zapis o pojavljanju navadne plamenke, Pterois miles (Bennett, 1828), v Jadranskem morju

Alan DEIDUN, Bruno ZAVA, Maria CORSINI-FOKA, Arnold SCIBERRAS & Alessio MARRONE First Record of the Devil Firefish *Pterois miles* (Bennett, 1828) (Actinopterygii: Scorpaenidae) from Maltese Coastal Waters 29
Prvi zapis o pojavljanju plamenke Pterois miles (Bennett, 1828) (Actinopterygii: Scorpaenidae) v malteških obalnih vodah

Chirine HUSSEIN, Amir IBRAHIM, Firas ALSHAWY, Mouina BADRAN & Rahaf ABO ASA First Mediterranean Record of the Diagonal Butterflyfish, *Chaetodon fasciatus* Forsskal, 1775, Reported from Syrian Waters 35
Prvi sredozemski zapis o pojavljanju poševnoprogame ščetinozobke, Chaetodon fasciatus Forsskal, 1775, iz sirskih voda

Houssein ELBARAASI, Tarek SHOEIB, Mona SAID & Laith A. JAWAD A Further Record of the Barred Knifejaw, *Oplegnathus fasciatus* (Temminck & Schlegel, 1844), a Pacific Species, from the Mediterranean: A New Record from Benghazi, Libya 41
Novi zapis o vrsti Oplegnathus fasciatus (Temminck & Schlegel, 1844), pacifiški vrsti, v Sredozemlju: nova najdba iz Bengazija v Libiji

Alen SOLDO & Rigers BAKIU Rapid Increase in Records of the Invasive Silver-cheeked Toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Adriatic Sea 49
Hitro naraščanje pojavov invazivne srebrnoproge napihavalke Lagocephalus sceleratus (Gmelin, 1789) v Jadranskem morju

Gerasimos KONDYLATOS, Konstantinos KALAENTZIS, Styliani MINOUDI & Maria CORSINI-FOKA Further Molecular Identification Confirms the Occurrence of *Lagocephalus guentheri* Miranda Ribeiro, 1915 in the Aegean Coastal Waters of Greece 55
Nadaljnja molekularna identifikacija potrjuje prisotnost vrste Lagocephalus guentheri Miranda Ribeiro, 1915 v egejskih obalnih vodah Grčije

Deniz ERGÜDEN & Cem ÇEVİK Length–Weight and Length–Length Relationships, and Condition Factor of *Ambassis dussumieri* Cuvier, 1828, in the Northeastern Mediterranean, Türkiye 65
Dolžinsko-masni odnos med dolžino in kondicijskim faktorjem pri vrsti Ambassis dussumieri Cuvier, 1828, v severovzhodnem Sredozemskem morju, Turčija

Okan AKYOL & HALİL ŞEN Unexpected Occurrence of Prussian Carp *Carassius gibelio* (Cyprinidae) in Homa Lagoon (Izmir Bay, Aegean Sea) 73
Nepričakovani pojav srebrnega koreslja Carassius gibelio (Cyprinidae) v laguni Homa (Izmirski zaliv, Egejsko morje)

SREDOZEMSKE HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
MEDITERRANEAN SHARKS AND RAYS

Terry CARBON, Emily GIGNON, Justine LALLAU-VAZZOLER, Hugo MENARD, Claudio BARRIA, Ana I. COLMENERO & Nicolas ZIANI Precopulatory Behaviour of *Pteroplatytrygon violacea* (Myliobatiformes: Dasyatidae) in the Northwestern Mediterranean 81
Predkopulacijsko vedenje vijoličnega morskega biča (Pteroplatytrygon violacea) (Myliobatiformes: Dasyatidae) v severozahodnem Sredozemlju

Hakan KABASAKAL Distribution of <i>Odontaspis ferox</i> in the Mediterranean Sea: Insights from Spatial and Temporal Analyses 87	MORSKA FAVNA FAUNA MARINA MARINE FAUNA
<i>Razširjenost vrste Odontaspis ferox v Sredozemskem morju: Spoznanja na podlagi prostorskih in časovnih analiz</i>	
Hristina GELEVSKA, Borut MAVRIČ, Lovrenc LIPEJ & Christian CAPAPÉ Is the Gulf of Trieste a potential nursery area for some elasmobranch species? 99	Francesco TIRALONGO, Paola LEOTTA & Riccardo MARTELLUCCI <i>Physalia physalis</i> in the central Mediterranean Sea: Recent observations associated with mass strandings of <i>Velella velella</i> and surface circulation dynamics 165
<i>Ali je Tržaški zaliv potencialno območje odraščanja (jaslice) za določene vrste hrustančnic?</i>	<i>Portugalska ladjica (Physalia physalis) v osrednjem Sredozemskem morju: nedavna opažanja v povezavi z masovnimi nasedanji morskega jadrčka (Velella velella) in dinamiko površinskega kroženja vode</i>
IHTIOFAVNA ITTIOFAUNA ICHTHYOFAUNA	
Cem DALYAN, Yunus GÖNÜL, Mahmud Samed ŞAHİNOĞULLARI & Hakan KABASAKAL New data on the Occurrence and Morphology of the Armless Snake Eel, <i>Dalophis imberbis</i> (Ophichthidae), from the Northeastern Mediterranean Sea 119	Andrea LOMBARDO, Juba CHABANE & Rachida GHALMI First Contribution to the Study of the "Sea Slug" Fauna (Gastropoda, Heterobranchia) of the Algerian coast 177
<i>Novi podatki o pojavljanju in morfologiji kačaste jegulje, Dalophis imberbis (Ophichthidae), iz severovzhodnega Sredozemskega morja</i>	<i>Prvi prispevek k proučevanju favne „goliš morskih polžev“ (Gastropoda, Heterobranchia) alžirske obale</i>
Chirine HUSSEIN, Firas ALSHAWY & Amir IBRAHIM First Record of the Mediterranean Dealfish, <i>Trachipterus trachipterus</i> (Gmelin, 1789) (Trachipteridae), in Syrian Marine Waters 127	Borut MAVRIČ, Tjaša PREMRL, Martin MAVRIČ, Tina MIRT, Neža LEBAN, Tihomir MAKOVEC & Lovrenc LIPEJ Assessing Soft-Bottom Epibenthic Communities: Methodological Insights into Dredging and Video Surveys 203
<i>Prvi zapis o pojavljanju kosice, Trachipterus trachipterus (Gmelin, 1789) (Trachipteridae), v sirskih morskih vodah</i>	<i>Ocenjevanje epibentoških združb mehkega dna: metodološki vpogledi v dredžanje in video pregleda</i>
Igor AGOSTINI & Okan AKYOL On the Occurrence of the Currently Largest Recorded <i>Pagellus erythrinus</i> (Sparidae) in the Tyrrhenian Sea (Italy) 133	FAVNA FAUNA FAUNA
<i>O pojavljanju trenutno največjega evidentiranega primerka vrste Pagellus erythrinus (Sparidae) v Tirenskem morju (Italija)</i>	Kim LEBAN, Iztok ŠKORNIK, Špela ČONČ, Mateja BREG VALJAVEC, Žan KURALT, Lenart ŠTAUT, Jure TIČAR, Katarina POLAJNAR HORVAT, Primož GAŠPERIČ, Aleš SMREKAR & Melita VAMBERGER Habitat Insights for the European Pond Turtle (<i>Emys orbicularis</i>) in the Coastal Wetland Area Sečovlje Salina Nature Park, Slovenia 221
Ismail Burak DABAN & Yusuf ŞEN Substantiated Record and Preliminary Biological Insights of <i>Microlophrys dalmatinus</i> (Steindachner & Kolombatović, 1883) from the Sea of Marmara, Türkiye 141	<i>Habitatne značilnosti močvirske sklednice (Emys orbicularis) v krajinskem parku Sečoveljske soline, Slovenija</i>
<i>Potrjen zapis o pojavljanju in predhodna biološka spoznanja o vrsti Microlophrys dalmatinus (Steindachner & Kolombatović, 1883) iz Marmarskega morja, Turčija</i>	Leon CIMERMAN & Matija KRIŽNAR Najdbe pleistocenske avifavne v kamnolomu Črni Kal (Primorska, Slovenija) 243
Zeliha ERDOĞAN, Gülçin ULUNEHIR AYDIN, Hatice TORCU-KOÇ & Tuğba ANBAROĞLU Population Study of <i>Trachurus mediterraneus</i> Focused on Reproductive Biology in Edremit Bay, Northern Aegean Sea 151	<i>Pleistocene Avifauna Finds in the Črni Kal Quarry (Primorska Region, Slovenia)</i>
<i>Razmnoževalna biologija sredozemskega šura (Trachurus mediterraneus) v sklopu populacijske raziskave v Edremitnem zalivu (severno Egejsko morje)</i>	Kazalo k slikam na ovitku 255
	Index to images on the cover 255

received: 2026-02-03

DOI 10.19233/ASHN.2026.15

NEW DATA ON THE OCCURRENCE AND MORPHOLOGY OF THE ARMLESS SNAKE EEL, *DALOPHIS IMBERBIS* (OPHICHTHIDAE), FROM THE NORTHEASTERN MEDITERRANEAN SEA

Cem DALYAN

Department of Biology, Division of Hydrobiology, Faculty of Science, Istanbul University, Istanbul, Türkiye

Yunus GÖNÜL & Mahmud Samed ŞAHİNOĞULLARI

Department of Biology, Institute of Graduate Studies in Sciences, Istanbul University, Istanbul, Türkiye

Hakan KABASAKAL

Hydrobiological Research Association, Gökçeada, Çanakkale, Türkiye
e-mail: kabasakal.hakan@gmail.com

ABSTRACT

The armless snake eel, *Dalophis imberbis*, is an ophichthid eel widely distributed in the Mediterranean Sea, but comparatively rare in the eastern basin. This study examined three specimens from Iskenderun Bay (northeastern Mediterranean Sea, Türkiye), which were identified based on standard morphological diagnostic characters and measured for key morphometric traits. The individuals were captured at shallow coastal depths, contributing new distributional and ecological data for the species in the eastern Levant Basin. Morphometric results are consistent with previously published Mediterranean data, although minor variations were observed among selected proportional measurements. These findings provide additional evidence for the presence of *D. imberbis* in Turkish waters and support the view that the species forms part of the established ichthyofauna of the eastern Mediterranean.

Key words: Anguilliformes, Ophichthidae, Levant Basin, occurrence

NUOVI DATI SULLA PRESENZA E LA MORFOLOGIA DELLA BISCIA DI MARE, *DALOPHIS IMBERBIS* (OPHICHTHIDAE), NEL MEDITERRANEO NORD-ORIENTALE

SINTESI

La biscia di mare, *Dalophis imberbis*, è un'anguilla della famiglia Ophichthidae ampiamente distribuita nel Mediterraneo, ma relativamente rara nel bacino orientale. In questo studio gli autori hanno esaminato tre esemplari provenienti dalla baia di Iskenderun (Mediterraneo nord-orientale, Türkiye), identificati sulla base dei caratteri diagnostici morfologici standard e misurati per i principali parametri morfometrici. Gli individui sono stati catturati in acque costiere poco profonde, fornendo nuovi dati distributivi ed ecologici per la specie nel bacino levantino orientale. I risultati morfometrici sono coerenti con i dati mediterranei precedentemente pubblicati. L'articolo fornisce ulteriori prove della presenza di *D. imberbis* nelle acque turche e supporta l'ipotesi che la specie faccia parte dell'ittiofauna stabilmente insediata del Mediterraneo orientale.

Parole chiave: Anguilliformes, Ophichthidae, Bacino del Levante, presenza

INTRODUCTION

The family Ophichthidae includes 62 genera and 374 species, among which is the armless snake eel, *Dalophis imberbis* (Delaroche, 1809) (Fricke *et al.*, 2026). The distribution range of *D. imberbis* extends from Mauritania to Spain and throughout the Mediterranean Sea (Froese & Pauly, 2026), where the species occurs at depths ranging from 5.4 m (Sabrina Lo Brutto, pers. comm.) to 115 m (Busalacchi *et al.*, 2010). *D. imberbis* is considered an established member of the Mediterranean anguilliform fauna (Kovačić *et al.*, 2021) and has been found in the Turkish Aegean and Mediterranean waters (Bilecenoğlu, 2024). However, most occurrence records originate from the western and central Mediterranean basins (e.g., Italian waters, Bonifazi *et al.*, 2019; Libyan waters, Elbaraasi *et al.*, 2019; Adriatic Sea, Soldo & Bakiu, 2021; Algerian waters, Alkhalili *et al.*, 2025). In contrast, its presence in the eastern Mediterranean has been documented only through sporadic records (e.g., Israeli waters, Golani, 2005; Syrian waters, Capapé *et al.*, 2021). Despite the species' broad distribution in the Mediterranean Sea, information on its occurrence and biological characteristics in the eastern Mediterranean remains limited. Therefore, the present study provides new data on the distribution and morphometric characteristics of *D. imberbis* in the northeastern Mediterranean Sea based on three specimens collected from Iskenderun Bay.

MATERIAL AND METHODS

Three specimens of *Dalophis imberbis* were captured in Iskenderun Bay (Fig. 1) by a commercial otter trawler (M/V Faik Baba; 400 hp) fitted with a net with a cod-end mesh size of 44 mm (stretched mesh, knot-to-knot) and a net opening width of 20 m. Trawling operations were conducted over a mixed mud–sand bottom at depths ranging from 54 to 69 m, at towing speeds of 2.2 to 2.8 knots. The specimens were captured at the following locations and depths: specimen no. 1 was caught at a depth of 63 m (36°47'4.5"–36°41'15.7" N; 36°6'43.2"–36°5'19.7" E); specimen no. 2 at 61 m (36°41'43.8"–36°44'47.0" N; 36°4'52.3"–36°3'30.7" E); and specimen no. 3 at 55 m (36°34'1.5"–36°34'57.8" N; 36°1'9.7"–36°3'14.4" E). Sampling was performed on 25 and 26 February 2013.

After capture, the specimens were kept frozen at –18 °C on board until fixation and were subsequently preserved in a 5% borax-buffered formalin solution. Species identification followed Bauchot (1986) and the scientific nomenclature conforms to the Fricke *et al.* (2026). Previous records of *D. imberbis* covering the period 1845–2026 were

obtained from the Global Biodiversity Information Facility (GBIF, 2026), available at the following web page: <https://www.gbif.org/species/2405106>. The coordinates of these records were plotted using cartographic software QGIS version 3.40 to generate a distribution map of the species throughout the Mediterranean Sea (Fig. 1). The present description is based on the examination of the three specimens from Iskenderun Bay, which are deposited in the museum of the Department of Hydrobiology, Istanbul University, under catalogue numbers 2013-1052, 2013-1053, and 2013-1054.

RESULTS AND DISCUSSION

Description of the examined specimens (nos. 1–3, Fig. 2): body extremely cylindrical and elongated, snout pointed, lower jaw symphysis closer to the anterior border of the eye than to the tip of the snout, eyes minute and covered by skin; anterior nostrils short and tube-shaped, posterior nostrils located above upper lip; teeth conical and slightly curved, uniserial in both jaws, five premaxillary teeth forming a V-shape, small teeth present on vomer; gill openings ranging from midlateral to entirely ventral, crescent-shaped; all fins absent; caudal tip hardened; lateral line with 62–63 preanal pores and 7–8 prebranchial pores; body depth 2.2–2.5% of total length, head length 7.5–8.0% of total length; snout length 18.2–21.2%, eye diameter 5.9–8.1%, and interorbital width 8.0–10.2% of head length; colouration: dorsal surface brownish, tip of upper jaw dark brown, ventral surface whitish. The characteristics of the examined specimens match those reported in the literature, confirming the identification of the collected individuals as *Dalophis imberbis* (Delaroche, 1809). Their length measurements are presented in Tab. 1.

In light of the available literature, *D. imberbis* is considered a native member of the Mediterranean ichthyofauna, exhibiting a broad but highly patchy distribution across the basin. The apparent scarcity of records in several areas is most likely attributable to the species' cryptic behaviour, burrowing lifestyle, and the limited efficiency of conventional sampling methods targeting benthic eel-like species, rather than reflecting true absence. *D. imberbis* is rarely reported off the French coast (Bauchot & Pras, 1980; Béarez *et al.*, 2017), while it appears relatively more frequently in Italian waters (Bonifazi *et al.*, 2019). In the Adriatic Sea, it occurs at low densities (Likić *et al.*, 2015). Further south, spawning activity for the species has been reported from the Bay of Algiers (Bauchot, 1986), whereas its presence along the Maghreb coast remains poorly documented, including in Algeria (Refes *et al.*, 2010) and Tunisia (Bradäi *et al.*, 2004). This likely reflects undersampling rather than true ab-




Fig. 1: Previous records (solid black circles) and present specimens (solid red circles, indicated by a solid red arrow) of *Dalophis imberbis* throughout the Mediterranean Sea (a), and a detailed map of the capture localities of the present specimens in Iskenderun Bay (b). The blue bathymetric colour gradient represents depth variation from 0 to 1000 m, from light to dark blue. The isobaths indicate depth contours at 200 m and 1000 m, progressing from the outer to the inner limits of the mapped area.

Sl. 1: Prejšnji zapisi o pojavljanju vrste *Dalophis imberbis* po celotnem Sredozemskem morju (polni črni krogi) in pričujoči osebk (polni rdeči krogi, označeni s polno rdečo puščico) (a) ter podroben zemljevid lokacij ulova pričujočih osebkov v zalivu Iskenderun (b). Modri batimetrični barvni gradient predstavlja spreminjanje globine od 0 do 1000 m, od svetlo do temno modre. Izobate označujejo globinske konture pri 200 m in 1000 m, ki potekajo od zunanjih proti notranjim mejam usmerjenega območja.



Fig. 2: Lateral view of *Dalophis imberbis*; scale (tape measure units) = 10 mm (a). Lateral view of the head of the same specimen; scale (tape measure units) = 10 mm (b). Dentition of the lower jaw; scale bar = 1 mm (c). Photo: Cem Dalyan.

Sl. 2: Stranski pogled na primerek *Dalophis imberbis*; merilo (enote na merilnem traku) = 10 mm (a). Stranski pogled na glavo istega primerka; merilo (enote na merilnem traku) = 10 mm (b). Zobovje spodnje čeljusti; merilo = 1 mm (c). Foto: Cem Dalyan.

sence. Eastwards, records from Greek (Papaconstantinou, 2014) and Turkish waters (Bilecenoğlu, 2024) confirm a continuous occurrence of the species, with the Levant Basin representing the easternmost extent of its distribution (Golani, 2005; El Sayed *et al.*, 2017; Bariche & Fricke, 2020). Therefore, *D. imberbis* should be regarded as a native but under-detected species in the eastern Mediterranean, with long-term monitoring occasionally revealing previously unrecorded occurrences in the region (Saad, 2005; Ali, 2018; Capapé *et al.*, 2021).

However, records derived from citizen science observations, particularly those including georeferenced photographs accessible through the GBIF platform (GBIF, 2026), have significantly improved the resolution of the known distribution of *D. imberbis* in recent years. Still, these opportunistic data cannot replace systematic scientific surveys, which remain essential to address persistent gaps in knowledge regarding the species' life history and population ecology. In this context, the most detailed morphometric study of *D. imberbis* to date has been conducted by Bonifazi *et al.* (2019), who reported selected measurements of 20 specimens out of approximately 200 stranded individuals in

the central Tyrrhenian Sea. More recently, Capapé *et al.* (2021) presented morphometric data based on a single specimen from Syrian waters. Although the specimens presented in this study were collected several years prior to their detailed examination, their morphometric data remain valuable for the understanding of *D. imberbis* populations in the eastern Mediterranean. In fact, by reporting measurements of three further specimens captured in the eastern Mediterranean Sea, this study contributes additional morphometric information and enhances knowledge of the species' morphology and distribution in the region.

Although based on a limited number of specimens, the morphometric results of this study provide comparative data for *D. imberbis* from the eastern Mediterranean. Overall, proportional measurements show a high degree of consistency with previous studies, particularly for head length (7.50% TL in the present study vs. 7.03% in Bonifazi *et al.* (2019) and 7.73% in Capapé *et al.* (2021)) and eye diameter (0.52% TL vs. 0.41% and 0.50%, respectively). Similarly, body depth values (2.34% TL) closely match those reported by Bonifazi *et al.* (2019) (2.26% TL), suggesting limited morphological

Tab. 1: Morphometric data of the examined *Dalophis imberbis* specimens (nos. 1–3), presented as individual measurements and percentages of total length (TL). Data from relevant literature are also included for comparison. SE: standard error of the mean; N/A: not available.

Tab. 1: Morfometrični podatki preučenih primerkov vrste *Dalophis imberbis* (št. 1–3), predstavljeni kot posamezne meritve in odstotki celotne dolžine (TL). Za primerjavo so vključeni tudi podatki iz ustrezne literature. SE: standardna napaka aritmetične sredine; N/A: ni na voljo.

Measurements (mm)	Sp. no 1	Sp. no 2	Sp. no 3	Mean	±SE	% of mean TL	Bonifazi <i>et al.</i> (2019) % of mean TL	Capapé <i>et al.</i> (2021) % of mean TL
TL	234	261	208	234.3	12.5	100	100	100
Preal length	100.7	117	88	101.9	6.9	43.5	N/A	37.5
Predorsal length	N/A	N/A	N/A	N/A	N/A	N/A	N/A	9.7
Prepectoral length	N/A	N/A	N/A	N/A	N/A	N/A	N/A	8.4
Body depth	5.9	5.6	4.9	5.5	0.3	2.3	2.3	1.9
Head length	18.6	18.6	15.5	17.6	0.8	7.5	7.0	7.7
Eye diameter	1.5	1.1	1.1	1.2	0.1	0.5	0.4	0.5
Preorbital length	N/A	N/A	N/A	N/A	N/A	N/A	N/A	2.4
Snout length	3.4	3.9	3.3	3.5	0.2	1.5	1.3	N/A
Upper jaw length	6.3	6.6	4.4	5.8	0.6	2.5	2.3	N/A
Lower jaw length	5.3	5.8	3.6	4.9	0.6	2.1	N/A	4.9
Interorbital distance	1.8	1.5	1.6	1.6	0.1	0.7	N/A	0.6

variability across Mediterranean populations. Some differences, however, were observed in specific measurements, particularly preanal length, which was slightly higher in the present material (43.49% TL) compared to Capapé *et al.* (2021) (37.50% TL). Such variation may be a reflection of intraspecific variability, limited sample size, or methodological differences among studies, including measurement protocols and specimen preservation conditions. In particular, long-term preservation history, including prolonged freezing, may induce minor tissue shrinkage or deformation, potentially affecting linear morphometric ratios. The absence of several morphometric parameters in previous studies (e.g., predorsal, prepectoral, and preorbital lengths in

Bonifazi *et al.* (2019)) further limits direct comparisons but also highlights the added value of the present dataset in complementing the morphological knowledge of the species.

From a broader perspective, the limited availability of morphometric data for *D. imberbis* likely reflects its low economic importance and, consequently, the lack of research priority given to a species not targeted by commercial fisheries. Such situations often result in reliance on opportunistic sampling and incidental captures, with specimens frequently deposited in collections and examined only years later, when taxonomic or faunistic interest arises. In this context, a 13-year interval between collection and detailed analysis, as in the

present case, is not unusual for non-commercial, rarely studied benthic species. Museum and institutional collections are thus particularly important, as they provide essential material for retrospective morphological and faunistic studies of poorly known taxa. Such archived specimens allow the reconstruction of distributional and morphological baselines that would otherwise remain undocumented, especially for cryptic and low-priority species such as *D. imberbis*.

ACKNOWLEDGEMENTS

The present study was supported by the Research Fund of Istanbul University (project no. 27457). We thank three anonymous reviewers for their constructive comments, which improved the manuscript. Special thanks are due to Prof. Dr. Sabrina Lo Brutto (University of Palermo, Italy) for kindly sharing information on the unusually shallow depth record of *Dalophis imberbis*, cited herein as personal communication.

NOVI PODATKI O POJAVLJANJU IN MORFOLOGIJI KAČASTE JEGULJE, *DALOPHIS IMBERBIS* (OPHICHTHIDAE), IZ SEVEROVZHODNEGA SREDOZEMSKEGA MORJA

Cem DALYAN

Department of Biology, Division of Hydrobiology, Faculty of Science, Istanbul University, Istanbul, Türkiye

Yunus GÖNÜL & Mahmud Samed ŞAHİNOĞULLARI

Department of Biology, Institute of Graduate Studies in Sciences, Istanbul University, Istanbul, Türkiye

Hakan KABASAKAL

Hydrobiological Research Association, Gökçeada, Çanakkale, Türkiye
e-mail: kabasakal.hakan@gmail.com

POVZETEK

Kačasta jegulja, *Dalophis imberbis*, je vrsta iz družine kačastih jegulj (*Ophichthidae*), ki je široko razširjena v Sredozemskem morju, vendar razmeroma redka v njegovem vzhodnem bazenu. Avtorji so preučili tri osebkke iz zaliva Iskenderun (severovzhodno Sredozemsko morje, Turčija), ki so jih določili na podlagi standardnih morfoloških diagnostičnih znakov, izmerjene pa so bile tudi njihove ključne morfometrične lastnosti. Primerke so ujeli v obalnih plitvinah, kar prinaša nove podatke o razširjenosti in ekologiji te vrste v vzhodnem Levantskem bazenu. Rezultati morfometričnih meritev so skladni s predhodno objavljenimi podatki za Sredozemlje, čeprav so bila med izbranimi proporcionalnimi meritvami opažena manjša odstopanja. Te ugotovitve zagotavljajo dodatne dokaze o prisotnosti vrste *D. imberbis* v turških vodah in podpirajo stališče, da ta vrsta predstavlja del ustaljene ihtiofavne vzhodnega Sredozemlja.

Ključne besede: Anguilliformes, Ophichthidae, Levantski bazen, pojavljanje

REFERENCES

- Ali, M. (2018):** An updated Checklist of the Marine fishes from Syria with emphasis on alien species. *Medit. Mar. Sci.*, 19, 388–393.
- Alkhalili, H., A.A. Taleb Bendiab, B. Seret, L. Bensahla-Talet, N. Benaissa & M. Bouderbala (2025):** Bony fish of Algerian marine fauna: a historical review with an updated checklist and new records. *Appl. ecol. environ. res.*, 23, 7957–7995. http://dx.doi.org/10.15666/aeer/2304_79577995.
- Bariche, M. & R. Fricke (2020):** The marine ichthyofauna of Lebanon: an annotated checklist, history, biogeography, and conservation status. *Zootaxa*, 4775, 1–157.
- Bauchot, M.L. (1986):** Ophichthidae (including Echelidae, Clofnam 84). In: P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese (eds.), *Fishes of the North-western Atlantic and the Mediterranean*, Vol II, UNESCO, Paris, pp. 577–585.
- Bauchot, M. L. & A. Pras (1980):** Guide des poissons marins d'Europe. Delachaux & Niestlé. Lausanne-Paris, 427 pp.
- Béarez, P., P. Pruvost, É. Feunteun, S. Iglésias, P. Francour, R. Causse, J. De Mazières, S. Tercerie & N. Bailly (2017):** Checklist of the marine fishes from metropolitan France. *Cybum*, 41(4), 351–371.
- Bilecenoglu, M. (2024):** Diversity of fishes along the coast of Türkiye. *Turk. J. Zool.*, 48, 589–616. <https://doi.org/10.55730/1300-0179.3197>.
- Bonifazi, A., F. Fratini & D. Ventura (2019):** Unusual massive beaching of *Dalophis imberbis* (Delaroche, 1809) (Anguilliformes, Ophichthidae) from the central Tyrrhenian Sea. *Thalassas*, 35(2), 577–586.
- Bradaï, M. N., J.P. Quignard, A. Bouaïn, O. Jarboui, A. Ouannes-Ghorbel, L. Ben Abdallah, J. Zaouali & S. Ben Salem (2004):** Ichtyofaune Autochtone Et Exotique des côtes tunisiennes: recensement et biogéographie. *Cybum*, 28, 315–328.
- Busalacchi, B., P. Rinelli, F. De Domenico, A. Profeta, F. Perdichizzi & T. Bottari (2010):** Analysis of demersal fish assemblages off the southern Tyrrhenian Sea (central Mediterranean). *Hydrobiologia*, 654, 111–124. <https://doi.org/10.1007/s10750-010-0374-9>.
- Capapé, C., A. Saad, A. Solaiman, I. Barakat & W. Sabour (2021):** First substantiated record of armless snake eel *Dalophis imberbis* (Osteichthyes: Ophichthidae) from the Syrian coast (Eastern Mediterranean Sea). *Annales, Ser. Hist. Nat.*, 31, 101–106. <https://doi.org/10.19233/ASHN.2021.12>
- Elbaraasi, H., B. Elabar, S. Elaabidi, A. Bashir, O. Elsilini, E. Shakman & E. Azzurrio (2019):** Updated checklist of bony fishes along the Libyan coast (southern Mediterranean Sea). *Medit. Mar. Sci.*, 20, 90–105.
- El Sayed, H., K. Akel. & P.K. Karachle (2017):** The marine ichthyofauna of Egypt. *Egyptian J. Aquat. Biol. Fish.*, 21(3), 81–116.
- Fricke, R., W.N. Eschmeyer & R. Van der Laan (eds.) (2026):** Eschmeyer's Catalog of Fishes: Genera, Species, References. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp> (accessed 5 May 2026).
- Froese, R. & D. Pauly (eds.) (2026):** FishBase. World Wide Web electronic publication. www.fishbase.org, version. (accessed 5 May 2026).
- GBIF (2026):** Global Biodiversity Information Facility. Available at: <https://www.gbif.org/species/2405106>. (accessed 5 May 2026).
- Golani, D. (2005):** Check-list of the Mediterranean Fishes of Israel. *Zootaxa*, 2005(947), 1–200.
- Kovačić, M., L. Lipej, J. Dulčić, S. Iglesias & M. Goren (2021):** Evidence-based checklist of the Mediterranean Sea fishes. *Zootaxa*, 4998, 1–115. <https://doi.org/10.11646/zootaxa.4998.1.1>.
- Likić, J., T. Safner, P. Peles, M. Dobos, B. Božić, M. Gredelj, M. Mesarić, D. Delić & I. Gudac (2015):** A strategic study of the likely significant environmental impact of the framework plan and programme of exploration and production of hydrocarbons in the Adriatic. Ministry of the Economy of the Republic of Croatia, 450 pp.
- Papaconstantinou, C. (2014):** Fauna Graeciae: an updated list the fishes of the Hellenic Seas. Biodiversity Conservation Zoology. Hellenic Center for Marine Research (HCMR), 340 pp.
- Refes, W., N. Semahi, M. Boulahdid & J.-P. Quignard (2010):** Inventaire de la faune ichthyologique du secteur oriental de la côte algérienne (El Kala; Skikda; Jijel; Bejaïa). *Rapports de la Commission internationale pour la mer Méditerranée*, 39, 646.
- Saad, A. (2005):** Check-list of bony fish collected from the coast of Syria. *Turk. J. Fish. Aquat. Sci.*, 5(2), 99–106.
- Soldo, A. & R. Bakiu (2021):** Checklist of marine fishes of Albania. *Acta Adriat.*, 62, 63–73.