

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 36, 2026, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 36, 2026, 1

ISSN 1408-533X
e-ISSN 2591-1783

UDK 5

Letnik 36, leto 2026 številka 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Ana Fortič, Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, James Pierson (USA), Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednici/Redattrici/Editors:

Martina Orlando-Bonaca, Ana Fortič

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:Zgodovinsko društvo za južno Primorsko - Koper / Società storica del Litorale - Capodistria®
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment / Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente®**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**Nacionalni inštitut za biologijo, Morska biološka postaja Piran / Istituto nazionale di biologia, Stazione di biologia marina di Pirano / National Institute of Biology, Marine Biology Station Piran
SI-6330 Piran / Pirano, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2026.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / La rivista Annales, series Historia Naturalis è inserita nei seguenti data base / Articles appearing in this journal are abstracted and indexed in: BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / Quest'opera è distribuita con Licenza / This work is licensed under a Creative Commons BY 4.0.

Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>

VSEBINA / *INDICE GENERALE* / *CONTENTS*

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Alen SOLDÓ, Rigers BAKIU & Sherif DURMISHAJ
The First Record of Bastard Grunt *Pomadasys incisus* (Bowdich, 1825) in Albanian Waters (Adriatic Sea) 1
Prvi zapis o pojavljanju neprave prašičevke Pomadasys incisus (Bowdich, 1825) v albanskih vodah (Jadransko morje)

Pero UGARKOVIĆ, Ilija ČETKOVIĆ, Olivera MARKOVIĆ, Aleksandar JOKSIMOVIĆ, Nikola ĐORĐEVIĆ & Jakov DULČIĆ Additional Records of the Bastard Grunt, *Pomadasys incisus* (Bowdich, 1825), from the Eastern Adriatic Sea 7
Dodatne najdbe neprave prašičevke, Pomadasys incisus (Bowdich, 1825), iz vzhodnega Jadranskega morja

Rüştü KIRMAN & Murat BİLECENOĞLU Significant Range Expansion of *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) in the Aegean Sea Based on Scuba and Photographic Observations 15
Znatno razširjanje areala vrste Sepioteuthis lessoniana (Cephalopoda: Loliginidae), v Egejskem morju na podlagi podvodnih opazovanj in fotografiranja

Adrian BRAJKOVIĆ, Iris MATULJA & Neven IVEŠA
Conquering the North: New Occurrence of the Common Lionfish, *Pterois miles* (Bennett, 1828), in the Adriatic Sea 21
Osvajanje severa: novi zapis o pojavljanju navadne plamenke, Pterois miles (Bennett, 1828), v Jadranskem morju

Alan DEIDUN, Bruno ZAVA, Maria CORSINI-FOKA, Arnold SCIBERRAS & Alessio MARRONE First Record of the Devil Firefish *Pterois miles* (Bennett, 1828) (Actinopterygii: Scorpaenidae) from Maltese Coastal Waters 29
Prvi zapis o pojavljanju plamenke Pterois miles (Bennett, 1828) (Actinopterygii: Scorpaenidae) v malteških obalnih vodah

Chirine HUSSEIN, Amir IBRAHIM, Firas ALSHAWY, Mouina BADRAN & Rahaf ABO ASA First Mediterranean Record of the Diagonal Butterflyfish, *Chaetodon fasciatus* Forsskal, 1775, Reported from Syrian Waters 35
Prvi sredozemski zapis o pojavljanju poševnoprogame ščetinozobke, Chaetodon fasciatus Forsskal, 1775, iz sirskih voda

Houssein ELBARAASI, Tarek SHOEIB, Mona SAID & Laith A. JAWAD A Further Record of the Barred Knifejaw, *Oplegnathus fasciatus* (Temminck & Schlegel, 1844), a Pacific Species, from the Mediterranean: A New Record from Benghazi, Libya 41
Novi zapis o vrsti Oplegnathus fasciatus (Temminck & Schlegel, 1844), pacifiški vrsti, v Sredozemlju: nova najdba iz Bengazija v Libiji

Alen SOLDÓ & Rigers BAKIU Rapid Increase in Records of the Invasive Silver-cheeked Toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Adriatic Sea 49
Hitro naraščanje pojavov invazivne srebrnoproge napihvalke Lagocephalus sceleratus (Gmelin, 1789) v Jadranskem morju

Gerasimos KONDYLATOS, Konstantinos KALAENTZIS, Styliani MINOUDI & Maria CORSINI-FOKA Further Molecular Identification Confirms the Occurrence of *Lagocephalus guentheri* Miranda Ribeiro, 1915 in the Aegean Coastal Waters of Greece 55
Nadaljnja molekularna identifikacija potrjuje prisotnost vrste Lagocephalus guentheri Miranda Ribeiro, 1915 v egejskih obalnih vodah Grčije

Deniz ERGÜDEN & Cem ÇEVİK Length–Weight and Length–Length Relationships, and Condition Factor of *Ambassis dussumieri* Cuvier, 1828, in the Northeastern Mediterranean, Türkiye 65
Dolžinsko-masni odnos med dolžino in kondicijskim faktorjem pri vrsti Ambassis dussumieri Cuvier, 1828, v severovzhodnem Sredozemskem morju, Turčija

Okan AKYOL & HALİL ŞEN Unexpected Occurrence of Prussian Carp *Carassius gibelio* (Cyprinidae) in Homa Lagoon (Izmir Bay, Aegean Sea) 73
Nepričakovani pojav srebrnega koreslja Carassius gibelio (Cyprinidae) v laguni Homa (Izmirski zaliv, Egejsko morje)

SREDOZEMSKE HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
MEDITERRANEAN SHARKS AND RAYS

Terry CARBON, Emily GIGNON, Justine LALLAU-VAZZOLER, Hugo MENARD, Claudio BARRIA, Ana I. COLMENERO & Nicolas ZIANI Precopulatory Behaviour of *Pteroplatytrygon violacea* (Myliobatiformes: Dasyatidae) in the Northwestern Mediterranean 81
Predkopulacijsko vedenje vijoličnega morskega biča (Pteroplatytrygon violacea) (Myliobatiformes: Dasyatidae) v severozahodnem Sredozemlju

Hakan KABASAKAL Distribution of <i>Odontaspis ferox</i> in the Mediterranean Sea: Insights from Spatial and Temporal Analyses 87	MORSKA FAVNA FAUNA MARINA MARINE FAUNA
<i>Razširjenost vrste Odontaspis ferox v Sredozemskem morju: Spoznanja na podlagi prostorskih in časovnih analiz</i>	
Hristina GELEVSKA, Borut MAVRIČ, Lovrenc LIPEJ & Christian CAPAPÉ Is the Gulf of Trieste a potential nursery area for some elasmobranch species? 99	Francesco TIRALONGO, Paola LEOTTA & Riccardo MARTELLUCCI <i>Physalia physalis</i> in the central Mediterranean Sea: Recent observations associated with mass strandings of <i>Velella velella</i> and surface circulation dynamics 165
<i>Ali je Tržaški zaliv potencialno območje odraščanja (jaslice) za določene vrste hrustančnic?</i>	<i>Portugalska ladjica (Physalia physalis) v osrednjem Sredozemskem morju: nedavna opažanja v povezavi z masovnimi nasedanji morskega jadrčka (Velella velella) in dinamiko površinskega kroženja vode</i>
IHTIOFAVNA ITTIOFAUNA ICHTHYOFAUNA	
Cem DALYAN, Yunus GÖNÜL, Mahmud Samed ŞAHİNOĞULLARI & Hakan KABASAKAL New data on the Occurrence and Morphology of the Armless Snake Eel, <i>Dalophis imberbis</i> (Ophichthidae), from the Northeastern Mediterranean Sea 119	Andrea LOMBARDO, Juba CHABANE & Rachida GHALMI First Contribution to the Study of the "Sea Slug" Fauna (Gastropoda, Heterobranchia) of the Algerian coast 177
<i>Novi podatki o pojavljanju in morfologiji kačaste jegulje, Dalophis imberbis (Ophichthidae), iz severovzhodnega Sredozemskega morja</i>	<i>Prvi prispevek k proučevanju favne „goliš morskih polžev“ (Gastropoda, Heterobranchia) alžirske obale</i>
Chirine HUSSEIN, Firas ALSHAWY & Amir IBRAHIM First Record of the Mediterranean Dealfish, <i>Trachipterus trachipterus</i> (Gmelin, 1789) (Trachipteridae), in Syrian Marine Waters 127	Borut MAVRIČ, Tjaša PREMRL, Martin MAVRIČ, Tina MIRT, Neža LEBAN, Tihomir MAKOVEC & Lovrenc LIPEJ Assessing Soft-Bottom Epibenthic Communities: Methodological Insights into Dredging and Video Surveys 203
<i>Prvi zapis o pojavljanju kosice, Trachipterus trachipterus (Gmelin, 1789) (Trachipteridae), v sirskih morskih vodah</i>	<i>Ocenjevanje epibentoških združb mehkega dna: metodološki vpogledi v dredžanje in video pregleda</i>
Igor AGOSTINI & Okan AKYOL On the Occurrence of the Currently Largest Recorded <i>Pagellus erythrinus</i> (Sparidae) in the Tyrrhenian Sea (Italy) 133	FAVNA FAUNA FAUNA
<i>O pojavljanju trenutno največjega evidentiranega primerka vrste Pagellus erythrinus (Sparidae) v Tirenskem morju (Italija)</i>	Kim LEBAN, Iztok ŠKORNIK, Špela ČONČ, Mateja BREG VALJAVEC, Žan KURALT, Lenart ŠTAUT, Jure TIČAR, Katarina POLAJNAR HORVAT, Primož GAŠPERIČ, Aleš SMREKAR & Melita VAMBERGER Habitat Insights for the European Pond Turtle (<i>Emys orbicularis</i>) in the Coastal Wetland Area Sečovlje Salina Nature Park, Slovenia 221
Ismail Burak DABAN & Yusuf ŞEN Substantiated Record and Preliminary Biological Insights of <i>Microlipophrys dalmatinus</i> (Steindachner & Kolombatović, 1883) from the Sea of Marmara, Türkiye 141	<i>Habitatne značilnosti močvirske sklednice (Emys orbicularis) v krajinskem parku Sečoveljske soline, Slovenija</i>
<i>Potrjen zapis o pojavljanju in predhodna biološka spoznanja o vrsti Microlipophrys dalmatinus (Steindachner & Kolombatović, 1883) iz Marmarskega morja, Turčija</i>	Leon CIMERMAN & Matija KRIŽNAR Najdbe pleistocenske avifavne v kamnolomu Črni Kal (Primorska, Slovenija) 243
Zeliha ERDOĞAN, Gülçin ULUNEHIR AYDIN, Hatice TORCU-KOÇ & Tuğba ANBAROĞLU Population Study of <i>Trachurus mediterraneus</i> Focused on Reproductive Biology in Edremit Bay, Northern Aegean Sea 151	<i>Pleistocene Avifauna Finds in the Črni Kal Quarry (Primorska Region, Slovenia)</i>
<i>Razmnoževalna biologija sredozemskega šura (Trachurus mediterraneus) v sklopu populacijske raziskave v Edremitnem zalivu (severno Egejsko morje)</i>	Kazalo k slikam na ovitku 255
	Index to images on the cover 255

received: 2026-03-09

DOI 10.19233/ASHN.2026.10

LENGTH-WEIGHT AND LENGTH-LENGTH RELATIONSHIPS, AND CONDITION FACTOR OF *AMBASSIS DUSSUMIERI* CUVIER, 1828, IN THE NORTHEASTERN MEDITERRANEAN, TÜRKIYE

Deniz ERGÜDEN

Marine Science Department, Faculty of Marine Science and Technology, Iskenderun Technical University, 31200 Iskenderun, Hatay, Türkiye
e-mail: deniz.erguden@iste.edu.tr; derguden@gmail.com

Cem ÇEVİK

Faculty of Fisheries, Çukurova University Balcalı Campus, 01330, Sarıçam/Adana, Türkiye

ABSTRACT

*This study provides the first data on the length–weight relationship (LWR), length–length relationship (LLR), and condition factor (K) of the Malabar glassy perchlet, *Ambassis dussumieri*, in the northeastern Mediterranean (Tuzla coast, Türkiye). A total of 132 specimens (68 males and 64 females) were examined. The specimens ranged from 4.5 to 8.9 cm in total length (TL), 3.7 to 7.7 cm in fork length (FL), 3.4 to 6.9 cm in standard length (SL) for both sexes. The calculated *b* values indicated positive allometric growth for the overall population (*t*-test, $p < 0.05$). The length–length relationships among the three length measurements (TL, FL, and SL) were highly correlated ($r^2 > 0.970$). The mean condition factor (*K*) was calculated as 0.812 ± 0.074 for both sexes, with no significant difference between females and males. These results will be useful for studies on the biology of non-indigenous fish species.*

Key words: The Malabar glassy perchlet, biological parameters, Levantine Sea, Türkiye

RELAZIONI LUNGHEZZA–PESO E LUNGHEZZA–LUNGHEZZA, E FATTORE DI CONDIZIONE DI *AMBASSIS DUSSUMIERI* CUVIER, 1828, NEL MEDITERRANEO NORD-ORIENTALE, TURCHIA

SINTESI

*Lo studio fornisce i primi dati sulla relazione lunghezza–peso (LWR), sulla relazione lunghezza–lunghezza (LLR) e sul fattore di condizione (*K*) del pesce vetro di Dussumier, *Ambassis dussumieri*, nel Mediterraneo nord-orientale (costa di Tuzla, Turchia). È stato esaminato un totale di 132 esemplari (68 maschi e 64 femmine). Gli esemplari presentavano una lunghezza totale (TL) compresa tra 4,5 e 8,9 cm, una lunghezza alla forcilla (FL) tra 3,7 e 7,7 cm e una lunghezza standard (SL) tra 3,4 e 6,9 cm per entrambi i sessi. I valori di *b* calcolati hanno indicato una crescita allometrica positiva per l'intera popolazione (*t*-test, $p < 0,05$). Le relazioni lunghezza–lunghezza tra le tre misure di lunghezza (TL, FL e SL) hanno mostrato un'elevata correlazione ($r^2 > 0,970$). Il fattore di condizione medio (*K*) è stato calcolato pari a $0,812 \pm 0,074$ per entrambi i sessi, senza differenze significative tra femmine e maschi. Questi risultati saranno utili per studi sulla biologia delle specie ittiche non indigene.*

Parole chiave: pesce vetro di Dussumier, parametri biologici, Mare del Levante, Turchia

INTRODUCTION

The Malabar glassy perchlet, *Ambassis dussumieri* Cuvier, 1828, is widely distributed in the Mediterranean Sea (Stern *et al.*, 2022) and throughout the Indo-West Pacific, from Algoa Bay in South Africa to the Red Sea and Seychelles, and eastwards to India, Australasia, the Philippines, and China (Anderson & Heemstra, 2003).

Ambassis dussumieri inhabits marine, estuarine, and riverine environments (Froese & Pauly, 2026). In the Mediterranean, the species was first recorded in the eastern basin, on the Israeli coast, by Stern *et al.* (2022) and subsequently reported by Çevik & Ergüden (2026) from Turkish marine waters in the northeastern Mediterranean. The pathway by which it entered the Mediterranean Sea remains uncertain, although Stern *et al.* (2022) suggested that the species likely migrated from the Red Sea to the Israeli coast via the Suez Canal.

The length–weight relationship (LWR) is an important tool in fish biology, ecology, and fisheries assessment, providing valuable information for stock assessment studies (Moutopoulos & Stergiou, 2002; Gonzalez Acosta *et al.*, 2004), particularly for converting length observations into weight estimates and assessing biomass (Gonzalez Acosta *et al.*, 2004).

The length–length relationship (LLR) is useful for standardising length types when summarising data and is also effective for comparative growth studies (Froese, 2006; Moutopoulos & Stergiou, 2002; Bakhshalizadeh *et al.*, 2025). Understanding relationships among total length, fork length, and standard length allows for the estimation of species distribution, assessment of population condition, and morphological comparisons. All these data contribute to informed decision-making and effective conservation strategies (Froese, 2006).

The present study aimed to investigate the length–frequency distribution, length–weight relationship (LWR), length–length relationship (LLR), and Fulton's condition factor (*K*) of the non-indigenous fish species of Malabar glassy perchlet *Ambassis dussumieri* in the northeastern Mediterranean Sea, Türkiye.

MATERIAL AND METHODS

A total of 132 specimens of *A. dussumieri* were collected in October, November, and December 2025 from the Kara Göçer Drainage Canal, Tuzla coast (Karataş, Adana), using a cast net (Fig. 1). Sampling was conducted approximately 400–500 m upstream from the canal mouth at a depth of




Fig. 1: Map showing the capture locality (•) of *Ambassis dussumieri* in the northeastern Mediterranean (Tuzla, Türkiye).

Sl. 1: Zemljevid obravnavanga območja z lokaliteto ulova primerkov vrste *Ambassis dussumieri* v severovzhodnem Sredozemlju (Tuzla, Turčija).

Tab. 1: Descriptive statistics and estimated parameters of the length–weight relationships of *Ambassis dussumieri* from the northeastern Mediterranean, Türkiye. *a*: = intercept, *b*: = slope, *SE.*: = Standard error, *C.I.*: = Confidence interval, *r*²: = Coefficient of determination, *GT*: = Growth type.

Tab. 1: Opisna statistika in ocenjeni parametri dolžinsko-masnih odnosov za vrsto *Ambassis dussumieri* iz severovzhodnega Sredozemlja, Turčija. *a*: = začetni koeficient (odsek), *b*: = naklon, *SE.*: = standardna napaka, *C.I.*: = interval zaupanja, *r*²: = koeficient determinacije, *GT*: = tip rasti.

Sample	No. of individuals	Total Length, TL (cm)	Weight, W (g)	Parameters of the relationship					
<i>Ambassis dussumieri</i>		$L_{min}-L_{max}$ (mean ±SD)	$W_{min}-W_{max}$ (mean ±SD)	a	b	SE(b)	95% CI of b	r ²	GT
Female	64	4.50–8.60 (6.86±0.949)	0.75–6.51 (2.83±1.348)	0.0036	3.417	0.068	3.282–3.553	0.976	A+
Male	68	4.50–8.90 (6.93±1.127)	0.72–6.99 (2.99±1.467)	0.0058	3.175	0.058	3.060–3.292	0.978	A+
Combined	132	4.50–8.90 (6.90±1.040)	0.72–6.99 (2.90±1.382)	0.0050	3.253	0.044	3.167–3.340	0.976	A+

about 1.5 m. After collection, the specimens were transported in a small iced Styrofoam box to the Basic Sciences Laboratory at the Faculty of Fisheries Science, Çukurova University.

Species identification was carried out based on the diagnostic criteria provided by Maugé (1986) and Anderson and Heemstra (2003). For each fish specimen, total length (TL), fork length (FL), and standard length (SL) were measured to the nearest 0.1 cm, and total weight (TW) was recorded to the nearest 0.01 g.

The sex was determined by macroscopic observation of the gonads in all individuals. Specimens of the Malabar glassy perchlet were grouped into size classes, and percentage frequency and TL were used to construct the length–frequency distribution. The sex ratio (males:females) was calculated and significant deviations from the expected ratio (1:1) were tested using the χ^2 test (Zar, 1996).

The functional relationship between length and weight of the specimens was fitted to the equation $W = aL^b$, where W is total weight (g), L is length (cm), and a and b are parameters to be estimated, with b representing the coefficient of allometry (Ricker, 1975). The parameters a and b were estimated using the least-squares regression on log-transformed length and weight values (Le Cren, 1951), according to the linearized form $\log W = \log a + b \log L$ (Sokal & Rohlf, 1987).

The b value is useful in describing fish growth type and associated changes in body shape under specific environmental conditions. When $b = 3$, growth is isometric; values of $b < 3$ indicate negative allometric growth, whereas values of $b > 3$ indicate positive allometric growth (Ricker, 1975; Sparre &

Venema, 1992; Avsar, 2016). Significant deviations of the estimated b value from the isometric value ($b = 3.0$) were tested using a t-test.

Fulton's condition factor was calculated from weight (g) and length (cm) and used to estimate changes in nutritional condition. The formula used was $K = 100 \times W / L^3$, where K is the condition factor, W is body weight, and L is total length (Le Cren, 1951; Pauly, 1980). Data were analysed using SPSS v22.0. All statistical tests were considered significant at the 0.05 level ($p < 0.05$).




Fig. 2: Total length frequencies of female and male *Ambassis dussumieri* from the northeastern Mediterranean coast of Türkiye.

Sl. 2: Frekvenčna porazdelitev celotne dolžine samice in samcev vrste *Ambassis dussumieri* s severovzhodne sredozemske obale Turčije.

RESULTS AND DISCUSSION

A total of 132 specimens were examined. The specimens ranged from 4.5 to 8.9 cm in total length (TL), 3.7 to 7.7 cm in fork length (FL), 3.4 to 6.9 cm in standard length (SL), and 0.72 to 6.99 g in total weight (TW) for the two sexes combined (Tab. 1). The maximum total length recorded during the study period was 8.9 cm. In males, TL ranged from 4.5 to 8.9 cm, and TW from 0.72 to 6.99 g, while in females, TL ranged from 4.5 to 8.6 cm and TW from 0.75 to 6.51 g.

The results show that the studied population consisted of 51.52% males ($n=68$) and 48.48% females ($n=64$), with maximum length and weight values higher in males. The overall sex ratio was M:F = 1.06:1, indicating a slight male bias (χ^2 test; $p < 0.05$). There was no significant difference between females and males regarding mean length (t-test, $t = 0.154$, $p > 0.05$) and weight (t-test, $t = 0.276$, $p > 0.05$).

The length–frequency relationship indicated that the 6.5 cm class size had the highest frequency, followed by the 8.0 cm class in both females and males, the 7.5 cm class in males, and the 5.5 cm class in females (Fig. 2).

The LWR indicated a positive allometric growth for both sexes (t-test, $p < 0.05$); the coefficient of determination (r^2) was 0.9762 for females, 0.9784 for males, and 0.9769 for the two sexes combined, indicating a strong correlation between length and weight. The length–weight equations were $W = 0.0036 \times L^{3.4171}$ for females, $W = 0.0058 \times L^{3.1757}$ for males, and $W = 0.0050 \times L^{3.2534}$ for the combined sexes (Fig. 3). The calculated parameters and length characteristics of the length–weight relationships are given in Table 1.

The relationship between total length (cm) and total weight (g) for all samples is shown in Fig. 3. All regression models were highly significant ($p < 0.05$), with coefficients of determination (r^2) exceeding 0.970 for all sexes.

In the length–length relationship analysis, the present study found that the b value of the LLRs was greater than 1, indicating positive allometric growth for females, males, and the two sexes combined. The relationships among the three length measurements (TL–FL–SL) were highly correlated ($r^2 > 0.970$, $p < 0.001$). Conversions among length measurements (TL, FL, and SL) are given in Table 2.

Fulton's condition factor (K) for females, males, and combined sexes was 0.807 ± 0.078 , 0.820 ± 0.072 , and 0.812 ± 0.074 , respectively, showing no significant variation ($p > 0.05$).

The maximum length values for *A. dussumieri* recorded in the present study are higher than those reported in previous studies (Stern *et al.*, 2022; Afrand & Sourinejad, 2023; Çevik & Ergüden, 2026;

Tab. 3). The enhanced growth and larger body size may be driven by nutrient-rich conditions associated with river inputs into the northeastern Mediterranean coastal waters of Türkiye, which likely boost primary production in the region.

Growth studies provide important scientific data for fisheries. The observed growth patterns showed positive allometry in the combined sample as well as the separate male and female samples, as the b value exceeded the isometric value ($b = 3.0$). The deviation




Fig. 3: Length–weight relationships for female, male, and combined of *Ambassis dussumieri* from the northeastern Mediterranean coast of Türkiye.

Sl. 3: Dolžinsko-masni odnosi za samice, samce in skupno populacijo vrste *Ambassis dussumieri* s severovzhodne sredozemske obale Turčije.

Tab. 2: Estimated parameters for the conversions between the length measurements (TL, FL, and SL in cm) of *Ambassis dussumieri* from the northeastern Mediterranean coast of Türkiye.

Tab. 2: Ocenjeni parametri za pretvorbe med merami dolžine (TL, FL in SL v cm) pri primerkih vrste *Ambassis dussumieri* s severovzhodne sredozemske obale Turčije.

Sample	No. of individuals	Equation	Constant a	Slope b	SE (b)	r ²
Female	64	TL = a + bSL	0.212	1.033	0.021	0.974
		SL = a + bFL	0.204	0.831	0.024	0.971
		FL = a + bTL	-0.379	1.106	0.016	0.971
Male	68	TL = a + bSL	0.215	1.023	0.018	0.980
		SL = a + bFL	0.141	0.866	0.015	0.982
		FL = a + bTL	-0.343	1.097	0.019	0.980
Combined	132	TL = a + bSL	0.236	1.013	0.014	0.977
		SL = a + bFL	0.149	0.862	0.024	0.971
		FL = a + bTL	-0.358	1.100	0.013	0.973

of the *b* value for the combined sexes (3.253) was statistically significant ($p < 0.05$).

Bagenal & Tesch (1978) stated that *b* values can vary primarily depending on sample size, length interval, study season, sea conditions, and environmental conditions at the sampling location. Our results are not fully comparable with previous records, since

not all of them included LWR studies for this species. However, a comparison between the previously reported length–weight parameters and those obtained in our study is provided in Table 3.

Additional factors are known to influence the length–weight parameters in fish, including gonad maturity, stomach fullness, health, and preservation techniques (Bagenal & Tesch, 1978; Wootton, 1998). Our length and weight data were collected during autumn and winter, however, the season and other above-mentioned factors were not considered in the study.

Fulton's condition factor (*K*), an expression of the relative fatness of fish, which indicates the interaction between biotic and abiotic factors and general fish condition, was 0.812 for the sexes combined. This factor is suitable for comparing the condition of fish of the same species across seasons, locations, or sexes, and reflects ecological and physiological conditions, providing information on the fish's physiological state. This metric also relates to fish welfare in terms of reproduction and nutrition (Le Cren, 1951).

In our study, the condition value was less than 1 ($K < 1$), possibly reflecting food/prey scarcity, increased competition, or environmental factors, including water quality. Nevertheless, our findings indicate that the population of *A. dussumieri* primarily exhibited allometric growth (+), which suggests that the northeastern Mediterranean waters of Türkiye still provide, on average, a suitable feeding environment for this species.

Prior to this study, no information was available on the length–weight and length–length relationships or condition factor of *A. dussumieri* along the Mediterranean coasts, and no biological data on this species were available in the FishBase database. Therefore, this study provides the first comprehensive information on the species in Mediterranean waters.

Tab. 3: Geographic comparison of maximum length and length–weight relationship data for *Ambassis dussumieri*.

Tab. 3: Geografska primerjava največje dolžine in podatkov o dolžinsko-masnem odnosu za vrsto *Ambassis dussumieri*.

Study	Locality	Country	Samples	No. of individuals	Length Type	$L_{min}-L_{max}$ (cm)	$W_{min}-W_{max}$ (g)
Stern <i>et al.</i> (2022)	Eastern Mediterranean Sea	Israel	Combined	6	TL	3.94–4.88	0.51–0.97
Afrand & Sourinejad (2023)	Persian Gulf, Khuran Strait	Iran	Combined	2	TL	4.98–5.20	-
Çevik & Ergüden (2026)	Northeastern Mediterranean Sea	Türkiye	Combined	4	TL	7.17–8.65	3.46–6.67
This study	Tuzla coast, northeastern Mediterranean Sea	Türkiye	Combined	132	TL	4.50–8.90	0.72–6.99

It also contributes to scientific knowledge of the Mediterranean Sea by documenting a newly recorded fish species and supporting the inclusion of its biological parameters in the FishBase database.

ACKNOWLEDGMENTS

The authors thank the Gözlem I captain and crew for their assistance in collecting the specimens.

DOLŽINSKO-MASNI ODNOS MED DOLŽINO IN KONDICIJSKIM FAKTORJEM
PRI VRSTI *AMBASSIS DUSSUMIERI* CUVIER, 1828,
V SEVEROVZHODNEM SREDOZEMSKEM MORJU,
TURČIJA

Deniz ERGÜDEN

Marine Science Department, Faculty of Marine Science and Technology, Iskenderun Technical University, 31200 Iskenderun, Hatay, Türkiye
e-mail: deniz.erguden@iste.edu.tr; derguden@gmail.com

Cem ÇEVİK

Faculty of Fisheries, Çukurova University Balcalı Campus, 01330, Sarıçam/Adana, Türkiye

POVZETEK

Avtorja obravnavata prve podatke o dolžinsko-masnem odnosu (LWR), razmerju med dolžinami (LLR) in kondicijskem faktorju (K) za Malabarskega steklenega ostriža *Ambassis dussumieri* v severovzhodnem Sredozemskem morju (obala Tuzle, Turčija). Pregledala sta skupaj 132 osebkov (68 samcev in 64 samic). Primerki obeh spolov so merili od 4,5 do 8,9 cm celotne dolžine (TL), od 3,7 do 7,7 cm viličaste dolžine (FL) in od 3,4 do 6,9 cm standardne dolžine (SL). Izračunane vrednosti b so nakazovale pozitivno alometrično rast za celotno populacijo (t-test, $p < 0,05$). Odnos med tremi merami dolžine (TL, FL in SL) je bil statistično zelo značilen ($r^2 > 0,970$). Povprečni koeficient kondicije (K) je bil $0,812 \pm 0,074$ za oba spola, brez pomembnih razlik med samicami in samci. Ti rezultati bodo uporabni za raziskave o biologiji tujerodnih vrst rib.

Ključne besede: Malabarski stekleni ostriž, biološki parametri, Levantsko morje, Turčija

REFERENCES

- Afrand, M. & I. Sourinejad (2023):** Morphological identification and DNA barcoding confirm the discovery of *Ambassis dussumieri* (Ambassidae) in the Persian Gulf. *J. Ichthyol.*, 63, 1072–1079.
- Anderson, M.E. & P.C. Heemstra, (2003):** Review of the glassfishes (Perciformes: Ambassidae) of the Western Indian Ocean. *Cybium*, 27(3), 199–209.
- Avsar, D. (2016):** Fisheries Biology and Population Dynamics (in Turkish). Akademisyen Publishing, Ankara, Turkey, 148 pp.
- Bakhshalizadeh, S., K. Abbasi, A.R. Liafuie & F. Tiralongo (2025):** Evidence for reducing fitness and competition of the native fish with invasive fish species in the south Caspian Sea. *Mar. Environ. Res.*, 212, 107566.
- Bagenal, T.B. & F.W. Tesch (1978):** Age and growth. In: Bagenal T. (ed.): *Methods for assessment of fish production in fresh waters*, 3rd ed. IBP Handbook No. 3. Blackwell Science Publications, Oxford, pp. 101–136.
- Çevik, C. & D. Ergüden (2026):** First record of *Ambassis dussumieri* Cuvier, 1828 (Osteichthyes: Ambassidae), in the North-Eastern Mediterranean, Türkiye. *Zool. Mid. East.*, 72, in press. <https://doi.org/10.1080/09397140.2026.2684418>.
- Froese, R. (2006):** Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *J. Appl. Ichthyol.*, 22(4), 241–253.
- Froese, R.D. & D. Pauly (2026):** Fishbase. World Wide Web Electronic Publication. www.fishbase.org. version (11/2025) (accessed: 25 February 2026).
- González Acosta, A.F., G. De La Cruz Agüero & J. De La Cruz Agüero (2004):** Length-weight relationships of fish species caught in a mangrove swamp in the Gulf of California (Mexico). *J. Appl. Ichthyol.*, 20(2), 154–155.
- Le Cren, E.D. (1951):** The length-weight relationship and seasonal cycle in the gonad weight and condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.*, 20, 201–219.
- Maugé, L.A. (1986):** Ambassidae. In: Daget, J. & J. P. Gosse & D. F. E. Thys van den Audenaerde (eds.): *Check-list of the freshwater fishes of Africa (CLOFFA)*. Vol. 2. ISNB, Brussels; MRAC, Tervuren; and ORSTOM, Paris, pp. 297–298.
- Moutopoulos, D.K. & K.I. Stergiou (2002):** Length-weight and length-length relationships of fish species from the Aegean Sea (Greece). *J. Appl. Ichthyol.*, 18(3), 200–203.
- Pauly, D. (1980):** Some simple methods for the assessment of tropical fish stocks. *FAO Fisheries Technical Paper*, 51, FAO, Rome, 234 pp.
- Ricker, W.E. (1975):** Computation and interpretation of biological statistics of fish populations. *Bulletin of the Fisheries Research Board of Canada*. *Bulletin* No. 191, Ottawa, Canada, 382 pp.
- Sokal, R. & F. Rohlf (1987):** *Introduction to Biostatistics*. Freeman, New York, USA, 363 pp.
- Sparre, P. & S.C. Venema (1992):** *Introduction to tropical fish stock assessment: Part I. Manual* FAO Fisheries Technical Paper. No. 306. 1. FAO, Rome, Italy, 337 pp.
- Stern, N., K. Gayer & A. Morov (2022):** A transparent invasion: a first Mediterranean record and an established population of the glassfish *Ambassis dussumieri* Cuvier 1828. *Medit. Mar. Sci.*, 23(1), 191–195.
- Wootton, R.J. (1998):** *Ecology of teleost fishes*. Kluwer Academic Publishers. 2nd Edition, Dordrecht, Boston, 386 pp.
- Zar, J.H. (1996):** *Biostatistical Analysis*. 3rd Ed. Prentice-Hall, Englewood Cliffs, NJ, 662 pp.