

ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterranee
Annals for Istrian and Mediterranean Studies
Series Historia Naturalis, 36, 2026, 1





ANNALES

Anali za istrske in mediteranske študije
Annali di Studi istriani e mediterraneei
Annals for Istrian and Mediterranean Studies

Series Historia Naturalis, 36, 2026, 1

**UREDNIŠKI ODBOR/
COMITATO DI REDAZIONE/
BOARD OF EDITORS:**

Alessandro Acquavita (IT), Nicola Bettoso (IT), Christian Capape (FR), Darko Darovec, Dušan Devetak, Jakov Dulčić (HR), Edy Fantinato (IT), Ana Fortič, Andrej Gogala, Daniel Golani (IL), Danijel Ivajnskič, Hakan Kabasakal (TR), Mitja Kaligarič, Marcelo Kovačič (HR), Petar Kružić (HR), Lovrenc Lipej, Vesna Mačić (ME), Alenka Malej, Borut Mavrič, Patricija Mozetič, Martina Orlando-Bonaca, James Pierson (USA), Michael Stachowitsch (AT), Francesco Tiralongo (IT), Tom Turk, Al Vrezec

**Glavni urednik/Redattore capo/
Editor in chief:**

Darko Darovec

**Odgovorni urednik naravoslovja/
Redattore responsabile per le scienze
naturali/Natural Science Editor:**

Lovrenc Lipej

Urednici/Redattrici/Editors:

Martina Orlando-Bonaca, Ana Fortič

Prevajalci/Traduttori/Translators:

Martina Orlando-Bonaca (sl./it.)

**Oblikovalec/Progetto grafico/
Graphic design:**

Dušan Podgornik, Lovrenc Lipej

Tisk/Stampa/Print:

Založništvo PADRE d.o.o.

Izdajatelj/Editori/Published by:

Zgodovinsko društvo za južno Primorsko - Koper / *Società storica del Litorale - Capodistria*[®]
Inštitut IRRIS za raziskave, razvoj in strategije družbe, kulture in okolja / *Institute IRRIS for Research, Development and Strategies of Society, Culture and Environment* / *Istituto IRRIS di ricerca, sviluppo e strategie della società, cultura e ambiente*[®]

**Sedež uredništva/Sede della redazione/
Address of Editorial Board:**

Nacionalni inštitut za biologijo, Morska biološka postaja Piran / *Istituto nazionale di biologia, Stazione di biologia marina di Pirano* / *National Institute of Biology, Marine Biology Station Piran*
SI-6330 Piran / *Pirano*, Fornače/Fornace 41, tel.: +386 5 671 2900, fax +386 5 671 2901;
e-mail: annalesSHN@nib.si, internet: www.zdjp.si

Redakcija te številke je bila zaključena 20. 06. 2026.

**Sofinancirajo/Supporto finanziario/
Financially supported by:**

Javna agencija za znanstveno-raziskovalno in inovacijsko dejavnost Republike Slovenije (ARIS)

Annales - Series Historia Naturalis izhaja dvakrat letno.**Naklada/Tiratura/Circulation:** 300 izvodov/copie/copies

Revija Annales, Series Historia Naturalis je vključena v naslednje podatkovne baze / *La rivista Annales, series Historia Naturalis è inserita nei seguenti data base* / *Articles appearing in this journal are abstracted and indexed in:* BIOSIS-Zoological Record (UK); Aquatic Sciences and Fisheries Abstracts (ASFA); Elsevier B.V.: SCOPUS (NL); Directory of Open Access Journals (DOAJ).

To delo je objavljeno pod licenco / *Quest'opera è distribuita con Licenza* / *This work is licensed under a Creative Commons BY 4.0.*



Navodila avtorjem in vse znanstvene revije in članki so brezplačno dostopni na spletni strani <https://zdjp.si/en/p/annalesshn/>
The submission guidelines and all scientific journals and articles are available free of charge on the website <https://zdjp.si/en/p/annalesshn/>
Le norme redazionali e tutti le riviste scientifiche e gli articoli sono disponibili gratuitamente sul sito <https://zdjp.si/en/p/annalesshn/>



VSEBINA / *INDICE GENERALE* / *CONTENTS*

BIOTSKA GLOBALIZACIJA
GLOBALIZZAZIONE BIOTICA
BIOTIC GLOBALIZATION

Alen SOLDI, Rigers BAKIU & Sherif DURMISHAJ
The First Record of Bastard Grunt *Pomadasys incisus* (Bowdich, 1825) in Albanian Waters (Adriatic Sea) 1
Prvi zapis o pojavljanju neprave prašičevke Pomadasys incisus (Bowdich, 1825) v albanskih vodah (Jadransko morje)

Pero UGARKOVIĆ, Ilija ČETKOVIĆ, Olivera MARKOVIĆ, Aleksandar JOKSIMOVIĆ, Nikola ĐORĐEVIĆ & Jakov DULČIĆ Additional Records of the Bastard Grunt, *Pomadasys incisus* (Bowdich, 1825), from the Eastern Adriatic Sea 7
Dodatne najdbe neprave prašičevke, Pomadasys incisus (Bowdich, 1825), iz vzhodnega Jadranskega morja

Rüştü KIRMAN & Murat BİLECENOĞLU Significant Range Expansion of *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) in the Aegean Sea Based on Scuba and Photographic Observations 15
Znatno razširjanje areala vrste Sepioteuthis lessoniana (Cephalopoda: Loliginidae), v Egejskem morju na podlagi podvodnih opazovanj in fotografiranja

Adrian BRAJKOVIĆ, Iris MATULJA & Neven IVEŠA
Conquering the North: New Occurrence of the Common Lionfish, *Pterois miles* (Bennett, 1828), in the Adriatic Sea 21
Osvajanje severa: novi zapis o pojavljanju navadne plamenke, Pterois miles (Bennett, 1828), v Jadranskem morju

Alan DEIDUN, Bruno ZAVA, Maria CORSINI-FOKA, Arnold SCIBERRAS & Alessio MARRONE First Record of the Devil Firefish *Pterois miles* (Bennett, 1828) (Actinopterygii: Scorpaenidae) from Maltese Coastal Waters 29
Prvi zapis o pojavljanju plamenke Pterois miles (Bennett, 1828) (Actinopterygii: Scorpaenidae) v malteških obalnih vodah

Chirine HUSSEIN, Amir IBRAHIM, Firas ALSHAWY, Mouina BADRAN & Rahaf ABO ASA First Mediterranean Record of the Diagonal Butterflyfish, *Chaetodon fasciatus* Forsskal, 1775, Reported from Syrian Waters 35
Prvi sredozemski zapis o pojavljanju poševnoprogame ščetinozobke, Chaetodon fasciatus Forsskal, 1775, iz sirskih voda

Houssein ELBARAASI, Tarek SHOEIB, Mona SAID & Laith A. JAWAD A Further Record of the Barred Knifejaw, *Oplegnathus fasciatus* (Temminck & Schlegel, 1844), a Pacific Species, from the Mediterranean: A New Record from Benghazi, Libya 41
Novi zapis o vrsti Oplegnathus fasciatus (Temminck & Schlegel, 1844), pacifiški vrsti, v Sredozemlju: nova najdba iz Bengazija v Libiji

Alen SOLDI & Rigers BAKIU Rapid Increase in Records of the Invasive Silver-cheeked Toadfish *Lagocephalus sceleratus* (Gmelin, 1789) in the Adriatic Sea 49
Hitro naraščanje pojavov invazivne srebrnoproge napihvalke Lagocephalus sceleratus (Gmelin, 1789) v Jadranskem morju

Gerasimos KONDYLATOS, Konstantinos KALAENTZIS, Styliani MINOUDI & Maria CORSINI-FOKA Further Molecular Identification Confirms the Occurrence of *Lagocephalus guentheri* Miranda Ribeiro, 1915 in the Aegean Coastal Waters of Greece 55
Nadaljnja molekularna identifikacija potrjuje prisotnost vrste Lagocephalus guentheri Miranda Ribeiro, 1915 v egejskih obalnih vodah Grčije

Deniz ERGÜDEN & Cem ÇEVİK Length–Weight and Length–Length Relationships, and Condition Factor of *Ambassis dussumieri* Cuvier, 1828, in the Northeastern Mediterranean, Türkiye 65
Dolžinsko-masni odnos med dolžino in kondicijskim faktorjem pri vrsti Ambassis dussumieri Cuvier, 1828, v severovzhodnem Sredozemskem morju, Turčija

Okan AKYOL & HALİL ŞEN Unexpected Occurrence of Prussian Carp *Carassius gibelio* (Cyprinidae) in Homa Lagoon (Izmir Bay, Aegean Sea) 73
Nepričakovani pojav srebrnega koreslja Carassius gibelio (Cyprinidae) v laguni Homa (Izmirski zaliv, Egejsko morje)

SREDOZEMSKE HRUSTANČNICE
SQUALI E RAZZE MEDITERRANEE
MEDITERRANEAN SHARKS AND RAYS

Terry CARBON, Emily GIGNON, Justine LALLAU-VAZZOLER, Hugo MENARD, Claudio BARRIA, Ana I. COLMENERO & Nicolas ZIANI Precopulatory Behaviour of *Pteroplatytrygon violacea* (Myliobatiformes: Dasyatidae) in the Northwestern Mediterranean 81
Predkopulacijsko vedenje vijoličnega morskega biča (Pteroplatytrygon violacea) (Myliobatiformes: Dasyatidae) v severozahodnem Sredozemlju

Hakan KABASAKAL Distribution of <i>Odontaspis ferox</i> in the Mediterranean Sea: Insights from Spatial and Temporal Analyses 87 <i>Razširjenost vrste Odontaspis ferox v Sredozemskem morju: Spoznanja na podlagi prostorskih in časovnih analiz</i>	
Hristina GELEVSKA, Borut MAVRIČ, Lovrenc LIPEJ & Christian CAPAPÉ Is the Gulf of Trieste a potential nursery area for some elasmobranch species? 99 <i>Ali je Tržaški zaliv potencialno območje odraščanja (jaslice) za določene vrste hrustančnic?</i>	
IHTIOFAVNA ITTIOFAUNA ICHTHYOFAUNA	
Cem DALYAN, Yunus GÖNÜL, Mahmud Samed ŞAHİNOĞULLARI & Hakan KABASAKAL New data on the Occurrence and Morphology of the Armless Snake Eel, <i>Dalophis imberbis</i> (Ophichthidae), from the Northeastern Mediterranean Sea 119 <i>Novi podatki o pojavljanju in morfologiji kačaste jegulje, Dalophis imberbis (Ophichthidae), iz severovzhodnega Sredozemskega morja</i>	
Chirine HUSSEIN, Firas ALSHAWY & Amir IBRAHIM First Record of the Mediterranean Dealfish, <i>Trachipterus trachipterus</i> (Gmelin, 1789) (Trachipteridae), in Syrian Marine Waters 127 <i>Prvi zapis o pojavljanju kosice, Trachipterus trachipterus (Gmelin, 1789) (Trachipteridae), v sirskih morskih vodah</i>	
Igor AGOSTINI & Okan AKYOL On the Occurrence of the Currently Largest Recorded <i>Pagellus erythrinus</i> (Sparidae) in the Tyrrhenian Sea (Italy) 133 <i>O pojavljanju trenutno največjega evidentiranega primerka vrste Pagellus erythrinus (Sparidae) v Tirenskem morju (Italija)</i>	
Ismail Burak DABAN & Yusuf ŞEN Substantiated Record and Preliminary Biological Insights of <i>Microlophrys dalmatinus</i> (Steindachner & Kolombatović, 1883) from the Sea of Marmara, Türkiye 141 <i>Potrjen zapis o pojavljanju in predhodna biološka spoznanja o vrsti Microlophrys dalmatinus (Steindachner & Kolombatović, 1883) iz Marmarskega morja, Turčija</i>	
Zeliha ERDOĞAN, Gülçin ULUNEHİR AYDIN, Hatice TORCU-KOÇ & Tuğba ANBAROĞLU Population Study of <i>Trachurus mediterraneus</i> Focused on Reproductive Biology in Edremit Bay, Northern Aegean Sea 151 <i>Razmnoževalna biologija sredozemskega šura (Trachurus mediterraneus) v sklopu populacijske raziskave v Edremitnem zalivu (severno Egejsko morje)</i>	
MORSKA FAVNA FAUNA MARINA MARINE FAUNA	
Francesco TIRALONGO, Paola LEOTTA & Riccardo MARTELLUCCI <i>Physalia physalis</i> in the central Mediterranean Sea: Recent observations associated with mass strandings of <i>Velella velella</i> and surface circulation dynamics 165 <i>Portugalska ladjica (Physalia physalis) v osrednjem Sredozemskem morju: nedavna opažanja v povezavi z masovnimi nasedanji morskega jadrčka (Velella velella) in dinamiko površinskega kroženja vode</i>	
Andrea LOMBARDO, Juba CHABANE & Rachida GHALMI First Contribution to the Study of the "Sea Slug" Fauna (Gastropoda, Heterobranchia) of the Algerian coast 177 <i>Prvi prispevek k proučevanju favne „goliš morskih polžev“ (Gastropoda, Heterobranchia) alžirske obale</i>	
Borut MAVRIČ, Tjaša PREMRL, Martin MAVRIČ, Tina MIRT, Neža LEBAN, Tihomir MAKOVEC & Lovrenc LIPEJ Assessing Soft-Bottom Epibenthic Communities: Methodological Insights into Dredging and Video Surveys 203 <i>Ocenjevanje epibentoških združb mehkega dna: metodološki vpogledi v dredžanje in video pregleda</i>	
FAVNA FAUNA FAUNA	
Kim LEBAN, Iztok ŠKORNIK, Špela ČONČ, Mateja BREG VALJAVEC, Žan KURALT, Lenart ŠTAUT, Jure TIČAR, Katarina POLAJNAR HORVAT, Primož GAŠPERIČ, Aleš SMREKAR & Melita VAMBERGER Habitat Insights for the European Pond Turtle (<i>Emys orbicularis</i>) in the Coastal Wetland Area Sečovlje Salina Nature Park, Slovenia 221 <i>Habitatne značilnosti močvirske sklednice (Emys orbicularis) v krajinskem parku Sečoveljske soline, Slovenija</i>	
Leon CIMERMAN & Matija KRIŽNAR Najdbe pleistocenske avifavne v kamnolomu Črni Kal (Primorska, Slovenija) 243 <i>Pleistocene Avifauna Finds in the Črni Kal Quarry (Primorska Region, Slovenia)</i>	
Kazalo k slikam na ovitku 255 Index to images on the cover 255	

IS THE GULF OF TRIESTE A POTENTIAL NURSERY AREA FOR CERTAIN ELASMOBRANCH SPECIES?

Hristina GELEVSKA

FAMNIT, University of Primorska, Koper, Slovenia
e-mail: hgelevska@gmail.com

Borut MAVRIČ & Lovrenc LIPEJ

Marine Biology Station Piran, National Institute of Biology, Piran, Slovenia

Christian CAPAPÉ

Laboratoire d'Ichtyologie, Université de Montpellier, 34 095 Montpellier cedex 5, France

ABSTRACT

This study aimed to determine whether the Gulf of Trieste meets the criteria for a nursery area as defined by different sources, primarily Heupel et al. (2007). The analysis was based on elasmobranch specimens from the collection of the Marine Biology Station Piran (National Institute of Biology, Slovenia), comprising five species (Mustelus mustelus, M. punctulatus, Torpedo marmorata, Raja asterias, and Aetomylaeus bovinus) caught as bycatch over a 24-year period. Four of the five species (excluding A. bovinus) fulfilled all three criteria. According to the data analyzed, the Gulf of Trieste could be considered a communal juvenile nursery area for the assessed species. In addition, records of juveniles belonging to other elasmobranch species reported during the study period in the area suggest the potential importance of this area as a nursery ground for other species as well.

Key words: elasmobranchs, parturition, nursery area criteria, abiotic conditions, bycatch, northern Adriatic

PUÒ IL GOLFO DI TRIESTE VENIR CONSIDERATO UN'AREA DI NURSERY PER ALCUNE SPECIE DI ELASMOBRANCHI?

SINTESI

Lo studio ha l'obiettivo di verificare se il Golfo di Trieste soddisfa i criteri per essere definito un'area di nursery secondo diverse fonti, in particolare secondo Heupel et al. (2007). L'analisi si è basata sugli esemplari di elasmobranchi conservati nella collezione della Stazione di biologia marina di Pirano (Istituto nazionale di biologia, Slovenia), comprendente cinque specie (Mustelus mustelus, M. punctulatus, Torpedo marmorata, Raja asterias e Aetomylaeus bovinus) catturate come catture accessorie nell'arco di 24 anni. Quattro delle cinque specie (esclusa A. bovinus) hanno soddisfatto tutti e tre i criteri. Sulla base dei dati analizzati, il Golfo di Trieste potrebbe essere considerato un'area di nursery giovanile comune per le specie valutate. Le segnalazioni di individui giovanili di altre specie di elasmobranchi registrate nell'area suggeriscono l'importanza potenziale del Golfo come nursery anche per altre specie.

Parole chiave: elasmobranchi, parto, criteri di area di nursery, condizioni abiotiche, catture accessorie, Adriatico settentrionale

INTRODUCTION

Most elasmobranch species occupy high trophic levels in marine ecosystems and play a major role in their structure and function, which means that their removal would likely be detrimental to these ecosystems (Pough & Janis, 2019). They display K-selected life-history traits, which include slow growth rates, long lifespans, late maturity, long gestation periods, and few offspring (Dulvy et al., 2014). Along with a general tendency toward age and sex segregation, these traits make them especially vulnerable to human impacts, as many species require more than one suitable habitat for different segments of the population (i.e., juveniles and adults, males and females) (Muñoz-Chapuli, 1984; Bradai et al., 2005; Pough & Janis, 2019). In the Mediterranean Sea, volumes and species composition of elasmobranchs captured as bycatch are poorly documented, with available data often too inconsistent to be incorporated in national and international statistics (Dulvy et al., 2014) and the taxonomic resolution of catch statistics remaining poor (Cashion et al., 2019). Recently, the Important Shark and Ray Area (ISRA) identification process in the Mediterranean Sea has classified 2 ISRAs in the northern Adriatic, namely, the western Vir Sea (Croatia), important for angel sharks (*Squatina squatina*), and Cervia–Marina di Ravenna (Italy), important for sandbar sharks (*Carcharhinus plumbeus*) (ISRA, 2025). Additionally, the entire northwestern Adriatic has been classified as an ISRA for threatened species such as the spiny dogfish (*Squalus acanthias*), and as a reproductive area for the common smooth-hound (*Mustelus mustelus*) (ISRA, 2025). The Gulf of Trieste and the broader northern Adriatic have previously

been suggested as nursery areas for certain elasmobranch species by Lipej et al. (2008; 2024; Fig. 1), but they have never been directly evaluated as such.

Beck et al. (2009) define a nursery area as an environment in which juveniles of a certain species occur at higher densities, experience lower predation risk, or exhibit faster growth rates than in other areas (Tab. 2). Also, the species using an area as a nursery are expected to enter the area as neonates, accumulate biomass, and then move offshore. However, some of these criteria are vague and difficult to assess, even with modern technology (Heupel et al., 2007). Heupel et al. (2007) proposed an alternative set of criteria, according to which an area can be defined as an elasmobranch nursery if: i) newborn or young-of-the-year individuals are encountered there more frequently than in other areas; ii) newborn or young-of-the-year individuals tend to remain in or return to for extended periods; iii) the area or habitat is repeatedly used across years (Tab. 2). Martins et al. (2018) adopted the criteria proposed by Heupel et al. (2007) for viviparous species, but added criteria for discriminating between nursery areas of ovoviviparous and oviparous species, and distinguished juvenile habitats (secondary nursery areas) from nursery habitats (primary nursery areas), as first introduced by Bass (1978). Primary and secondary nursery areas may overlap, but not necessarily. Distinguishing between them is important in areas where many species co-occur, given that cannibalism is common among some elasmobranch groups. In these cases, juvenile aggregation areas that are spatially separated from the adults may in fact function as nursery areas, with the adult–juvenile segregation serving as an important qualifying



Fig. 1: Juvenile and newborn specimens of the Mediterranean starry ray (*Raja asterias*) and adult and newborn specimens of the marbled electric ray (*Torpedo marmorata*) from the Gulf of Trieste. (Photos by Jernej Uhan and Domen Trkov, respectively)

Sl. 1: Mladostni osebki in novorojenčki zvezdaste raže (*Raja asterias*) ter odrasli osebki in novorojenčki marmoriranega električnega skata (*Torpedo marmorata*) iz Tržaškega zaliva. (Avtorja fotografij: J. Uhan in D. Trkov).

characteristic (Heupel *et al.*, 2007; Martins *et al.*, 2018). There have also been records of different species using the same area as a nursery, called a communal nursery, where juveniles of multiple species co-occur while adults are mostly absent (Rosa *et al.*, 2023). However, in such nurseries, juveniles face a trade-off between lower predation risk and increased competition, both with conspecifics and juveniles of other species. This effect may be reduced through resource partitioning, provided these are sufficient within the nursery area (Rosa *et al.*, 2023). The purpose of this study was to examine the available data on elasmobranch species in the Gulf of Trieste and determine whether sufficient evidence exists to classify the area as an elasmobranch nursery.

MATERIAL AND METHODS

Study area

The Gulf of Trieste is the northernmost part of the Adriatic Sea, shared by Italy, Slovenia, and Croatia (Fig. 2). The Istrian peninsula partially closes it

off to the south, creating a relatively isolated area (Malačič & Petelin, 2009). Covering around 600 km², the gulf has an average depth of 18.7 m and a maximum depth of 25 m (DEIMS-SDR, 2025). Apart from several flat islets at the entrance to the Marano–Grado lagoon, no islands occur within the gulf. Its main freshwater input (Fig. 2) is provided by the Isonzo River (Monti *et al.*, 2012). Although tidal amplitudes in the Gulf of Trieste are among the largest in the Adriatic Sea, they rarely exceed 1 m (Malačič & Petelin, 2009). The southeastern coast, extending from Miramare to Muggia, has been extensively modified by human activity and artificial structures such as piers and harbor facilities (Monti *et al.*, 2012). The majority of the Slovenian part of the gulf is characterized by sedimentary bottom (mud, muddy sand, and sand) (UNEP, 2021). The remaining rocky coastal areas include sandstone habitat types colonized by algal vegetation, whereas sheltered bays support seagrass meadows. Offshore, the soft-bottom seabed is interspersed with thousands of biogenic outcrops formed by calcareous organisms growing on various hard substrates (Falace *et al.*, 2024). The northern Adriatic is one of the



Fig. 2: Map of the Gulf of Trieste.

Sl. 2: Zemljevid obravnavanega območja Tržaškega zaliva.

regions with the highest permanent production in the Mediterranean Sea, where mesozooplankton biomass exhibits a much broader range than elsewhere (Conversi *et al.*, 2009).

Species analyzed

A total of 549 specimens belonging to 5 species were assessed, including 2 shark species and 3 batoid species (Tab. 1): the common smooth-hound (*M. mustelus*), the blackspotted smooth-hound (*M. punctulatus*), the marbled electric ray (*Torpedo marmorata*), the Mediterranean starry ray (*Raja asterias*), and the bull ray (*Aetomylaeus bovinus*). All specimens included in the analysis were caught as bycatch in the Gulf of Trieste between 2000 and 2024 and were subsequently handed over to researchers at the Marine Biology Station (National Institute of Biology) in Piran. Specimens were sexed based on the presence or absence of secondary sexual characters (*i.e.*, claspers). Sexual maturity and age were determined using indicators such as clasper calcification, umbilical scars, and body size compared to known size at maturity. For specimens without direct age/maturity data but for which biometric data were available, maturity was estimated as newborn, juvenile, or adult based on publicly available literature (CIESM, 2025; FishBase, 2025). For each specimen, we calculated Fulton's condition factor using the formula (Fulton, 1902): $K = (W/L^3) \times 100$, where K is Fulton's condition factor, W is the weight of the fish (g), and L is the length (typically total length in cm). We then compared these values to species-specific condition factors from previous studies conducted in the Mediterranean Sea. Condition factors serve as indicators of fish welfare and population health, as they reflect environmental quality and suitability, including water quality and food availability.

They also provide information on the interaction between biotic and abiotic factors influencing the physiological condition of different species (Ragheb, 2023).

The occurrence of the species in the Adriatic Sea was also included in the analysis. Five occurrence classes were defined by Soldo & Lipej (2022): rare – when species is recorded after a long period of time (decades); occasional – recorded every few years; common – with several single records on a yearly basis; abundant – multiple records in catches or sighted every year; questionable/not confirmed – when a record needs confirmation. All five species included in this study have been reported to favor shallow coastal waters along the continental shelf, characterized by high food availability, as reproductive and nursery habitats (Serena & Relini, 2006; Consalvo *et al.*, 2007; Enajjar *et al.*, 2015; Catalano *et al.*, 2021); therefore, the Gulf of Trieste was considered a strong candidate for a potential nursery area.

Criteria for nursery areas

Among several nursery area definitions, we decided to use the criteria defined by Heupel *et al.* (2007) (Tab. 2), as they can be applied to areas such as the northern Adriatic, where bycatch and sighting reports are scant and tracking data for juveniles or adults are unavailable. The biotic and abiotic conditions of the Gulf of Trieste were also compared to the environmental preferences of each of the five species through a systematic literature review. Analyzing these parameters alongside comparable data from the rest of the northern Adriatic and the Mediterranean Sea, we assessed whether the Gulf of Trieste can be considered a nursery ground for certain elasmobranch species found in the Adriatic Sea, based on the three nursery area criteria described by Heupel *et al.* (2007).

Tab. 1: Species analyzed, occurrence status in the Adriatic, total number of specimens caught, number of specimens with available biometric data (males:females), and total length (TL) at maturity for each sex.

Tab. 1: Analizirane vrste, njihov status v Jadranu, število ujetih osebkov, biometrični podatki (samci: samice) in celotna dolžina (TL) ob spolni zrelosti za vsak spol.

Latin name	Common name	Occurrence in the Adriatic Sea (Soldo & Lipej, 2022)	No. of specimens caught (2000-2024)	No. of male : female specimens with biometric data	Total length at maturity for males : females in cm	Reproductive Period / parturition
<i>Mustelus mustelus</i>	common smooth-hound	abundant	138	13 : 20	70–80 cm : 70–80 (Lipej <i>et al.</i> , 2011; FishBase 2025)	late spring (Saïdi <i>et al.</i> , 2008; Özcan & Başusta, 2018; Boscolo Palo <i>et al.</i> , 2022)
<i>Mustelus punctulatus</i>	blackspotted smooth-hound	common	175	111 : 56	90 cm : 100 (Lipej <i>et al.</i> , 2011; FishBase, 2025)	mid-May to early June (Saïdi <i>et al.</i> , 2009; Boscolo Palo <i>et al.</i> , 2022)
<i>Torpedo marmorata</i>	marbled electric ray	common	74	47 : 27	24.1 : 26.1 (Bajt <i>et al.</i> , 2024)	May, August, September–February (Capapé, 1979; Consalvo <i>et al.</i> , 2007; Chatzispayrou <i>et al.</i> , 2021)
<i>Raja asterias</i>	Mediterranean starry ray	common	127	88 : 22	45–54 : 56.1–60 (Sviben <i>et al.</i> , 2019; CIESM, 2025)	summer to early autumn (Stehmann & Bürkel, 1984; Serena & Relini, 2006)
<i>Aetomylaeus bovinus</i>	bull ray	rare	37	20 : 17	80–95 : 90–100 (CIESM, 2025)	late spring and summer (Michael, 1993)

Tab. 2: Summary of nursery area definitions and criteria according to different authors.**Tab. 2: Pregled definicij in meril za območja odraščanja (ang. nursery areas) po različnih avtorjih.**

Publication	Nursery Area Criteria
Heupel et al., 2007	(1) Newborn/young-of-the year individuals are more commonly encountered there than in other areas. (2) Newborn/young-of-the-year individuals have a tendency to remain or return for extended periods. (3) Newborn/young-of-the-year individuals repeatedly use the area or habitat across years.
Beck et al., 2009	> an area in which juveniles occur at higher densities, > they avoid predation more successfully, > they grow faster compared to a different habitat, > the species moves into the area as larvae/neonates, accumulate biomass, and then move offshore, > contribution per unit area to the production of individuals that recruit to adult populations is greater, on average, than production from other habitats in which juveniles occur, from any combination of four factors, the ideal being an area having all 4 factors at high values: (1) density (2) growth (3) survival of juveniles (4) movement to habitats inhabited by adults
Martins et al., 2018	> Egg case nursery/Primary nursery area/For oviparous species: (1) high densities of eggs and egg cases in contact with the benthos or permanent structures (2) area must be used as an egg-laying area over multiple years by adults (3) newborns should leave the area promptly after hatching > Juvenile nursery area/Secondary nursery area: (1) should have a high abundance of neonates and juveniles, (2) may or may not be distinct from the egg case nursery (3) should strongly contribute to population recruitment > Nursery area for viviparous species (Heupel et al., 2007): (1) Newborn/young-of-the-year individuals are more commonly encountered there than in other areas. (2) Newborn/young-of-the-year individuals have a tendency to remain or return for extended periods. (3) Newborn/young-of-the-year individuals repeatedly use the area or habitat across years.

RESULTS AND DISCUSSION

Smooth-hound, *Mustelus mustelus* (Linnaeus, 1758)

A total of 138 *M. mustelus* specimens were caught as bycatch during the study period and included in the analysis. All individuals examined were juveniles, and no newborns with postnatal scars were reported. Juvenile females were more frequent than juvenile males, with a female-to-male ratio of 60:40 (Fig. 3). All specimens were caught over a 10-year span, specifically in September and October between 2011 and 2021. The average Fulton's condition factor across all specimens was 0.0032 (Tab. 3) with a standard error of ± 0.000221 , and females had a slightly higher mean than males (Tab. 4). Since all specimens were juveniles, no comparisons of condition factor between juveniles and adults could be made.

Blackspotted smooth-hound, *Mustelus punctulatus* (Risso, 1827)

Of the 175 *M. punctulatus* specimens included in the analysis, males were more frequent than females, with a sex ratio of 66:33 (Fig. 3). While no newborns were recorded, juvenile specimens of both sexes

accounted for 65.3% of all specimens. Both juvenile and adult specimens were reported from September to November, as well as in January and from April to June. The specimens included in this study were caught in 2002, 2003, 2011, and 2014, with juveniles reported in each of those years. The average Fulton's condition factor was 0.0031 (Tab. 3) with a standard error of ± 0.000302 , and males had a higher average than females (Tab. 4). Since biometric data were only available for juvenile specimens, no comparisons of Fulton's condition factors could be made between juvenile and adult specimens.

Marbled electric ray, *Torpedo marmorata* (Risso, 1810)

A total of 74 *T. marmorata* specimens were analyzed. The adult sex ration was almost 50:50, whereas juvenile males were more frequent than juvenile females, with a 3:1 ratio (Fig. 3). Three newborn males were recorded, and together with the other juveniles, they accounted for 55.4% of all specimens caught. *T. marmorata* was the most frequently caught species on a monthly basis. From September to March and in May, both adult and juvenile specimens were caught, with one newborn specimen caught in January and two in




Fig. 3: Maturity and sex distribution in adults and juveniles of the five species included in this study. Males (grey) and females (dark red) refer to adult individuals.

Sl. 3: Porazdelitev zrelosti in spola pri odraslih osebkih in mladostnih primerkih petih vrst, vključenih v to raziskavo. Samci (sivo) in samice (temno rdeče) se nanašajo na odrasle osebke.

Tab. 3: Fulton's condition factor values for the five species included in this study compared with values reported from other Mediterranean studies.

Tab. 3: Vrednosti Fultonovega kondicijskega faktorja za pet vrst, vključenih v to raziskavo, v primerjavi z vrednostmi iz drugih sredozemskih študij.

Species	Range	Average	Study
<i>M. mustelus</i>	0.0028–0.0036	0.032	this study
	0.002–0.003	/	Filiz & Mater, 2002
	0.003–0.004	/	Ismen et al., 2007
	/	0.004	Ozcan & Başusta, 2018
	0.003–0.01	/	Colombelli & Bonanomi, 2022
<i>M. punctulatus</i>	0.026–0.036	0.0031	this study
	/	0.003	Colombelli & Bonanomi, 2022
<i>T. marmorata</i>	0.0119–0.0870	0.0285	this study
	0.022–0.027	/	Filiz & Mater, 2002
	0.019–0.023	/	Ismen et al., 2007
	/	0.025	Duman & Başusta, 2013
<i>R. asterias</i>	0.0007–0.0068	0.0055	this study
	/	0.007	Karakulak et al., 2006
	/	0.007	Ferrà et al., 2016
<i>A. bovinus</i>	0.0022–0.0141	0.0052	this study
	0.012–0.022	/	Başusta et al., 2012
	/	0.004	Tsikliras & Dimarchopoulou, 2021
	0.014–0.021	/	Colombelli & Bonanomi, 2022

Tab. 4: Fulton's condition factor values for the five species included in this study by sex.**Tab. 4: Vrednosti Fultonovega kondicijskega faktorja po spolu za pet vrst, vključenih v to raziskavo.**

Species	Males		Females	
	Range	Average	Range	Average
<i>M. mustelus</i>	0.0028–0.0035	0.0031	0.0028–0.0036	0.0032
<i>M. punctulatus</i>	0.0026–0.0036	0.0032	0.0028–0.0034	0.0030
<i>T. marmorata</i>	0.0119–0.0870	0.0255	0.0168–0.0795	0.0337
<i>R. asterias</i>	0.0043–0.0065	0.0055	0.0007–0.0068	0.0057
<i>A. bovinus</i>	0.0022–0.0050	0.0031	0.0026–0.0141	0.0059

Tab. 5: Fulton's condition factor values for the five species included in this study by maturity stage.**Tab. 5: Vrednosti Fultonovega kondicijskega faktorja po stopnjah zrelosti za pet vrst, vključenih v to raziskavo.**

Species	Newborns		Juveniles		Adults	
	Range	Average	Range	Average	Range	Average
<i>M. mustelus</i>	/	/	0.0028–0.0036	0.0032	/	/
<i>M. punctulatus</i>	/	/	0.0026–0.0036	0.0031	/	/
<i>T. marmorata</i>	0.0204–0.0870	0.0380	0.0119–0.0795	0.0305	0.0168–0.0738	0.0252
<i>R. asterias</i>	/	/	0.0007–0.0068	0.0056	0.0048–0.0066	0.0055
<i>A. bovinus</i>	/	/	0.0022–0.0040	0.0030	0.0026–0.0141	0.0056

December. In April and August, only adult specimens were caught, while no specimens were caught in June or July. The *T. marmorata* specimens analyzed in this study were caught between 2011 and 2014, with both juveniles and adults reported in each year. The average Fulton's condition factor was 0.0285 (Tab. 3) with a standard error of ± 0.017578 . Females had a higher average than males (Tab. 4), and newborns and juveniles had a higher average than adults (Tab. 5).

**Mediterranean starry ray, *Raja asterias*
(Delaroche, 1809)**

Male *R. asterias* juveniles and adults outnumbered females, with a combined sex ratio of 4:1 (Fig. 3). No newborns were recorded; however, juveniles of both sexes accounted for 74.6% of all individuals.

Specimens were caught in April and from July to December during the period 2011–2016, with juveniles recorded in each year. The average Fulton's condition factor was 0.0055 (Tab. 3) with a standard error of ± 0.000908 . Females had a higher average than males (Tab. 4), and juveniles had a higher average than adults (Tab. 5).

**Bull ray, *Aetomylaeus bovinus*
(Geoffroy Saint-Hilaire, 1817)**

While *A. bovinus* was the only species in which juvenile specimens were less abundant than adults, juveniles still accounted for 40.54% of all caught individuals. Adult males and females occurred at an almost equal 1:1 frequency, while juvenile males outnumbered females by 2:1 (Fig. 3). Four male embryos were

found inside pregnant females; however, these were not included in the analysis. Both adult and juvenile specimens were reported from August to September 2005. The average Fulton's condition factor was 0.0052 (Tab. 3) with a standard error of ± 0.003550 . Females had a higher average than males (Tab. 4), and adults had a higher average than juveniles (Tab. 5).

Fulfillment of the three criteria defining nursery areas for the studied species

The first criterion – that juveniles are more frequently encountered in the area compared to other areas – was met by four of the five species. While catches of juveniles and adults of the examined species have been sporadically reported in the northern Adriatic, mostly along the Croatian coast, no solid nursery areas for any of the examined species have been confirmed yet. In the southern Adriatic, a nursery area for *R. asterias* has been proposed along the Albanian coast (Bakiu & Kule, 2024), however, it is unlikely that the *R. asterias* specimens caught in the northern Adriatic originated there.

For *M. mustelus*, all individuals recorded in the Gulf of Trieste during the study period were juveniles; therefore, the first criterion is considered fulfilled for this species. Given that no established nursery area for *M. mustelus* has been identified nearby and that the entire northwestern Adriatic has been classified as an ISRA important for the reproduction of this species (ISRA, 2025), it is possible that the Gulf of Trieste – as a confined part of the wider northwestern Adriatic area – plays an important role in its reproduction.

Juveniles also accounted for 65.3% of all *M. punctulatus* specimens, 55.4% of *T. marmorata* specimens, and 74.6% of *R. asterias* specimens; therefore, the first criterion is considered fulfilled for these species as well. In the Mediterranean, the closest suggested nursery area for *M. punctulatus* is in the Gulf of Gabes, southern Tunisia (Enajjar *et al.*, 2015), while for *R. asterias*, a nursery area has been identified within a restricted coastal zone of the southern Ligurian and northern Tyrrhenian Seas (Serena & Relini, 2006); therefore, the high proportion of juveniles of these two species recorded in the Gulf of Trieste is unlikely the result of a spillover from nearby nursery areas. For *T. marmorata*, both juvenile and adult specimens were recorded during the same period, suggesting that the species uses the Gulf of Trieste both as a reproductive habitat and a nursery area (pers. observation). This interpretation is further supported by observations from the shallow waters off the Sečovlje salina and within the salina channels, where numerous juvenile electric rays have been captured, sighted, photographed, and filmed (Kristina Gorišek, pers. comm.).

While juvenile specimens only accounted for 40.54% of all *A. bovinus* records, the occurrence of adults and pregnant females in the Gulf of Trieste during the same period suggests that reproduction may also take place in or near the area. Since *A. bovinus* is considered a thermophilic species, individuals may migrate seasonally into the Gulf of Trieste from more southern regions, potentially for foraging, given that all our records originate from the summer months. The closest potential nursery areas for *A. bovinus* in the Mediterranean are located along the southern Tunisian coast (Capapé and Quignard, 1975) and in the Amvrakikos Gulf in Greece (Ciprian *et al.*, 2026). The specimens recorded in the Gulf of Trieste may have originated from the latter area, given that the juvenile individuals for which TL/DW data were available were relatively large and close to the reported size at maturity for the species (87.5 cm for the male and 86.0 and 87.9 cm for the two females).

The second criterion – that juveniles remain in or return to the area for extended periods – was also fulfilled by four of the five species. Juvenile specimens were recorded over a period of several months, indicating that they are present not only following the reproductive period of adults, but also throughout the year. For all species except *A. bovinus*, both juvenile and adult specimens were recorded during periods corresponding to reproduction and immediately thereafter, also suggesting an increased abundance of juveniles shortly after the reproductive season.

Studies on the reproductive biology of *M. mustelus* place parturition in late spring (Saïdi *et al.*, 2008; Ozcan & Başusta, 2018; Boscolo Palo *et al.*, 2022). However, juveniles in the Gulf of Trieste were recorded in autumn, suggesting that they use the area independently of adult reproductive grounds, potentially as a secondary or juvenile nursery area. Accordingly, the second criterion can be considered met for this species.

Studies on *M. punctulatus* suggest that gestation begins in November, with parturition occurring from mid-May to early June (Saïdi *et al.*, 2009; Boscolo Palo *et al.*, 2022). Both juvenile and adult *M. punctulatus* individuals were reported in the Gulf of Trieste in autumn, winter, and spring, suggesting that the species may be using the basin as both a primary and a secondary nursery area. Therefore the second criterion is met for this species, too.

The *T. marmorata* records are in compliance with data from other studies on this species' reproductive biology in the Mediterranean Sea, where late pregnant individuals were found from July to October and post-pregnant individuals from September to February, as well as in May and August (Capapé, 1979; Consalvo *et al.*, 2007; Chatzisprou *et al.*, 2021).




Fig. 4: Annual bycatch (kg) of four of the five species included in this study, based on data from the Slovenian Fisheries Institute (BIOSWEB, 2025).

Fig. 4: Letni prilov (kg) za štiri od petih vrst, vključenih v to raziskavo, na podlagi podatkov Inštituta za ribištvo (BIOSWEB, 2025).

Tab. 6: Suitability of the abiotic conditions in the Gulf of Trieste for the five species included in this study.**Tab. 6: Primernost abiotskih dejavnikov (pogojev) v Tržaškem zalivu za pet vrst, vključenih v to raziskavo.**

	Substrate type	Climate	Salinity (psu)	Depth (m)
<i>M. mustelus</i>	sandy, muddy, or detritic bottoms, sometimes swim midwater	temperate	tolerates Mediterranean salinity level (38–40)	5–624 (usually 5–50)
<i>M. punctulatus</i>	inshore, sandy and muddy bottoms	subtropical	tolerates Mediterranean salinity level (38–40)	0–250
<i>T. marmorata</i>	seagrass areas, rocky reefs and adjacent soft bottoms	subtropical (< 20 °C)	tolerates Mediterranean salinity level (38–40)	2–270
<i>R. asterias</i>	benthic in inshore waters on sandy or muddy bottom	subtropical	tolerates Mediterranean salinity level (38–40)	2–700 (usually 20–50)
<i>A. bovinus</i>	demersal and semi-pelagic in estuaries, lagoons and on the continental shelf	subtropical	tolerates Mediterranean salinity level (38–40) as well as lower salinity	0–500
Gulf of Trieste	rocky coastal area, sandstone habitat types overgrown by algal vegetation, seagrass meadows, silty clay, biogenic detritus and soft bottom	temperate–humid subtropical (Cfa)	35–38	max. 37

These results indicate that the occurrence of juvenile, newborn, and adult *T. marmorata* in the Gulf of Trieste is typical of an area where a potential nursery might be located. Therefore, the second criterion is met for this species as well.

Research on the reproductive biology of *R. asterias* in the Mediterranean Sea suggests that juveniles are especially common from January to March and from July to September (Stehmann & Bürkel, 1984; Serena & Relini, 2006). Given that both juveniles and adults were caught in the Gulf of Trieste during and after the typical period when this species reproduces, this may indicate that the area is being used as a nursery. Therefore, the second criterion is also met for this species.

While not much information is available on the reproductive habits of *A. bovinus* in the Mediterranean Sea, the species' mating season generally begins in spring and ends in autumn (Michael, 1993). The reported catches of *A. bovinus* in the Gulf of Trieste align with this general pattern, as they occurred during the assumed reproductive period, with parturition probably occurring elsewhere. Since the juveniles recorded were close to the size at maturity reported for the species, they were likely born the previous year and migrated to the northern Adriatic from another area, possibly Amvrakikos Gulf, during the summer. This means

that both adults and juveniles may be transient in the area rather than using it as a nursery.

The third criterion – that juveniles use the area or habitat repeatedly over the years – was also fulfilled by four of the five species. Since no direct tracking data were available, bycatch records from this study, as well as data from the website of the Slovenian Fisheries Institute (ZZRS) reported for different years were used as a substitute. *M. mustelus* was by far the most abundantly caught species included in this study (Fig. 4), with catch records on the ZZRS website exceeding 1000 kg each year (BIOSWEB, 2025). Additionally, the specimens included in the analysis were caught 10 years apart. Therefore, the third criterion is met for this species.

The only catch records available for *M. punctulatus* span the years from 2005 to 2012 (Fig. 4), with no records thereafter (BIOS, 2025). It is possible that *M. punctulatus* and *M. mustelus* records have been conflated, as the two closely related species are known to be frequently misidentified (Marino et al., 2018). One distinguishing feature between the species is size at sexual maturity, estimated at 90 cm for males and 100 cm for females in *M. punctulatus*, compared with 70–80 cm for both sexes in *M. mustelus* (Lipej et al., 2011). Consequently, individuals representing adult *M. mustelus* may have been misidentified as

Tab. 7: Abundance of selected prey items of juveniles of all five species included in this study.

Tab. 7: Številčnost (abundanca) izbranega plena mladih primerkov vseh petih vrst, vključenih v to raziskavo.

Species	Food preference / specialization of juveniles	Abundance	Sources
<i>M. mustelus</i>	Crustaceans		Maynou <i>et al.</i> , 2004; Farrugio & Soldo, 2015; BIOSWEB, 2025
	<i>Squilla mantis</i>	seasonally abundant (summer)	
	<i>Maja squinado</i>	no data	
<i>M. punctulatus</i>	Crustaceans		Maynou <i>et al.</i> , 2004; Farrugio & Soldo, 2015; Bacci <i>et al.</i> , 2024
	<i>Squilla mantis</i>	seasonally abundant (summer)	
	<i>Ethusa mascarone</i>	present	
<i>T. marmorata</i>	Teleost fish		Farrugio & Soldo, 2014; 2015; BIOSWEB, 2025
	<i>Boops boops</i>	seasonally abundant (summer)	
	<i>Spicara</i> spp.	no data	
	Soleidae	seasonally abundant (fall)	
	Crustaceans		
	<i>Cypridina mediterranea</i>	no data	
	Cephalopods		
	<i>Loligo vulgaris</i>	seasonally abundant (winter)	
	<i>Sepia elegans</i>	seasonally abundant (summer)	
<i>R. asterias</i>	Crustaceans		Farrugio & Soldo, 2014; 2015; Bacci <i>et al.</i> , 2024
	<i>Ethusa mascarone</i>	present	
	<i>Polybius</i> sp.	present	
	Cephalopods		
	<i>Loligo vulgaris</i>	seasonally abundant (winter)	
	<i>Sepia</i> sp.	seasonally abundant (summer)	
	Teleost fish		
	<i>Gobius niger</i>	present	
	<i>Cepola macrophthalma</i>	no data	
<i>A. bovinus</i>	Gastropods		Bacci <i>et al.</i> , 2024
	<i>Gibbula magus</i>	no data	
	Crustaceans		
	<i>Paguristes eremita</i>	present	

juvenile *M. punctulatus*, which could explain why only juveniles of *M. mustelus* have been recorded throughout the study period. The specimens analyzed in this study were caught in 2002, 2003, 2011, and 2014, suggesting that not all catches may be represented in the ZZRS database. As these records span several years, the third criterion is considered met for *M. punctulatus*, too.

There are virtually no catch records of *T. marmorata* on the ZZRS website before 2016, after which irregular annual records are reported, ranging from 30 to 283 kg (Fig. 4). The specimens analyzed in this study were caught between 2011 and 2014, suggesting that not all historical catches are represented in the ZZRS database. Given that juveniles were caught in multiple years, the third criterion is met for this species as well.

R. asterias has been reported every year since 2000, except for 2003 and 2004; however, it should be noted that similar reporting gaps are present for several other species during the mentioned two years, suggesting that the pattern may not accurately reflect landings of *R. asterias* for that period (BIOSWEB, 2025). The *R. asterias* specimens included in this study were caught between 2011 and 2016, with juveniles captured in each year. Therefore, the third criterion is also met for this species.

For *A. bovinus*, no assumptions can be made regarding whether juveniles were recorded across multiple years. Two studies included specimens caught in 2005 (Dulčić et al., 2008; Lipej et al., 2009), while Uhan (2016) reported two adult specimens added to the collection of the Marine Biology Station of the National Institute of Biology in 2014. However, since the ZZRS website does not provide catch records for this species, no conclusion can be drawn regarding the third criterion based on the available data.

Fulton's condition factor and additional criteria

Combining Fulton's condition factor values with the biotic and abiotic conditions of the Gulf of Trieste allowed for an extension of the analysis to ecological factors that could not be assessed through the main nursery area criteria alone.

The mean Fulton's condition factor for *M. mustelus* (Tab. 3) was well within the expected range for the species, as both lower and higher values have been reported in the Mediterranean Sea (Filiz & Mater, 2002; Ismen et al., 2007; Ozcan & Başusta, 2018; Colombelli & Bonanomi, 2022). However, only juvenile specimens had available biometric data, therefore the condition factor values could not be directly compared between juveniles and adults.

As with *M. mustelus*, only juvenile specimens of *M. punctulatus* had available biometry; therefore, condition factor values in juveniles could not be directly compared with those in adults from this study (Tab. 5). However, when compared to similar studies from the region, the mean value (Tab. 3) is slightly higher than those reported for the Mediterranean (Colombelli & Bonanomi, 2022).

For *T. marmorata*, comparable studies in the Mediterranean have reported lower mean values for Fulton's condition factor (Filiz & Mater, 2002; Ismen et al., 2007; Duman & Başusta, 2013); accordingly, the specimens in this study appear to exhibit above-average mean condition factor values (Tab. 3).

Fulton's condition factor mean for *R. asterias* was lower than values reported by comparable studies from the northern Adriatic, which also included a higher proportion of juveniles than adults (Karakulak et al., 2006; Ferrà et al., 2016). It should be noted that some of the specimens in Ferrà et al. (2016) originated from the Gulf of Trieste, and the authors recognized the area as an aggregation site for the species (Ferrà et al., 2016).

The mean Fulton's condition factor for *A. bovinus* was within the range reported in comparable studies from the Mediterranean (Başusta et al., 2012; Tsikliras & Dimarchopoulou, 2021; Colombelli & Bonanomi, 2022).

Analyses of the stomach contents of *M. mustelus* suggest that it is an opportunistic predator, feeding on a wide range of prey items, including benthic invertebrates and fishes, as well as prey of varying sizes and morphologies (Ahmed et al., 2022). In the northern Adriatic, decapod crustaceans are considered the most important and frequent prey for juveniles and are present in all seasons (Jardas et al., 2007). This indicates that prey resources for juveniles are likely available throughout the year, supporting the suitability of the Gulf of Trieste as a potential nursery area.

Juveniles of *M. punctulatus* in the northern Adriatic have also been shown to mainly feed on crustaceans – unlike adults, which preferentially consume cephalopods (Lipej et al., 2011) – however, there is no clear indication of competition for food resources with *M. mustelus* juveniles (Tab. 6 and Tab. 7), suggesting that the two species may co-occur within the same nursery area.

Analyses of the diet of *T. marmorata* in the Mediterranean show that both juveniles and adults primarily feed on teleost fishes (Chatzisprou et al., 2021) and cephalopods (Capapé et al., 2007). The species has also shown local dietary specialization (Chatzisprou et al., 2021). This ability might be especially useful in communal nursery areas to avoid competition for food with juveniles of other species.

Diet analyses of *R. asterias* in the Adriatic Sea show that both juveniles and adults primarily feed on decapods, with cephalopods and teleost fishes occurring in smaller proportions (Sviben *et al.*, 2019). Sviben *et al.* (2019) further report that specimens from the Slovene sector of the Gulf of Trieste exhibited the highest average prey weight, prey diversity, and meal value, suggesting plentiful food resources for the species. This may be particularly relevant in the context of nursery function, especially in a multi-species setting where juveniles of other taxa also primarily consume decapods.

The diet of juvenile *A. bovinus* in the northern Adriatic consists almost exclusively of gastropods, while adults also consume fish (Lipej *et al.*, 2025). The disk width (DW) at birth of specimens from the northern Adriatic was greater than that observed in specimens from other parts of the Mediterranean. Correspondingly, adult individuals from this region attained some of the highest body weights recorded across the species distribution range (Dulčić *et al.*, 2008; Lipej *et al.*, 2009). Dulčić *et al.* (2008) and Lipej *et al.* (2009) suggested that these differences may be related to the apparent absence of parasitic infections in the examined specimens. If so, low parasite prevalence may be another habitat characteristic that favors the presence of this species (Tab. 6 and Tab. 7).

Based on these results, the Gulf of Trieste is most likely used as a communal juvenile habitat (secondary nursery area) by four of the five species examined, as it fulfills the three criteria for a nursery area for all species except *A. bovinus*. It also satisfies the first criterion of a juvenile nursery area as defined by Martins *et al.* (2018) (Tab. 2). However, due to the limited number of specimens and lack of tracking data, it is difficult to assess the contribution of the area to juvenile recruitment. The abiotic (Tab. 6) and biotic (Tab. 7) conditions in the Gulf of Trieste are consistent with those expected of a nursery area for all five species: the gulf is relatively shallow, features both soft-sediment and hard-substrate habitats, and contains abundant prey resources for each species. Some species have been shown to exhibit dietary flexibility in habitats where they overlap with other species in order to avoid intra- and inter-specific competition, which can also depend on their mode

of reproduction (Capapé *et al.*, 2003; Chatzispyrou *et al.*, 2021; Ahmed *et al.*, 2022; Finotto *et al.*, 2023). Additionally, primary production measurements suggest that the trophic conditions in the area have not significantly changed in over a decade (Cibic *et al.*, 2022) and have therefore been stable enough to potentially aid the development of a nursery area. None of the five species have any natural predators in the area, and they all occupy high trophic levels typical of predators at the top of the food chain (Lipej *et al.*, 2011; Sviben *et al.*, 2019; Ahmed *et al.*, 2022; Lipej *et al.*, 2025).

The characteristics of the Gulf of Trieste indicate that the habitat could support a communal nursery area for all five species. None of the species in this study are subject to protection in the region; in Slovenia, specifically, there are no ongoing regular monitoring programs regarding sharks and rays (Lipej *et al.*, 2024). Despite the presence of several suitable sites that could serve as potential nursery areas and already have some form of protection – both in Slovenia (the natural monuments Cape Madonna and Debeli rtič, the Strunjan Nature Reserve, the Sečovelje Salina Nature Park, and Landscape Park Strunjan) and in Italy (the Miramare Marine Protected Area, Duino Cliffs Nature Reserve, and Cavana di Monfalcone) – none have yet been designated or evaluated as potential nursery areas (Lipej *et al.*, 2024; WDPa, 2025). Additionally, juveniles of other elasmobranch species, such as *Carcharhinus plumbeus*, *Alopias vulpinus*, *Squalus acanthias*, *Raja clavata*, *Dasyatis pastinaca*, *Pteroplatytrigon violacea*, and *Myliobatis aquila*, have also been documented in the Gulf of Trieste over the 24-year period (Fig. 4). However, they were not included in this study as there were insufficient specimens of each species available for further analysis (Lipej *et al.*, 2008; 2016; 2022a; 2022b; 2023). Regardless, this highlights the potential importance of the area as a nursery for an even higher number of species than those considered in this study. Given the limited available data on elasmobranchs in the region, increased efforts to gather more information and conduct accurate future analyses are needed. A proper assessment of the real status of elasmobranch species in the area is essential for informing effective conservation and management measures.

ALI JE TRŽAŠKI ZALIV POTENCIALNO OBMOČJE ODRAŠČANJA (JASLICE) ZA DOLOČENE VRSTE HRUSTANČNIC?

Hristina GELEVSKA

FAMNIT, University of Primorska, Koper, Slovenia
e-mail: hgelevska@gmail.com

Borut MAVRIČ & Lovrenc LIPEJ

Marine Biology Station Piran, National Institute of Biology, Piran, Slovenia

Christian CAPAPÉ

Laboratoire d'Ichtyologie, Université de Montpellier, 34 095 Montpellier cedex 5, France

POVZETEK

Avtorji poročajo o izsledkih raziskave, katere namen je bil ugotoviti, ali Tržaški zaliv izpolnjuje merila za območje odraščanja, kot jih definirajo različni viri, predvsem Heupel s sodelavci (2007). Analiza je temeljila na osebkih hrustančnic iz zbirke Morske biološke postaje Piran (Nacionalni inštitut za biologijo, Slovenija) in je vključevala primerke petih vrst (*Mustelus mustelus*, *M. punctulatus*, *Torpedo marmorata*, *Raja asterias* in *Aetomylaeus bovinus*), ujetih kot prilov v obdobju 24 let. Štiri od petih vrst (razen vrste *A. bovinus*) so izpolnile vsa tri merila. Glede na analizirane podatke bi Tržaški zaliv lahko obravnavali kot skupno območje odraščanja mladice za preučevane vrste. Poleg tega zapisi o mladih primerkih drugih vrst hrustančnic, o katerih so poročali v obdobju raziskave na tem območju, kažejo na potencialen pomen tega okolja kot območja odraščanja tudi za druge vrste.

Ključne besede: hrustančnice, skotitev, merila za območje odraščanja, abiotski pogoji, prilov, severni Jadran

REFERENCES

- Ahmed, A.M.M., A. Azab, M.A. Ahmad, H.M.M. Khalaf-Allah, & M.A.M. El-Tabakh (2022): Observations on food and feeding habits of the common smooth hound shark, *Mustelus mustelus* in the Mediterranean Sea at Alexandria coast, Egypt. Egypt. J. Aquat. Biol. Fish., 26(3), 123–138.
- Akyol, O., İ. Aydin, E. Olfa & C. Capapé (2017): Bull ray, *Aetomylaeus bovinus* (Geoffroy Saint-Hilaire, 1817) (Myliobatidae) in the Mediterranean Sea and captures of juveniles from Izmir Bay (Aegean Sea, Turkey). J. Appl. Ichthyol., 33, 1200–1203.
- Akyol, O., H. Şen & C. Capapé (2022): Occurrence of a Shoal of Bull Ray *Aetomylaeus bovinus* (Myliobatidae) Around a Sea-Cage Farm in İskenderun Bay (Türkiye, NE Mediterranean Sea). COMU-JMSF, 5(2), 199–202.
- Bacci, T., M. Targusi, L. Lattanzi, V. Marusso, B. Trabucco & P. Tomassetti (2024): Decapod Crustacean Records from A Collection of Marine Soft-Bottom Macrofauna of the Italian Waters Supporting the Update of Checklists and Registers of the Mediterranean Basin. Diversity, 16(1), 2.
- Bajt, O., B. Mavrič, R. Milačič, J. Ščančar, T. Zuliani & L. Lipej (2024): Bioaccumulation of organotin compounds in the marbled electric ray (*Torpedo marmorata*) in the northern Adriatic Sea. Mar. Pollut. Bull., 204, 116511.
- Bakiu, R. & M. Kule (2024): Critical life stage and spatial analyses of shark and ray species in the Albanian territorial waters. Studia Marina, 37(2), 14–23.
- Bass, A.J. (1978): Problems in studies of sharks in the Southwest Indian Ocean. In: Hodgson ES, Mathewson RF (eds) Sensory biology of sharks, skates and rays. Office of Naval Research, Department of the Navy, Arlington, VA, pp. 545–594.
- Başusta, A., N. Başusta, J.A. Sulikowski, W.B. Driggers III., S.A. Demirhan & E. Çiçek (2012): Length–weight relationships for nine species of batoids from the Iskenderun Bay, Turkey. J. Appl. Ichthyol., 28, 850–851.
- Beck, M.W., K.L. Heck Jr., K.W. Able, D.L. Childers, D.B. Eggleston, B.M. Gillanders, B. Halpern, C.G. Hays, K. Hoshino, T.J. Minello, R.J. Orth, P.F. Sheridan & M.P. Weinstein (2009): The Identification, Conservation, and Management of Estuarine and Marine Nurseries for Fish and Invertebrates. BioScience, 51, 633–641.
- BIOSWEB (2025): Biološka Zbirka Podatkov Zavoda za Ribištva Slovenije (date of access: 01.04.2025) <http://www.biosweb.org/index.php?task=taxonsearch>
- Boscolo Palo, G., M. Di Lorenzo, G. Salvatore, S. Ragonese, C. Mazzoldi & F. Colloca (2022): Sharks Do Not Always Grow Slowly: Tagging Data Reveal a Different Pattern of Growth, Longevity and Maturity for Threatened Smooth-Hounds in the Central Mediterranean Sea. J. Mar. Sci. Eng., 10, 1647.
- Bradaï, M.N., B. Saïdi, A. Bouaïn, O. Guélorget & C. Capapé (2005): The Gulf of Gabès (southern Tunisia, central Mediterranean): nursery area for sandbar shark, *Carcharhinus plumbeus* (Nardo, 1827) (Chondrichthyes: Carcharhinidae). Ann. Ser. Hist. Nat., 15 (2), 187–194.
- Capapé, C. (1979): La torpille marbrée, *Torpedo marmorata* Risso, 1801 (Pisces, Rajiformes) des côtes tunisiennes: nouvelles données sur l'écologie et la biologie de la reproduction de l'espèce avec une comparaison entre les populations méditerranéennes et atlantiques. Ann. Sci. nat., Zool., Paris, 13 (1), 79–97.
- Capapé, C., O. Guélorget, C. Reynaud, A. Marquès, J.L. Bouchereau & J. Zaouali (2003): Effects of reproductive factors on interrelationships among three deep water sharks from northern Tunisia (Central Mediterranean). Annales, Ser. Hist. Nat., 13(2), 109–120.
- Capapé, C., S. Crouzet, C. Clement, Y. Vergne & O. Guelorget (2007): Diet of the marbled electric ray *Torpedo marmorata* (Chondrichthyes: Torpedinidae) off the Languedocian coast (Southern France, Northern Mediterranean). Annales, Ser. Hist. Nat., 17(1), 17–22.
- Capapé, C. & J.P. Quignard (1975): Contribution à la systématique et à la biologie de *Pteromylaeus bovinus* (Geoffroy Saint-Hilaire, 1817), (Pisces, Myliobatidae) des côtes tunisiennes. Bull. Mus. Hist. Nat., Paris, 3 (338) Zool., 240, 1329–1347.
- Cashion, M. S., Bailly, N., & D. Pauly (2019): Official catch data underrepresent shark and ray taxa caught in Mediterranean and Black Sea fisheries. Marine Policy, 105, 1–9.
- Catalano, G., V. Crobe, A. Ferrari, R. Baino, D. Massi, A. Titone, C. Mancusi, F. Serena, R. Cannas, L. Carugati, F. Hemida, C. Manfredi, R. Melis, G. Scarcella, L. Sion, M. Stagioni, F. Tinti & A. Cariani (2021): Strongly structured populations and reproductive habitat fragmentation increase the vulnerability of the Mediterranean starry ray *Raja asterias* (Elasmobranchii, Rajidae). Aquat. Conserv., 32, 10.1002/aqc.3739.
- Chatzisprou, A., C. Gubili, K. Touloumis, D. Karampetsis, S. Kioulouris, A. Anastasopoulou, A. Christidis, P. Peristeraki, I.E. Batjakas & C. Koutsikopoulos (2021): Life-history traits of the marbled electric ray, *Torpedo marmorata* Risso, 1810, from the Greek Seas, north-eastern Mediterranean Sea. Environ. Biol. Fishes, 10, 1575–1592.
- Ciprian, M., I. Giovos, C. Mazzoldi, & D. K. Moutopoulos (2026). Morphometric Relations Within Elasmobranch Species from the Amvrakikos Gulf (Central Mediterranean). Diversity, 18(1), 41.

- Cibic, T., L. Baldassarre, F. Cerino, C. Comici, D. Fornasaro, M. Kralj & M. Giani (2022):** Benthic and Pelagic Contributions to Primary Production: Experimental Insights from the Gulf of Trieste (Northern Adriatic Sea). *Front. Mar. Sci.*, 9, 877935.
- Coll, M., J. Navarro & I. Palomera (2013):** Ecological role, fishing impact, and management options for the recovery of a Mediterranean endemic skate by means of food web models. *Biol. Conserv.*, 157, 108–120.
- Colombelli, A. & S. Bonanomi (2022):** Length–weight relationships for six elasmobranch species from the Adriatic Sea. *J. Appl. Ichthyol.*, 38, 328–332.
- Compagno, L.J.V. (1984):** FAO species catalogue vol. 4. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Part 2. Carcharhiniformes. FAO Fisheries Synopsis 125 (4/2), Rome, pp. 251–655.
- Consalvo, I., U. Scacco, M. Romanelli & M. Vacchi (2007):** Comparative study on the reproductive biology of *Torpedo torpedo* (Linnaeus, 1758) and *T. marmorata* (Risso, 1810) in the central Mediterranean Sea. *Sci. Mar.*, 71(2), 213–222.
- Conversi, A., T. Peluso & S. Fonda-Umani (2009):** Gulf of Trieste: A changing ecosystem. *J. Geophys. Res.*, 114, C03S90, doi:10.1029/2008JC004763.
- Dulčić, J., L. Lipej, M. Orlando-Bonaca, R. Jenko, B. Grbec, O. Guélorget & C. Capapé (2008):** The bull ray, *Pteromylaeus bovinus* (Myliobatidae), in the northern Adriatic Sea. *Cybiu*, 32, 119–123.
- Dulvy, N.K., S.L. Fowler, J.A. Musick, R.D. Cavanagh, P.M. Kyne, L.R. Harrison, J.K. Carlson, L.N.K. Davidson, S.V. Fordham, M.P. Francis, C.M. Pollock, C.A. Simpfendorfer, G.H. Burgess, K.E. Carpenter, L.J.V. Compagno, D.A. Ebert, C. Gibson, M.R. Heupel, S.R. Livingstone, J.C. Sanciangco, J.D. Stevens, S. Valenti & W.T. White (2014):** Extinction risk and conservation of the world's sharks and rays. *eLife*, 3, e00590.
- Duman, O. & N. Başusta (2013):** Age and growth characteristics of marbled electric ray *Torpedo marmorata* (Risso, 1810) Inhabiting Iskenderun Bay, North-eastern Mediterranean Sea. *Turk. J. Fish. Aquat. Sci.*, 13, 551–559.
- Enajjar, S., B. Saïdi & M. Bradai (2015):** The Gulf of Gabès (Central Mediterranean Sea): a nursery area for sharks and batoids (Chondrichthyes: Elasmobranchs). *Cah. Biol. Mar.*, 56, 143–150.
- Falace, A., G. Alongi, M. Orlando-Bonaca & S. Bevilacqua (2024):** Species loss and decline in taxonomic diversity of macroalgae in the Gulf of Trieste (Northern Adriatic sea) over the last six decades. *Mar. Environ. Res.*, 202, 106828.
- Farrugio, H. & A. Soldo (2014):** Status and Conservation of Fisheries in the Adriatic Sea. Draft internal report for the purposes of the Mediterranean Regional Workshop to Facilitate the Description of Ecologically or Biologically Significant Marine Areas, UNEP/MAP-RAC/SPA, Malaga, 50 pp.
- Farrugio, H. & A. Soldo (2015):** Adriatic Sea: Status and conservation of fisheries. Cebrian, D. & S. Requena (eds.). UNEP/MAP-RAC/SPA, Tunis; 58 pp.
- Farrugio H., P. Oliver & F. Biagi (1993):** An overview of the history, knowledge, recent and future research trends in Mediterranean fisheries. *Sci. Mar.*, 57, 105–119.
- Ferrà, C., G. Fabi, P. Polidori, A.N. Tassetti, S. Leoni, G. Pellini & G. Scarcella (2016):** *Raja asterias* population assessment in FAO GFCM GSA17 area. *Mediterr. Mar. Sci.*, 17(3), 651–660.
- Filiz, H. & S. Mater (2002):** A preliminary study on length–weight relationships for seven Elasmobranch species from North Aegean Sea, Turkey. *EgeJFAS*, 19 (3-4), 401–409.
- Finotto, L., D. Berto, F. Rampazzo, S. Raicevich, S. Bonanomi & C. Mazzoldi (2023):** Trophic partitioning among three mesopredatory shark species inhabiting the Northwestern Adriatic Sea. *Diversity*, 15, 1163.
- Follesa, M.C., M.F. Marongiu, W. Zupa, A. Bellodi, A. Cau, R. Cannas, F. Colloca, M. Djurovic, I. Isajlovic, A. Jadaud, C. Manfredi, A. Mulas, P. Peristeraki, C. Porcu, S. Ramirez-Amaro, F. Salmerón Jiménez, F. Serena, L. Sion, I. Thasitis, A. Cau, P. & P. Carbonara (2019):** Spatial variability of Chondrichthyes in the northern Mediterranean. *Sci. Mar.*, 83(1), 81–100.
- Froese, R. & D. Pauly (eds.) (2024):** FishBase. Available at: www.fishbase.org. (date of access: 04.04.2025).
- Fulton, T. W. (1902):** Rate of growth of sea fishes. In: 20th Annual report of the Fishery Board of Scotland for the year ended 1902 (ed.). Neill & Company, Edinburgh, pp. 326–446.
- Goosen, A.J.J. & M.J. Smale (1997):** A preliminary study of age and growth of the smoothhound shark *Mustelus mustelus* (Triakidae). *S. Afr. J. Mar. Sci.*, 18(1), 85–91.
- Gračan, R., T. Polak & B. Lazar (2021):** Life history traits of the Blackspotted smooth-hound *Mustelus punctulatus* (Carcharhiniformes: Triakidae) in the Adriatic Sea. *Nat. Croat.*, 30, 475–492.
- Gulf of Trieste - Regional climate model data from EURO-CORDEX for the eLTER project** (date of access: 21.05.2025) <https://deims.org/f2ce5ae3-8873-4a8b-abad-d56d5d6da164>.
- Heithaus, M. (2007):** Nursery areas as essential shark habitats: A theoretical perspective. In: Shark Nursery Grounds of the Gulf of Mexico and the East Coast Waters of the United States. AFS, pp. 3–13.
- Heupel, M., J.K. Carlson & C. Simpfendorfer (2007):** Shark nursery areas: Concepts, definition, characterization and assumptions. *Mar. Ecol. Prog. Ser.*, 337, 287–297.
- İşmen, A., O. Ozen, U. Altınağaç, U. Ozekinci & A. Ayaz (2007):** Weight–length relationships of 63 fish species in Saros Bay, Turkey. *J. Appl. Ichthyol.*, 23, 707–708.

- IUCN and UNEP-WCMC (2025):** The World Database on Protected Areas (WDPA) [On-line], [8/2025], UNEP-WCMC, Cambridge. Available at: www.protectedplanet.net.
- Jardas, I., M. Šantić, V. Nerlović & A. Pallaoro (2007):** Diet of the smooth-hound *Mustelus mustelus* (Chondrichthyes: Triakidae) in the eastern Adriatic Sea. *Cybius*, 31(4), 459–464.
- Karakulak, F. S., H. Erk, H. & B. Bilgin (2006):** Length–weight relationships for 47 coastal fish species from the northern Aegean Sea, Turkey. *J. Appl. Ichthyol.*, 22, 274–278.
- Lipej, L. (ed.) (2021):** Slovenia Conservation of Mediterranean Marine and Coastal Biodiversity by 2030 and Beyond. UNEP/MAP-SPA/RAC, Tunis, 89 pp.
- Lipej, L., B. Mavrič & C. Capapé (2008):** On the occurrence of the sandbar shark, *Carcharhinus plumbeus* (Chondrichthyes: Carcharhinidae) off the Slovenian coast (northern Adriatic). *Acta Adriat.* 49(2), 137–145.
- Lipej, L., B. Mavrič & J. Dulčić (2009):** Size of the bull ray, *Pteromylaeus bovinus* (Geoffroy Saint-Hilaire, 1817), from the northern Adriatic. *J. Appl. Ichthyol.*, 25, 103105.
- Lipej, L., B. Mavrič, S. Rešek, M. Chérif & C. Capapé (2011):** Food and feeding habits of the blackspotted smooth-hound, *Mustelus punctulatus* (Elasmobranchii: Carcharhiniformes: Triakidae), from the northern Adriatic. *Acta Ichthyol. Piscat.*, 41(3), 171–177.
- Lipej, L., J. Uhan, B. Mavrič & S. Vujčić-Karlo (2016):** A record of porbeagle, *Lamna nasus* (Bonnaterre, 1788), in the Gulf of Trieste with discussion on its occurrence in the Adriatic Sea. *Acta Adriat.*, 57(2), 305–314.
- Lipej, L. & B. Mavrič (2022):** An unusual record of the sharpnose sevengill shark, *Heptranchias perlo* (Bonnaterre, 1788), in waters of Slovenia (northern Adriatic Sea). *Spixiana*, 45(1), 103–108.
- Lipej, L., D. Trkov, B. Mavrič, T. Fortibuoni, N. Bettoso, D. Donša & D. Ivajnsiĉ (2022):** Occurrence of bluntnose sixgill shark, *Hexanchus griseus* (Bonnaterre, 1788) in the Gulf of Trieste (northern Adriatic) with particular reference to historical and contemporary records in the Adriatic Sea. *Acta Adriat.*, 63(1), 93–108.
- Lipej, L., L.L. Zamuda & B. Mavrič (2023):** Note on a juvenile common Thresher shark *Alopias vulpinus* (Bonnaterre 1788) in the waters off Slovenia (Northern Adriatic Sea). *Croat. J. Fish.*, 81, 183–187.
- Lipej, L., B. Mavrič & A. Bolje (2024):** Chapter 7: Mediterranean and Black Seas Region. In: Jabardo, R. W., A. Z. A. Morata, R. H. Benett, B. Finucci, J. R. Ellis, S. Fowler, M. I Grant, A. P. Barbosa Martins, S. L. Sinclair (eds.): *The Global Status of Sharks, Rays and Chimaeras*. IUCN, Gland, pp. 138–143.
- Lipej, L., R. Battistella, B. Mavrič & D. Ivajnsiĉ (2025):** An insight into the diet of the Bull ray, *Aetomylaeus bovinus* (Geoffroy Saint-Hilaire, 1817) in the Northern Adriatic Sea. *Ann. Ser. Hist. Nat.*, 35(1), 43–54.
- Malaĉiĉ, V. & B. Petelin (2009):** Climatic circulation in the Gulf of Trieste (northern Adriatic), *J. Geophys. Res.*, 114, C07002.
- Marino, I., L. Finotto, F. Colloca, M. Di Lorenzo, M. Gristina, E. D. Farrell, L. Zane & C. Mazzoldi (2018):** Resolving the ambiguities in the identification of two smooth-hound sharks (*Mustelus mustelus* and *Mustelus punctulatus*) using genetics and morphology. *Mar. Biodivers.* 48(3), 1551–1562.
- Martins, A.P.B., M.R. Heupel, A. Chin & C.A. Simpfendorfer (2018):** Batoid nurseries: definition, use and importance. *Mar. Ecol. Prog. Ser.*, 595, 253–267.
- Maynou, F., P. Abello & P. Sartor (2004):** A review of the fisheries biology of the mantis shrimp, *Squilla mantis* (L., 1758) (Stomatopoda, Squillidae) in the Mediterranean. *Crustaceana*, 77, 1081–1099.
- Michael, S.W. (1993):** Reef Sharks and Rays of the World: A Guide to their Identification, Behavior, and Ecology. MD: Lighthouse Press, Annapolis, 112 pp.
- Mendez, L., A. Bacquet & F. Briand (2022):** Guide of Mediterranean Skates and Rays. Available at: www.ciesm.org/Guide/skatesandrays/.
- Monti, M., M. Minocci, L. Milani, S. Fonda Umani (2012):** Seasonal and interannual dynamics of microzooplankton abundances in the Gulf of Trieste (Northern Adriatic Sea, Italy). *Estuar. Coast. Shelf Sci.*, 115, 149–157.
- Muñoz-Chapuli, R. (1984):** Ethologie de la reproduction chez quelques requins de l'Atlantique nord-est. *Cybius*, 8(3), 1–14.
- Mytilineou, C., C.Y. Politou, C. Papaconstantinou, S. Kavadas, G. D'Onglia & L. G. Sion (2005):** Deep-water fish fauna in the Eastern Ionian Sea. *Belg. J. Zool.*, 135(2), 229–233.
- Ozcan, E.I. & N. Bařusta (2018):** Preliminary study on age, growth and reproduction of *Mustelus mustelus* (Elasmobranchii: Carcharhiniformes: Triakidae) inhabiting the Gulf of Iskenderun, north-eastern Mediterranean Sea. *Acta Ichthyol. Piscat.*, 48, 27–36.
- Pough, F.H. & C.M. Janis (2019):** Extant Chondrichthyans. In: Pough, F. H., W. E. Bemis, B. McGuire & C.M. Janis (eds.): *Vertebrate Life* (10th Edition). Oxford University Press, pp. 103–119.
- Ragheb, E. (2023):** Length-weight relationship and well-being factors of 33 fish species caught by gillnets from the Egyptian Mediterranean waters off Alexandria. *J. Aquat. Res.*, 49 (3), 361–367.

Rosa, R., E. Nunes, V. Pissarra, C.P. Santos, J. Varela, M. Baptista, J. Castro, J.R. Paula, T. Repolho, T.A. Marques, R. Freitas & C. Frazão (2023): Evidence for the first multi-species shark nursery area in Atlantic Africa (Boa Vista Island, Cabo Verde). *Front. Mar. Sci.*, 10, 2296–7745.

Saïdi, B., M.N. Bradaï & A. Bouaïn (2009): Reproductive biology and diet of *Mustelus punctulatus* (Risso, 1826) (Chondrichthyes: Triakidae) from the Gulf of Gabès, central Mediterranean Sea. *Sci. Mar.*, 73(2), 249–258.

Saïdi, B., M.N. Bradaï & A. Bouaïn (2008): Reproductive biology of the smooth-hound shark *Mustelus mustelus* (L.) in the Gulf of Gabès (south-central Mediterranean Sea). *J. Fish Biol.*, 72, 1343–1354.

Serena, F. & G. Relini (2006): Use of Scientific Campaigns (Trawl Surveys) for the Knowledge of the Sensitive Habitats. A Review of the MEDITS, GRUND and APHIA Data with Special Attention to the Italian Seas. In: N. Baştusta, Ç. Keskin, F. Serena, S. Bernard (eds.): International workshop on Cartilaginous fishes in the Mediterranean. Turkish Marine Research Foundation, Istanbul, pp. 135–148.

Soldo, A. & L. Lipej (2022): An annotated checklist and the conservation status of Chondrichthyans in the Adriatic. *Fishes*, 7, 245.

Stehmann, M. & D.L. Bürkel (1984): Rajidae. In: Whitehead, P.J.P., M.L. Bauchot, J.C. Hureau, J. Nielsen, E. Tortonese (eds.): *Fishes of the north-eastern Atlantic and Mediterranean*. UNESCO, Paris. Vol. 1, pp. 163–196.

Sviben, S., D. Trkov, B. Mavrič, P. Kružić & L. Lipej (2019): Observations on the diet of the Starry skate, *Raja asterias* Delaroche, 1809 (Elasmobranchii: Rajidae) in the Adriatic Sea. *Mediterr. Mar. Sci.*, 20(3), 577–584.

The International Union of Conservation of Nature (IUCN)(2025): Red List of Threatened Species (date of access: 01.04.2025) <https://www.iucnredlist.org/>.

The WEB-GIS Important Shark and Ray Areas (ISRA) (2025): Atlas - International Union of Conservation of Nature (IUCN) (date of access: 29.03.2025) <https://sharkrayareas.org/e-atlas/>.

Tserpes, G., C.D. Maravelias, M. Pantazi & P. Peristeraki (2013): Distribution of relatively rare demersal elasmobranchs in the eastern Mediterranean. *Estuar., Coast. Shelf Sci.*, 117, 48–53.

Tsikliras, A.C. & D. Dimarchopoulou (2021): Filling in knowledge gaps: Length–weight relations of 46 uncommon sharks and rays (Elasmobranchii) in the Mediterranean Sea. *Acta Ichthyol. Piscat.*, 51(3), 249–255.

Uhan, J. (2016): Biometrične značilnosti hrustančnic (Elasmobranchii) iz severnega Jadranskega morja. Diplomsko naloga, University of Ljubljana, Ljubljana, 71 pp.